

A TRIMEC(R) COMPLEX
ONE QUART COVERS 8,000 SQ. FT.

001
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under 2217-695
EPA Reg. No.

- KEEP FROM FREEZING -

KILLS: DANDELION, CRICKWEED, KNOTWEED, PLANTAIN, HENBIT,
SPURGE and many other broadleaf lawn weeds; some of which
are shown on the side panel.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	6.66
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid..	1.917
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid).....	0.817
INERT INGREDIENTS.....	90.623
TOTAL	100.007

THIS PRODUCT CONTAINS:

- *0.469 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid per
gallon or 5.50%.
- **0.135 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 1.59%.
- ***0.058 lb. 3,6-dichloro-o-anisic acid per gallon or 0.676%.
- **Isomer Specific by AOAC Method 6.D01-5.

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

See side panels for additional precautionary information.

TRIMEC(R) is a registered trademark of PBI/GORDON CORPORATION

U.S. Patent No. 3,284,186

NET CONTENTS ONE QUART

WEEDS

Bedstraw
Black Medic
Buckhorn
Bull Thistle
Burdock
Buttercup
Chicory
Chickweed
Clover
Cockleburr
Dandelion
Dock
Galinsoga
Ground Ivy
Hawkweed
Heal-All
Henbit
Knotweed
Lambsquarter
Lespedeza
Mallow
Morning Glory
Mustard
Oxalis
Peppergrass
Pigweed
Plantain
Poison Ivy
Poison Oak
Purslane
Ragweed
Sheep Sorrel
Shepherd's Purse
Speedwell
Spurge
Tansy Mustard
Thistle
Velvet Leaf
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow
and many other
broadleaf lawn weeds.

980/1082

EPA Reg. No. 2217-695
EPA Est. No. 2217-KS-1

MADE BY  **pbl/gordon**
corporation
KANSAS CITY, KANSAS 64108



STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes with plenty of water; for skin, wash with soap and water. If irritation develops, get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Do not store near food or feedstuffs. Do not store near fertilizers, seeds, insecticides or fungicides. Store in original container and place in locked storage area when not using.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse can. Rinse thoroughly before discarding in trash.

BEST AVAILABLE COPY

OTHER USE PRECAUTIONS:

Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solutions near these plants. Whether using a hose-end sprayer or pressure-sprayer, do not use sprays that are fine because they can drift and cause damage to non-target plants. Coarse spray droplets are less likely to wind-drift. Use only lawn-type sprayers. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of trees and other ornamental species. This product may be injurious to St. Augustine, bent and centipede grasses under conditions of high temperature and moisture. Under these conditions, or, in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage may occur.

1

37

GENERAL: TRIMEC^(R) Lawn Weed Killer For Southern Grasses is a TRIMEC(R) complex, a combination of three proven weed killers - 2,4-D, Mecoprop, and Dicamba. Together, they control a wide range of broadleaf lawn weeds; some of which are shown on the side panel. This formulation is low in 2,4-D for maximum safety to sensitive grasses such as St. Augustine, bentgrass, and buffalograss.

WHEN TO USE: Spraying should be done when broadleaf weeds are small and tender for best results. If weeds are older, they're going to be harder to kill. Avoid spraying St. Augustine grass under conditions of high temperature and moisture. Under these conditions, or, in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage may occur.

1. For best results, mow 1 to 2 days before treating.
2. Spray at any time during the growing season when weeds are actively growing, although spray works faster with temperatures above 70°F.
3. Irrigate the lawn 2 or 3 days prior to spraying to ensure that weeds are growing and able to absorb the herbicide.
4. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a "spot-treatment" may be made at any time with a small pressure-sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots.
5. Do not water for 24 hours after treatment.
6. Do not mow for 48 hours after treatment.

Poor weed control may result if spray is applied during drought or just before rain.

HOW TO USE IN THE SPREADER KING LAWN HERBICIDE APPLICATOR:

*Success
10/11/71*

1. Determine lawn size by multiplying length X width = square feet
2. Select the proper lawn size in column 1 below.
3. Select the proper amount of TRIMEC(R) LAWN WEED KILLER from column 2 and pour into the SPREADER KING bottle.
4. Fill bottle with water to amount shown in column 3.
5. Push SPREADER KING at average walking speed being careful not to overlap treated areas.

COLUMN 1 LAWN SIZE SQUARE FEET	COLUMN 2 AMT SPREADER KING LAWN WEED KILLER		COLUMN 3 FILL BOTTLE TO PINTS
	Tbsp.	Fl. Oz.	
1,000	8	4	1
2,000	16	8	2
3,000	24	12	3
4,000	36	16	4
5,000	40	20	5
6,000	48	24	6
7,000	56	28	7
8,000	64	32	8

HOW MUCH TO USE-HOSE-END SPRAYER: Use a hose-end sprayer designed for lawns. Put recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2), and spray on area shown in Column 3. For example, to spray 2,500 sq. ft., measure 10 ozs. (20 tbsp.) in jar, add water to the 10 gallon level, and mix. Turn on hose and spray as evenly as possible over 2,500 sq. ft.

COLUMN 1		COLUMN 2	COLUMN 3
Amt. To Use: (Tbsp.)	(Fl.Oz.)	Fill Bottle To: (Gallons)	Area To Treat: (Sq. Ft.)
10	5	5	1,250
20	10	10	2,500
30	15	15	3,750
40	20	20 (Full)	5,000

PRESSURE-SPRAYERS: Measure indicated amount into tank and dilute with water according to the following chart. Use a spray pattern that gives small spray droplets without any fine mist, and cover the area indicated.

Amt. To Use: (Tbsp.)	(Fl.Oz.)	Amt. Of Water In Sprayer (Gallons)	Area To Be Sprayed: (Sq.Ft.)
2	1	1	250
4	2	2	500
6	3	3	750

SPOT-SPRAYING: To spray small areas or individual weeds, do not use a hose-end sprayer. Mix 1 Oz. (2 tbsp.) in $\frac{1}{2}$ gallon of water in a pressure-sprayer, and spray as required, being certain to "wet" each weed.

BEST AVAILABLE COPY

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.