

READ THE ENTIRE LABEL FIRST
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops, get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Once injury may result. Do not spray where grass is emerging from dormant. Do not apply to newly seeded lawn grasses until well established. Seed can be safely sown 3-4 weeks after application. Do not use on carpetgrass, dichondra, or on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed labeled dosage. Be particularly careful when spraying trees. This product may be injurious to St. Augustine, berm, and Centipede grasses under conditions of high temperature and moisture. Under these conditions, or in cases of accidental overuse, some grillothermally sprouting of the St. Augustine may occur.

RECEIVED
JUL 9 1987

TRIMEC®
LIQUID WEED & FEED
For Sensitive Grasses

15-0-0

NET CONTENTS ONE QUART
(NET WEIGHT: 2 LBS. 5 OZ.)

Contains TRIMEC³ Turf Herbicide

— KEEP FROM FREEZING —

1 QUART COVERS 3500 TO 5000 SQ. FT.

KILLS: Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds; some of which are shown on this label.

The fast, easy way to weed control and fertilize your lawn.

No Spreading — No Complicated Mixing
Just attach to a garden hose & spray

ACTIVE INGREDIENTS

*Dimethylamine Salt of 2 (2-methyl-4-chlorophenoxy) propionic acid 3.85%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 1.11%
***Dimethylamine Salt of Dicamba (3,6-dichloro-p-anisic acid) 0.47%

INERT INGREDIENTS

TOTAL 100.00%

This Product Contains:

**0.293 lb. 2 (2-methyl-4-chlorophenoxy) propionic acid per gallon or 3.18%
*0.085 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 0.92%
***0.036 lb. Dicamba (3,6-dichloro-p-anisic acid) per gallon or 0.39%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

TRIMEC is a registered trademark of PBI Gordon Corporation
U.S. Patent No. 3,284,186

Mfd. by **G** pbi/gordon
corporation
KANSAS CITY, KANSAS 66118



GUARANTEED ANALYSIS

Total Nitrogen (N) 15.00%
15% Urea Nitrogen
Derived from Urea

No Spreading — No Complicated Mixing
Just attach to a garden hose & spray

Where to Use: Liquid Weed & Feed is especially formulated for use on sensitive southern grasses - St. Augustine, fescue, rye or zoysia lawns. For lawns containing sensitive grasses, use Liquid Weed & Feed at the 1 quart for 5,000 square feet rate. For lawns with non-sensitive grasses, use Liquid Weed & Feed at the 1 quart for 3,500 square feet rate. Liquid Weed & Feed is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep-sorrel and spurge. Liquid Weed & Feed also controls the easy-to-kill weeds such as chickweed, clover, dandelion, knotweed, and plantains. In addition, Liquid Weed & Feed also feeds your lawn with a 15% urea nitrogen. For best results, spraying should be done when weeds are young and tender. Older weeds are more difficult to kill and may require a retreatment in 30 days to achieve complete control.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or sod down in the bare spots.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE:

SHAKE WELL BEFORE USING.

*Connect spray nozzle to Garden hose, remove safety cap from container, attach container to end of nozzle.
*Turn on water, place finger over hole on right side of nozzle and proper ratio of LIQUID WEED & FEED will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of this liquid has been used.)

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on the label.

813-1185
EPA REG. NO. 2217-718
EPA EST. NO. 2217-KS-1

- WEEDS**
- Bedstraw
 - Black Medic
 - Buckhorn
 - Bull Thistle
 - Burdock
 - Buttercup
 - Chicory
 - Chickweed
 - Clover
 - Cocklebur
 - Dandelion
 - Dock
 - Gainsoya
 - Ground Ivy
 - Hawkweed
 - Heal All
 - Henbit
 - Knotweed
 - Lambsquarter
 - Lepedeza
 - Matow
 - Morning Glory
 - Mustard
 - Oxalis
 - Peppergrass
 - Pigweed
 - Plantain
 - Poison Ivy
 - Poison Oak
 - Purslane
 - Ragweed
 - Sheep Sorrel
 - Shepherd's Purse
 - Speedweed
 - Tansy Mustard
 - Thistle
 - Spurge
 - Vernal Leaf
 - Wild Carrot
 - Wild Garlic
 - Wild Lettuce
 - Wild Onion
 - Yarrow
 - and many other broadleaf lawn weeds

BEST AVAILABLE COPY