



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

**MEASURING
GUIDE**

1,000 sq. ft.
coverage

3,000 sq. ft.
coverage

5,000 sq. ft.
coverage
when empty

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Do not allow spray to drip to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Seed can be safely sown 3-4 weeks after application. Do not use on carpetgrass, dichondra, or on lawns or turf where desirable flowers are present. Do not spray mids of trees and ornamentals. Do not exceed specified dosages. Be particularly careful within dripline of trees. This product may be injurious to St. Augustine, bent and Centipede grasses under conditions of high temperature and moisture under these conditions, or in cases of Argentine, Argentine spurge, and temporary desiccation of the St. Augustine linkage may occur.

**TRIMEC®
LIQUID WEED & FEED
For Sensitive Grasses**

15-0-0

**NET CONTENTS ONE QUART
(NET WEIGHT: 2 LBS. 5 OZ.)**

**Contains TRIMEC® Turf Herbicide
— KEEP FROM FREEZING —**

1 QUART COVERS 3,500 TO 5,000 SQ. FT.

**KILLS: Dandelion, Chickweed, Knotweed, Henbit,
Plantains, Clover, Spurge and many other broadleaf
lawn weeds; some of which are shown on this label.**

The fast, easy way to weed control and fertilize your lawn

**No Spreading — No Complicated Mixing
Just attach to a garden hose & spray**

ACTIVE INGREDIENTS

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	3.85%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	1.11%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.47%
INERT INGREDIENTS	94.57%
This Product Contains	TOTAL 100.00%
**0.293 lb. 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 3.18%	
***0.085 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 0.92%	
***0.036 lb. Dicamba (3,6-dichloro-o-anisic acid) per gallon or 0.39%	

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautions and statements.

TRIMEC® is a registered trademark of PBI-Gordon Corporation.

Mfd. by  **pbi/gordon**
Corporation
KANSAS CITY, KANSAS 66116



GUARANTEED ANALYSIS

Total Nitrogen (N) 15.00%
15% Urea Nitrogen
Derived from Urea

**No Spreading — No Complicated Mixing
Just attach to a garden hose & spray**

WHERE TO USE:

LIQUID WEED & FEED is especially formulated for use on sensitive southern grasses — St. Augustine, bentgrasses, bahia, bermuda and centipede — as well as on bluegrass, fescue, rye or zoysia lawns. For lawns containing sensitive grasses, use **LIQUID WEED & FEED** at the 1 quart for 5,000 sq. ft. rate. For lawns with non-sensitive grasses, use **LIQUID WEED & FEED** at the 1 quart for 3,500 sq. ft. rate. **LIQUID WEED & FEED** is especially effective on all hard to kill weeds such as poison ivy, poison oak, sheepsores and spurge. **LIQUID WEED & FEED** also controls the easy to kill weeds such as chickweed, clover, dandelion, knotweed and plantains. In addition, **LIQUID WEED & FEED** also feeds your lawn with a 15% urea nitrogen. For best results, spraying should be done when weeds are young and tender. Older weeds are more difficult to kill and may require a retreatment in 30 days to achieve complete control.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or sod down in the bare spots.
3. Post-weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn with 24 hours after treatment.

HOW TO USE:

SHAKE WELL BEFORE USING

Connect spray nozzle to garden hose, remove safety cap from container and attach to nozzle. Turn on water, pull trigger, and spray on right side of nozzle at proper rate. **LIQUID WEED & FEED** will automatically be deposited. Spray evenly, covering the entire area to be treated, and thoroughly. Take care to mist spray nozzle to see how much water has been used.

NOTICE: Do not apply this product to trees, shrubs, flowers, lawns or other desirable plants since injury may result. Do not apply to newly seeded lawn grasses until well established. Seed can be safely sown 3-4 weeks after application. Do not use on carpetgrass, dichondra, or on lawns or turf where desirable flowers are present. Do not spray mids of trees and ornamentals. Do not exceed specified dosages. Be particularly careful within dripline of trees. This product may be injurious to St. Augustine, bent and Centipede grasses under conditions of high temperature and moisture under these conditions, or in cases of Argentine, Argentine spurge, and temporary desiccation of the St. Augustine linkage may occur.

411-987
EPA REG. NO. 221-1116
EPA EST. NO. 221-1116

- WEEDS**
- Bedstraw
 - Black Medic
 - Buckhorn
 - Bull Thistle
 - Burdock
 - Burrhead
 - Chicory
 - Chickweed
 - Clover
 - Cocklebur
 - Dandelion
 - Dock
 - Galinsoga
 - Ground Ivy
 - Hawkweed
 - Heal All
 - Henbit
 - Knotweed
 - Lambsquarter
 - Lespedeza
 - Mallow
 - Morning Glory
 - Mustard
 - Oxalis
 - Peppergrass
 - Pigweed
 - Plantain
 - Ragwort
 - Ragweed
 - Sheepsores
 - Shepherd's Purse
 - Spinehead
 - Tansy Mustard
 - Thistle
 - Trifolium
 - Witchgrass
 - Wild Onion
 - Wild Yam
 - Yarrow
 - and many other broadleaf lawn weeds.

BEST AVAILABLE COPY

Handwritten mark

