


PM 23 021-625 Notification 10/23

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number 195721
	Application for Pesticide:		

Section I

1. Company/Product Number 2217-695	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) TrimecTM Lawn Weed Killer For Southern or Northern Grasses	PM# (PM 23)	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation PO Box 14090 1217 West 12th Street Kansas City, MO 64101 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)



NOTIFICATION: Revision of the wetlands warning statement within the Environmental Hazards section according to definition stated in Pesticide Regulation (PR) Notice 93-3, "Labeling Statement Prohibiting Application to Water", March 9, 1993. This is to certify that this notification meets every requirement of PR-Notice 88-6. Three copies of the revised labeling are enclosed.

NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 93-3

Section III

1. Material This Product Will Be Packaged In:				DATE: <u>7/12/93</u>
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted.		If "Yes," Unit Package wgt. No. per container	If "Yes," Package wgt. No. per container	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other (_____) <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name James A. Armbruster, Ph.D.	Title Mgr. Regulatory Services	Telephone No. (Include Area Code) (816) 421-4070
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Mgr. Regulatory Services	
4. Typed Name James A. Armbruster, Ph.D.	5. Date May 20, 1993	

2016

TRIMEC® LAWN WEED KILLER

FOR SOUTHERN OR NORTHERN GRASSES

For use on Southern Grasses including St. Augustinegrass, Centipedegrass, Bentgrass, Bahiagrass, and Bermudagrass.
and

For use on Northern Grasses including Kentucky bluegrass, perennial ryegrass, and the fescue species.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	6.66%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	1.91%
Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.81%
INERT INGREDIENTS	90.62%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- *0.469 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 5.50%
 - **0.135 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 1.59%
 - ***0.058 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.676%
- Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**NOTIFICATION
LABEL NOT REVIEWED
PER FR NOTICE 88-8**
DATE 7/12/93

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional Precautionary Statements.

One Quart Covers Up To 8,000 Square Feet.


- KEEP FROM FREEZING -

NET CONTENTS ONE QUART

980/ AP052093

EPA REG. NO 2217-695
EPA EST. NO 2217-KS-1

Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101



326

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirt, shoes and gloves when mixing, loading, or applying this product. Wear face shield or goggles when mixing, loading or applying this product.

When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. After using this product, rinse gloves thoroughly before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops, get medical attention. Avoid contamination of food or feedstuffs.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.


STORAGE & DISPOSAL

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS: Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solutions near these plants. When using a hose-end sprayer or pressure sprayer, do not use fine sprays because they can drift and cause damage to non-target plants. Coarse spray droplets are less likely to wind drift. Use only lawn-type sprayers. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of trees and other ornamental species. This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass under conditions of high temperature and moisture. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustinegrass may occur.



4 of 6

WHERE TO USE: TRIMEC® Lawn Weed Killer For Southern Or Northern Grasses is a combination of three proven weed killers--2,4-D, Mecoprop, and Dicamba. This formulation will control a wide range of broadleaf weeds in residential and ornamental lawns. Established turf of northern (cool season) and southern (warm season) grasses may be treated. The northern (cool season) turfgrasses include Kentucky bluegrass, perennial ryegrass and the fescues. The southern (warm season) turfgrasses include St. Augustinegrass, centipedegrasses, bentgrass, bahiagrass and bermudagrass.

WHEN TO USE: For best results spraying should be done when broadleaf weeds are young and actively growing. Older weeds will be harder to kill.

The maximum application rate to turf is 0.3 pounds 2,4-D acid equivalent per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Spray at any time during the growing season when weeds are actively growing, although spray works faster with temperatures above 70° F. To avoid damage do not apply to St. Augustinegrass or centipedegrass while emerging from winter dormancy. Damage may also occur on these grasses if temperatures drop below 45° F within 72 hours of application. Avoid spraying St. Augustinegrass when temperatures are above 90° F and under conditions of high humidity. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustinegrass may occur.
2. Irrigate the lawn 2 or 3 days prior to spraying to ensure that weeds are growing and able to absorb the herbicide.
3. Mow 1 to 2 days before treating.
4. On new lawns, wait until the grass has reached a height of 2 inches before making a broadcast application. However, a "spot-treatment" may be made at any time with a pressure sprayer. Wait 3 weeks before seeding grass, planting sprigs or planting stolons.
5. Do not water for 24 hours after treatment.
6. Do not mow for 48 hours after treatment.

Poor weed control may result if spray is applied during drought or just before rain.

(Option 1) UNIT PACKAGING. (The entire contents of the container is to be applied in a single application. A siphon tube and a hose-end sprayer are attached to the immediate containers.)


HOW TO USE: No mixing of this product with water is required. For the southern (warm season) grasses, apply the entire contents of this container (32 fluid ounces) on 8,000 square feet of turf. For the northern (cool season) grasses, apply the entire contents of the container (32 fluid ounces) on 6,000 square feet of turf. The assembly and application instructions are presented as follows:

ASSEMBLY INSTRUCTIONS:

1. Attach siphon tube to hose-end sprayer attachment.
2. Remove cap from bottle and insert sprayer attachment.
3. Securely tighten sprayer attachment on bottle.
4. Screw hose into hose-end sprayer attachment.

APPLICATION INSTRUCTIONS:

1. Turn on water. **TO START FLOW OF PRODUCT:** Place Finger over hole on top right side of sprayer attachment. Hold about 3 feet above area to be treated, moving at a steady pace to ensure uniform coverage.
2. **TO STOP FLOW OF PRODUCT:** Remove finger from hole to break siphon action. **NOTE:** To ensure adequate coverage, use moderate to high water pressure when applying through hose-end sprayer attachment.



(Option 2) HOSE-END SPRAYERS:

HOW TO USE :

For southern (warm season) grasses, apply 4 fluid ounces of product per 1,000 square feet of turf. Refer to the spray preparation chart for southern grasses for areas larger than 1,000 square feet. Pour the recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2), and spray on area shown in Column 3. For example, to spray 2,500 square feet measure 10 fluid ounces (20 tablespoons) in jar, add water to the 10 gallon level, and mix. Turn on water and spray uniformly over 2,500 square feet.

SOUTHERN GRASSES including St. Augustinegrass, centipedegrass, bentgrass, bahiagrass and bermudagrass.

Column 1		Column 2	Column 3
Amount of Product		Fill Bottle To	Area to Treat
(Tbsp.)	(Fluid Ounces)	(Gallons)	(Square Feet)
10	5	3	1,250
20	10	5	2,500
30	15	8	3,750
40	20	10	5,000

For Northern (cool season) grasses apply 5 1/3 fluid ounces per 1,000 square feet of established turf. Refer to the spray preparation charts for cool season grasses for areas larger than 1,000 square feet.

NORTHERN GRASSES including Kentucky bluegrass, perennial ryegrass and the fescue species.


Column 1		Column 2	Column 3
Amount of Product		Fill Bottle To	Area to Treat
(Tbsp.)	(Fluid Ounces)	(Gallons)	(Square Feet)
10	5	3	938
20	10	5	1,876
30	15	8	2,814
40	20	10	3,752

(Option 3)

FOR SMALL AREAS WITH KNAPSACK SPRAYERS, BACKPACK SPRAYERS, CART SPRAYERS, PRESSURE SPRAYERS OR OTHER MANUALLY OPERATED SPRAYERS: Determine the area by measuring and prepare the appropriate amount of spray mixture. Refer to the spray preparation charts for southern and northern grasses presented below.

SOUTHERN GRASSES including St. Augustinegrass, centipedegrass, bentgrass, bahiagrass and bermudagrass.

Amount of Product		Water Volume	Area to Treat
(Tbsp.)	(Fluid Ounces)	(Gallons)	(Square Feet)
8	4	1	1,000
24	12	3	3,000



6216

NORTHERN GRASSES including Kentucky bluegrass, perennial ryegrass and the fescue species.

Amount of Product		Water Volume	Area to Treat
(Tbsp.)	(Fluid Ounces)	(Gallons)	(Square Feet)
10 2/3	5 1/3	1	1,000
32	16	3	3,000

For Spot treatments of individual weeds in southern or northern turfgrasses, mix 2 fluid ounces of product with 1 gallon of water. Thoroughly wet individual weeds. Do not use hose-end applicators for spot treatments.

--WEEDS--

- | | | | |
|--------------|--------------|----------------|----------------|
| Bedstraw | Galingosa | Peppergrass | Thistle |
| Black Medic | Ground Ivy | Pigweed | Velvetleaf |
| Buckhorn | Hawkweed | Plantain | Wild Garlic |
| Bull Thistle | Heal-All | Poison Ivy | Wild Lettuce |
| Burdock | Henbit | Poison Oak | Wild Onion |
| Buttercup | Knotweed | Purslane | Wild Carrot |
| Chickweed | Lambsquarter | Ragweed | Yarrow |
| Chicory | Lespedeza | Sheep Sorrel | and many other |
| Clover | Mallow | Shepherdspurse | broadleaf |
| Cocklebur | Morningglory | Speedwell | lawn weeds |
| Dandelion | Mustard | Spurge | |
| Dock | Oxalis | Tansy Mustard | |

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

