

GORDON'S 2,4-D WEED & FEED

12-3-5

PROVIDES FERTILIZATION
CONTROLS LAWN WEEDS.
TREATS 5,000 SQ. FT.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 1.14%
INERT INGREDIENTS 98.86%
TOTAL 100.00%

THIS PRODUCT CONTAINS:

19.0 lbs. or 0.95% of 2,4-Dichlorophenoxyacetic acid per ton or 1.49 lbs. per acre.

Isomer Specific by AOAC Method 6.D01-5.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

NET WEIGHT 18 POUNDS

Mfg. by **G** pbi / Gordon
CORPORATION
 KANSAS CITY, KANSAS 66108

ACCOUNTED
 APR 8 1988

2217-697



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

1061
 PM 23
 2217-697

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

STORAGE & DISPOSAL

Do not store near other pesticides or seeds. Do not reuse empty container. Wrap and put in trash collection.

USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS

Gordon's Weed & Feed 2,4-D controls dandelion, plantains, and other weeds commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dichondra, carpetgrass, bentgrass or St. Augustine lawns.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

SPREADER SETTINGS

Scotts Precision Flow	5.75
Scotts Broadcast	J
Sears Broadcast	4.7
Sears Drop	8.0
Ortho 4000	3.5
Ortho 3000	7.0

FERTILIZER GUARANTEE

Total Nitrogen (not less than)	12%
Water Insoluble Nitrogen (not less than)	3%
Available Phosphoric Acid (not less than)	3%
Soluble Potash (not less than)	5%

WEED CONTROL

2,4-D for Control of Broadleaf Plantain, Buckhorn Plantain, Dandelion, Purslane, Pigweed and Smartweed.

Notice: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 2217-697
 EPA EST. NO. 2217-KS-1

559/1182