J.	2217-70		1/2/3/19	7	1 1	Approval expires 2-28-95
Please read instructions on reverse before completing form. United States Environmental Protection Washington, DC 204			Form Approve	Registra Amendn Other	tion	OPP Identifier Number
	Application	n for Pestic	ide - Section	<u> </u>		
1. Company/Product Number 2217-709		2. EP/	2. EPA Product Manager James A. Tompkins			posed Classification
4. Company/Product (Name) Quadmec® Turf Herbicide		PM#	PM# (PM 25) 2 2 By			None Restricted
5. Name and Address of Application PBI/Gordon Corporation P.O. Box 014090 1217 West 12th Street Kansas City, MO 64101	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name					
	กดาเราริยุปโก	N Section -	- 11			
Notification - Explain bell Explanation: Use additional in NOTIFICATION of Minor Labeling containing 2,4-D. Jack Houseng 2,4-D products from dedicated of Hammond, Chair of the Industry labeling is enclosed. This notific made to the labeling or the confirstatement to the EPA. I further unviolation of FIFRA and I may be stated that the EPA is further unviolation of FIFRA and I may be contained to the Industry labeling or the confirmation of FIFRA and I may be stated that the EPA is further unviolation of FIFRA and I may be contained to the Industry labeling in the I	page(s) if necessary. (For section ng Revision, PR Notice 95-2: We arrier, OPP Special Review Branch Chrontainers larger than 5 gallons. This trask Force for 2,4-D Research Datation is consistent with the provision dential statement of formula of this understand that if this notification is subject to enforcement action and personal page 1. Init Peckaging Yes No "Yes" No. per container Ormation 4. Size(s) Research Datation and personal page 1.	n I and Section II. e submitting a minief, recognized that a notification is the ta, Pesticide and This of PR Notice 95 product. I understanot consistent with enables with the Notice Section 1 and 1	or label revision for the spigots may be more addition of pre-approximate the spigots may be more addition of pre-approximate the terms of PR Not lipp 12 and 14 of FIF Packaging No. per container	ated cation. below. ne engineering of the practical than wed language st., February 1, 19 ms at 40 CFR 1: n of 18 USC Seice 95-2 and 40 RA. 2. Type of the control of Late On Labe	control state pump and sated in a le 95. One (1) 52.46, and c. 1001 to v CFR 152.4 Container Metal Plastic Glass Paper Other (S	probe systems in dispensing fiter from the Agency to Larry copy of the proposed no other changes have been willfully make any false 16, this product may be in
6. Manner in Which Label is Aft	fixed to Product Lithog Paper Stenc	ʻgived Iraph	Other _			
	Stenc	Section -	11/			
1. Contact Point. (Complete its	ems directly helow for identification			oraceary to a	racase this	application !
Name	n of individual to be contacted, if necessary, to process this Title Telephol			T	e No. (Include Area Code)	
Craig Martens		Manager, Regulatory Services				16) 421-4070
· · · · · · · · · · · · · · · · · · ·	Certifica ents I have made on this form and knowingly false or misleading sta w.	s all attachments	•		•	6. Date Application Received (Stamped)
2. Signature	3. Title					
Caughantens_		Manager, Regulatory Services				
4. Typed Name	5. Date	-			e estat	
Craig Martens	August 29, 19	997			• • •	

295

QUADMEC® TURF HERBICIDE POSTEMERGENT GRASS & BROADLEAF HERBICIDE

For selective control of grass and broadleaf weeds in Bermudagrass, Bluegrass, Tall Fescue, Turf-Type Tall Fescues, and Zoysiagrass Turf.

One Gallon Covers Up To One Acre.

KEEP FROM FREEZING - SHAKE WELL BEFORE USING

ACTIVE INGREDIENTS:

*Monosodium acid methanearsonate,	18.00%
**Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	5.83%
· · · · · · · · · · · · · · · · · · ·	
***Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	
****Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	1.46%
INERT INGREDIENTS:	<u>68.85%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- *1.80 lbs, monosodium methanearsonate per gallon or 18.00%. Total Arsenic (as elemental) all in water soluble form 8.33%
- **0.48 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.84%

AND PROPERTY OF THE PROPERTY O

- ***0.48 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.84%
- *****0.12 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.21%


Isomer Specific by AOAC Method.

QUADMEC® and TRIMEC® are registered trademarks of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

WARNING

NOTIFICATION



See side panels for additional Precautionary Statements and Statement of Practical Treatment.

NET CONTENTS ONE U.S. CALLONS (5785 UTERS)

700/ AP082997
EPA REG NO 2217-709
EPA EST NO 2217-KS-1
MANUFACTURED BY:

pbi/Gardon

Corporation



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

Total Carlo Car

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Wear face shield or goggles when mixing, loading or applying this product. When mixing, loading, or applying this product, wear long-sleeved shirt, long pants, socks, shoes, chemical-resistant gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection. See protective clothing statement in Directions for Use.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more: Do not open-pour product from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Statement of Practical Treatment

IF IN EYES: Flush with plenty of water and get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash promptly with soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from treated area. Do not contaminate water used for domestic consumption, or by wildlife or aquatic life, or for irrigation purposes.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

FOR USE ON RESIDENTIAL AND OTHER TURF SITES EXCLUDING SOD FARMS.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dusts has settled.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pecticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Page 2 of 4

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

militaria de la composição de la composi

- GENERAL INFORMATION -

Refer to partial list of weeds controlled by Quadmec® Turf Herbicide. This product controls a wide variety of problem weed species including crabgrass, dallisgrass, nutsedge, spurge, dandelion and oxalis.

THE FOLLOWING PROTECTIVE CLOTHING REQUIREMENTS MUST BE FOLLOWED.

Wear protective clothing when handling or applying this product including long pants, long-sleeved shirt, and impermeable gloves and boots. Mixer/loaders should include an apron and full-face shield when handling or mixing concentrate. Spray rig applicator should wear a mask or respirator approved by the Mining Enforcement and Safety Administration and the National Institute for Occupational Safety and Health.

- APPLICATION INFORMATION AND USE PRECAUTIONS -

Do not apply this product through any type of irrigation system. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not use a hose end sprayer.

Coarse sprays are less likely to wind drift. Avoid fine mists. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area, be particularly careful within the dripline of trees or other ornamental species.

Do not spray while grass is emerging from dormancy. Do not apply to newly seeded lawns until after the third mowing. Seed can be sown 3 to 4 weeks after application.

Do not use on St. Augustinegrass, centipedegrass, carpetgrass, red fescues, dichondra, nor on lawns or turf where desirable clovers are present.

Weed control is best achieved when temperatures are above 70°F. However, DO NOT apply this product to Kentucky bluegrass, tall fescue, or turf-type tall fescues when air temperatures exceed 80°F nor when drought conditions exist. DO NOT apply this product to bermudagrass or zoysiagrass when air temperatures exceed 90°F nor when drought conditions exist.

Be sure soil is moist and lawn healthy before application. Temporary discoloration may occur and high spray volumes may reduce the turf discoloration. If turf is showing stress due to high temperatures and/or drought, irrigation is recommended 2 to 3 days after treatment.

Failure to observe these use precautions may result in injury.

Keep children and domestic animals off treated area until this material has been watered into the soil.

After using this product, clean sprayer with Gordon's Spray Tank Cleaner and rinse thoroughly with water before applying other pesticides.

ESTABLISHED BERMUDAGRASS AND ZOYSIAGRASS:

Mix 3 to 5 fluid ounces of Quadmec® Turf Herbicide in 1 to 3 gallons of water for treatment of 1000 square feet or 1 to 12/3 gallons of Quadmec® Turf Herbicide in 50 to 100 gallons of water for treatment of one acre of turf. Better control is obtained by making applications shortly after crabgrass emergence in the spring which will also control surviving winter annual broadleaf weeds and emerged summer annual broadleaf weeds. A second application may be necessary 10 to 14 days later for dense weed populations, especially dallisgrass. Do not apply this product to bermudagrass or zoysiagrass when air temperatures exceed 90°F nor when turf is under stress such as drought.

Zoysiagrass cultivars may vary in the levels of tolerance to applications containing MSMA.

The maximum application rate to turf is 0.9 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

ESTABLISHED KENTUCKY BLUEGRASS, TALL FESCUE, TURF-TYPE TALL FESCUES (IMPROVED AND LOW GROWING):

Mix 3 fluid ounces of Quadmec® Turf Herbicide in 1 to 3 gallons of water for treatment of 1,000 square feet of turf or 1 gallon of Quadmec® Turf Herbicide in 50 to 100 gallons of water per acre of turf. Make a second application at the same rate 10 to 14 days after the first application. Make spring applications to newly emerged crabgrass and for broadleaf weed control, including oxalis. Summer applications can be made to more mature crabgrass and to broadleaf weeds. Do not apply this product to Kentucky bluegrass, tall fescue, or turf-type tall fescues when air temperatures exceed 80°F nor when turf is under stress such as drought.

Injury from this product may occur to mixed stands of Kentucky bluegrass and red fescues (chewings fescue, creeping fescue, and spreading fescues). If you are unwilling to accept some degree of turf injury, then do not use this product for mixed stands.

- SPOT TREATMENTS -

SPOT TREATMENTS FOR ESTABLISHED BERMUDAGRASS, ZOYSIAGRASS, KENTUCKY BLUEGRASS, TALL FESCUE, AND TURF-TYPE TALL FESCUES:

Mix 3 fluid ounces of Quadmec[®] Turf Herbicide in 1 gallon of water for treatment of 1,000 square feet of turf. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants. Withhold sprinkling or irrigation for 24 hours after application. Two or more repeat applications at 10 to 14 day intervals may be required. Temporary discoloration of the turfgrass may occur.

- PARTIAL LIST OF WEEDS CONTROLLED -					
Aster	Healall	Purslane			
Bedstraw	Henbit	Ragweed			
Black Medic	Knotweed	Sandbur			
Buckhorn	Lambsquarters	Sheep Sorrel			
Burdock	Lespedeza	Shepherdspurse			
Chicory	Mallows	Speedwell			
Chickweed	Morningglory	Spurge			
Clovers	Nutsedge '	Wild Carrot			
Crabgrass	Oxalis	Wild Garlic .			
Dallisgrass	Peppergrass	Wild Lettuce			
Dandelion	Pigweeds	Wild Onion			
Docks	Plantains	Yarrow			
Goosegrass	Poison Ivy	Yellow Woodsorrel			
Ground Ivy	Poison Oak				

LIMITED WARRANTY AND DISCLAIMER.

Butter the second state of the time to the state of the property of the state of the property of the state of

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.