

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

James L. Kuntsman, Ph.D. PBI/Gordon Corporation 1217 W. 12<sup>th</sup> Street P.O. Box 014090 Kansas City, Missouri 64101

JAN 24 2013

Subject:

Notification; Per PR-Notice 98-10

Quadmec Turf Herbicide EPA Reg. No. 2217-709

Date Submitted: January 24, 2013 (emailed)

#### Dear Dr. Kuntsman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated January 24, 2013 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact me at (703) 306-0415 or davis.kable@epa.gov.

Sincerely.

Kable Bo Davis
Product Manager 25

Herbicide Branch

Registration Division (7505P)

2/12

Please read instructions on revi	erse before completing to	orm.		Form Approve	ed. OMB No.	2070-0060.	Approval Expires 2-28-95
<b>≎</b> EPA	Environmenta	nited States al Protection ngton, DC 20460	Agency		Registra Amend Other	2. 1. 1	OPP Identifier Number
	Apı	olication for I	Pesticide	- Section	<b>4</b>	V	
1. Company/Product Number				roduct Manag Kable Bo D	er	3. Pr	oposed Classification
4. Company/Product (Name Quadm	) ec Turf Herbicide		PM# Produ	uct Manager	—Team 25	X	None Restricted
5. Name and Address of Ap PBI/Gordon Corporatio Post Office Box 01409 Kansas City, Missouri (	n D		6. Exped (b)(i), my EPA Reg Product N	ited Review.	In accordan	ce with FII	
		Sec	<u>tion – II                                 </u>			D. W. Z. S. Z.	P1/74 8-9-1/7-18 7
Amendment - Explain below.  Resubmission in response to Agency letter dated  Notification - Explain below.			A <sub>2</sub>	nal.printed labe gency letter date fe Too" Applica ther - Explain be	ed tion.		FICATION 2 4 2013
Explanation: Use additional NOTIFICATION: Lab Amendments. Section I	eling notification p I(N): Other Revision	er PRN 98-10: I	Notificatio				
			ion – III		<del> </del>		
Material This Product Will Be	e Packaged In:						
Yes* No * Certification must		per If "Yes		kaging .  No. per  Container	2. Type of C	al tic ss	
3. Location of Net Contents Info	<b>!</b>	Size(s) Retail Contai	ner	5. Le	ocation of Lab On Label On Labeling	el Direction	
6. Manner in Which Label is Affi	xed to Product	Lithograph Paper glued Stenciled	[	Other			
		Sect	ion – IV				
1. Contact Point (Complete item	s directly below for identi	fication of individual	to be contact	ed, if necessary	, to process th		
Name James L. Ku	nstman, Ph.D.		ector of Re	egulatory Ser	vices	•	No. (Include Area Code) 816-460-6292
I certify that the statements I acknowledge that any kno both under applicable law.  2. Signature	I have made on this form						6. Date Application Received (Stamped)
Oa I Z		Direc	irector of Regulatory Services				
4. Typed Name 5. Da James L. Kunstman, Ph.D.		5. Date	January 24, 2013				•



1217 W. 12<sup>th</sup> STREET · P.O. BOX 014090 KANSAS CITY, MISSOURI 64101-0090 816-421-4070 · 1-800-821-7925 FAX: 816-474-0462

January 24, 2013

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard (South Building)
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Kable Bo Davis, (PM-25)

Dear Mr. Davis:

NOTIFICATION: Labeling notification per PRN 98-10: Notifications Non-notifications and Minor Formulation Amendments. Section II(N): Other Revisions. Quadmec Turf Herbicide (EPA Reg. No. 2217-709)

# I. Turf applications of MSMA have been extended by the Agency:

May 27, 2012: An Agency letter prepared by Mindy Ondish accepted the draft labeling of Quadmec Turf Herbicide (EPA Reg. No. 2217-709) and stated that the requirements were satisfied for the Reregistration Eligibility Decisions of 2,4-D, MCPP-p, dicamba and MSMA.

October 1, 2012: A letter prepared by the Organic Arsenical Products Task Force affirmed that the use of MSMA for turf applications on golf courses and sod farms would be extended beyond the phase out period established during January, 2009.

January 23, 2013: Now, we are requesting to extend the period for use of existing stocks of Quadmec Turf Herbicide. We ask to delete the expiration date of December 31, 2013, or we ask remove the prohibition for use of this product on golf courses and sod farms after December 31, 2013. Please refer to page 4 of the proposed draft labeling.

#### II. Certification statement:

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR\_Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

JAN 2.4 2013

## III. Enclosures:

- 1. Application for Pesticide Amendment (EPA Form 8570-1)
- 2. One (1) copy the draft labeling with annotations.
- 3. One (1) copy the draft labeling without annotations.

If you have any questions, please call me at 816-460-6292 or contact me at <a href="mailto:jkunstman@pbigordon.com">jkunstman@pbigordon.com</a>.

Sincerely,

James L. Kunstman, Ph.D.

Director of Regulatory Services

# QUADMEC® TURF HERBICIDE

# EPA Reg. No. 2217-709

#### **ACTIVE INGREDIENTS:**

Monosodium acid methanearsonate	18.00%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	5.83%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	2.93%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	
OTHER INGREDIENTS:	71.78%
TOTAL	

#### THIS PRODUCT CONTAINS:

- 1.80 lb monosodium methanearsonate per gallon or 18.00%. Total Arsenic (as elemental) all in water soluble form 8.33%.
- 0.48 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.84%
- 0.24 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 2.42%
- 0.12 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 1.21%

Isomer Specific by AOAC Method.

NOTIFICATION

## KEEP OUT OF REACH OF CHILDREN

JAN 2 4 2013

# **WARNING - AVISO**

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find some one to explain it to you in detail.)

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

# PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**WARNING:** Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- · protective eyewear,
- long-sleeved shirt and long pants,
- · shoes and socks,
- · chemical-resistant gloves and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.



### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

### **User Safety Recommendations**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	
If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or on clothing:	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled:	<ul> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
	ct container or label with you when calling a poison control center or doctor or going for may also contact 1-877-800-5556 for emergency medical treatment advice.

#### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- · coveralls worn over short-sleeved shirt and short pants,
- · chemical-resistant footwear plus socks,
- · chemical-resistant gloves made of any water-proof material,
- · chemical-resistant headgear for overhead exposure and
- · protective eyewear.

#### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

#### **APPLICATION INFORMATION AND PRECAUTIONS:**

Weed control is best achieved when temperatures are above 70°F. However, do not apply this product to Kentucky bluegrass, tall fescue, or turf-type tall fescues when air temperatures exceed 80°F nor when drought conditions exist. Do not apply this product to bermudagrass or zoysiagrass when air temperatures exceed 90°F nor when drought conditions exist.

Be sure soil is moist and lawn healthy before application. Temporary discoloration may occur and high spray volumes may reduce the turf discoloration. If turf is showing stress due to high temperatures and/or drought, irrigation is recommended 2 to 3 days after treatment.

Failure to observe these use precautions may result in injury.

After using this product, clean sprayer with Gordon's® Spray Tank Cleaner and rinse thoroughly with water before applying other pesticides.

#### RESTRICTIONS:

This product is for use on golf courses and sod farms only.

Do not apply this product through any type of irrigation system. Do not pour spray solutions near these plants. Do not use a hose end sprayer.

Do not exceed specified dosages for any area. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.

Do not spray while grass is emerging from dormancy. Do not apply to newly seeded turfgrass until after the third mowing. Seed can be sown 3 to 4 weeks after application.

Do not use on St. Augustinegrass, centipedegrass, carpetgrass, red fescues, dichondra, or turf where desirable clovers are present.

Do not apply this product in Florida. Aerial application is prohibited.

#### Application restrictions for golf courses:

For newly constructed golf courses, only one broadcast application is allowed. For all other applications to golf courses, only spot treatments (100 square feet maximum per spot) are allowed. A spot treatment must not exceed 25% of total golf course acreage per year.

#### Application restrictions for sod farms:

The maximum application rate is 1.67 gallons of product per acre per application. The maximum number of broadcast applications is limited to 2 per season with a minimum of 30 days between applications. The maximum seasonal rate is 3.34 gallons of product per acre.

Only 2 broadcast applications are allowed per season. Do not apply within 25 feet of permanent water bodies or aquatic habitat, including, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, and estuaries.

#### **ESTABLISHED BERMUDAGRASS AND ZOYSIAGRASS:**

Mix 3 to 5 fluid ounces of Quadmec® Turf Herbicide in 1 to 3 gallons of water for treatment of 1,000 square feet or 1 to 1 2/3 gallons of Quadmec® Turf Herbicide in 50 to 100 gallons of water for treatment of one acre of turf. Better control is obtained by making applications shortly after crabgrass emergence in the spring which will also control surviving winter annual broadleaf weeds and emerged summer annual broadleaf weeds. A second application may be necessary for dense weed populations, especially dallisgrass. Do not apply this product to bermudagrass or zoysiagrass when air temperatures exceed 90°F nor when turf is under stress such as drought.

Zoysiagrass cultivars may vary in the levels of tolerance to applications containing MSMA.

# ESTABLISHED KENTUCKY BLUEGRASS, PERENNIAL RYEGRASS, TALL FESCUE, TURF-TYPE TALL FESCUES (IMPROVED AND LOW GROWING):

Mix 3 fluid ounces of Quadmec® Turf Herbicide in 1 to 3 gallons of water for treatment of 1,000 square feet of turf or 1 gallon of Quadmec® Turf Herbicide in 50 to 100 gallons of water per acre of turf. Make a second application at the same rate. Make spring applications to newly emerged crabgrass and for broadleaf weed control including oxalis. Summer applications can be made to more mature crabgrass and to broadleaf weeds. Do not apply this product to Kentucky bluegrass, tall fescue, or turf-type tall fescues when air temperatures exceed 80°F nor when turf is under stress such as drought.

Injury from this product may occur to mixed stands of Kentucky bluegrass and red fescues (chewings fescue, creeping fescue, and spreading fescues). If you are unwilling to accept some degree of turf injury, then do not use this product for mixed stands.

# SPOT TREATMENTS FOR ESTABLISHED BERMUDAGRASS, ZOYSIAGRASS, KENTUCKY BLUEGRASS, PERENNIAL RYEGRASS, TALL FESCUE, AND TURF-TYPE TALL FESCUE:

Mix 3 fluid ounces of Quadmec® Turf Herbicide in 1 gallon of water for treatment of 1,000 square feet of turf. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants. Withhold sprinkling or irrigation for 24 hours after application. Temporary discoloration of the turfgrass may occur.

#### Limitations on spot treatments for golf course and sod farm turf:

The maximum application rate is 3.0 fl.oz. per 1,000 sq.ft. per application (0.25 lb MCPP-p acid equivalent per acre). Spot treatments are limited to a maximum of 100 sq.ft. per spot treatment, and the treated area must not exceed 25% of total golf course acreage per year.

### **Spray Drift Management**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

#### **Droplet Size**

Use only Medium or coarser spray nozzles according to ASABE (Standard S572.1) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

#### Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors ontarget deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

#### **Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

### Weed spectrum

This product controls a wide variety of problem weed species including crabgrass, dallisgrass, nutsedge, spurge, dandelion and oxalis. Refer to partial list of weeds controlled by Quadmec® Turf Herbicide.

WEEDS CONTROLLED			,
Aster, white heath & white	Dayflower	Lambsquarters	Ragweed
prairie	Deadnettle	Lawn burweed	Red sorrel (*sheep sorrel)
Bedstraw	Dock	Lespedeza, common	Sandbur
Barnyardgrass	Dogfennel	Mallow, common	Shepherdspurse
Beggarweed, creeping	English daisy	Matchweed	Spotted spurge
Bindweed	False dandelion (*spotted	Mouseear chickweed	Spurge
Black medic	catsear & common catsear)	Nutsedge, yellow	Thistle
Broadleaf plantain	Field bindweed	Oxalis (*yellow woodsorrel	Veronica (*corn speedwell)
Buckhorn plantain	(*morningglory & creeping	& creeping woodsorrel)	Virginia buttonweed
Bull thistle	jenny)	Parsley-piert	White clover (*Dutch clover,
Burclover	Field oxeye-daisy	Pennsylvania smartweed	honeysuckle clover, white
Burdock, common	(*creeping oxeye)	Pennywort (*dollarweed)	trefoil & purplewort)

Buttercup, creeping	Filaree, whitestem &	Pepperweed	Wild carrot
Carpetweed	redstem	Pigweed	Wild garlic
Chickweed, common	Florida pusley	Pineappleweed	Wild geranium
Chicory	Foxtail (Green and Yellow)	Plantain	Wild lettuce
Cinquefoil	Goosegrass	Poison ivy	Wild mustard
Clover	Ground ivy	Poison oak	Wild onion
Compassplant	Groundsel	Prostrate knotweed	Wild strawberry
Crabgrass, large & smooth	Hawkweed	(*knotweed)	Wild violet
Curly dock	Healall	Puncturevine	Yarrow
Dallisgrass	Henbit	Purslane	Yellow rocket
Dandelion			
*Synonyms			

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep from freezing.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning If burned, stay out of smoke.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate

for later use or disposal. Repeat this procedure two more times.

**IOR** 

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

#### [For Refillable Containers:]

**CONTAINER HANDLING:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

# **APPENDIX**

- 1. Statements which may appear on different label components depending on packaging configuration.
  - · See next panel for additional Precautionary Statements and First Aid
  - Net Contents: \_
  - EPA Est. No.
- 2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:
  - For selective control of grass and broadleaf weeds in bermudagrass, bluegrass, tall fescue, turf-type tall fescue, and zoysiagrass turf.
  - · One gallon covers up to one acré.
  - · Keep from freezing.
  - · Shake well before using.
  - QUADMEC® and TRIMEC® are registered trademarks of PBI/Gordon Corporation.
  - Postemergent Grass & Broadleaf Herbicide
  - · For use only on golf courses and sod farms

# **DOCUMENT CONTROL INFORMATION**

1. Unique Label Identifier: 002217-00709.20130124.notif-proposed-highlighted.doc

2. Reason for Issue: remove 2013 use restriction only