

GURDON'S
PROFESSIONAL TURF PRODUCTS



2217-710 1061
PM 23

READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals
DANGER Corrosive, causes eye damage and skin irritation. Do not get into eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed.
Environmental Hazards
Do not apply directly to water. Do not apply when weather conditions favor drift away from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area. Keep from freezing. To prevent cross-contamination, do not store near other pesticides, fertilizers, seeds, food or feed.

PRODUCT DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed, by state and local authorities, by burning. If burned, stay out of smoke.

USE PRECAUTIONS:

Avoid drift of spray mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Avoid fine mists. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area. Be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 85°F. Seed can be safely sown 3 to 4 weeks after application at recommended rate. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

TRIMEC CONTROLS:

- | | |
|-------------|------------------|
| Bedstraw | Morning Glory |
| Black Medic | Peppergrass |
| Buckhorn | Pigweed |
| Burdock | Plantain |
| Chicory | Poison Ivy |
| Chickweed | Poison Oak |
| Clover | Purslane |
| Dandelion | Ragweed |
| Dock | Sheep Sorrel |
| Ground Ivy | Shepherd's Purse |
| Heart-Ali | Speedwell |
| Henbit | Spurge |
| Knotted | Wild Carr |
| Lambquarter | Wild Garlic |
| Lespedeza | Wild Lettuce |
| Mallow | Wild Onion |
| | Yarrow |

TRIMEC® LAF 637

BROADLEAF HERBICIDE

FOR SALE TO USE AND STORAGE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL ONLY

KILLS: Dandelion, Knotweed, Henbit, Chickweed, Plantain, Spurge, and many other broadleaf weeds, some of which are listed on this label.

ONE GALLON COVERS UP TO 3½ ACRES
KEEP FROM FREEZING

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid.....	41.08
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	13.90
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)...	1.67
INERT INGREDIENTS.....	43.35
TOTAL	100.00

THIS PRODUCT CONTAINS:

- *3.35 lbs. 2,4-Dichlorophenoxyacetic acid per gallon or 33.48% a.i.
- **1.13 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 11.48% a.i.
- ***0.14 lb. 3,6-dichloro-o-anisic acid per gallon or 1.40% a.i.

KEEP OUT OF REACH OF CHILDREN
DANGER

Statement of Practical Treatment


If In Eyes: In case of eye contact immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.
If On Skin: Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.
If Swallowed: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.
If Inhaled: Remove victim to fresh air and apply respiration if indicated.

U.S. Patent No. 3,284,186

TRIMEC™ is a registered trademark of PBI/GORDON CORPORATION

© 1983, PBI-GORDON CORP.

NET CONTENTS 55 U.S. GALLONS



ORNAMENTAL LAWNS & TURF (Instructions):

Apply TRIMEC at the rate of 2.3 to 4 pints per acre in sufficient water for adequate coverage. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra and carpetgrass.

APPLICATION TYPE

Professional Lawn Maintenance:	Use 2.3 to 2.7 pints per acre (84 to 104 oz. per 1,000 sq. ft.) at application rate of 1 to 3 gallons of solution spray per 1,000 sq. ft.
Mature Weeds:	Use 2.5 to 3 pints per acre (82 to 110 oz. per 1,000 sq. ft.) at application rate of 4 to 5 gallons of solution spray per 1,000 sq. ft.
	Use 3.25 to 4 pints per acre when controlling mature weeds.

DIRECTIONS

Controlled droplet applicator:

Reduced rates of TRIMEC LAF-637 must be used when grass is stressed from heat, drought, etc. Do not use on warm season grasses. Add 2 pints TRIMEC LAF-637 to the Herbicide container then fill with 3 pints of water. Spray entire contents over 33,000 sq. ft. (approx. 3/4 acre). Do not overlap between spray patterns.

St. Augustine Grass:

Apply TRIMEC LAF-637 at 2.3 pints in up to 150 gallons of water per acre (1.2 oz. TRIMEC LAF-637 in 5 gallons water per 1,500 square feet). Do not spray St. Augustine if grass is stressed from heat, drought, etc. Slight turf yellowing should disappear after about one week.

Bentgrass:

On closely mowed bentgrass (putting and bowling greens) apply TRIMEC LAF-637 at a maximum rate of one fluid ounce in 5 gallons water per 1,500 sq. ft. —preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE:

Care should be taken to avoid overdosing bentgrass and St. Augustine grass or injury may result. Large volumes of spray (i.e. one fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.



EPA REG. NO. 2217
EPA CAT. NO. 2217-10