

READ THE ENTIRE LABEL FIRST
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area if accessible to children. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, insecticides, fungicides or sprays.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly, before discarding in trash.

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Seed can be safely sown 3-4 weeks after application. Do not use on carpet grass, dichondra or on lawns or other desirable covers already present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages. Be particularly careful with ornamental trees. This product may be poisonous to fish. Apply the herbicide through the grasses under conditions of high temperature and moisture. Under these conditions, in some cases of accidental overdose some slight irritation or discoloration of the St. Augustine foliage may occur.

TRIMEC®

LIQUID WEED & FEED

For Sensitive Grasses

15-0-0

NET CONTENTS ONE QUART
(NET WEIGHT: 2 LBS. 5 OZ.)

Contains TRIMEC® Turf Herbicide

— KEEP FROM FREEZING —

1 QUART COVERS UP TO 5,000 SQ. FT.

KILLS: Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds, some of which are shown on this label.

The fast, easy way to weed control and fertilize your lawn.

No Spreading — No Complicated Mixing
Just attach to a garden hose & spray

ACTIVE INGREDIENTS

1-(2-ethyl-4-methyl-5-pyridinyl)-3-methyl-4-propionic acid	3.85%
1-(2-ethyl-4-methyl-5-pyridinyl)-3-methyl-4-propionic acid	1.11%
1-(2-ethyl-4-methyl-5-pyridinyl)-3-methyl-4-propionic acid	0.47%
INERT INGREDIENTS	94.57%

This Product Contains:

10.253 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 3.18%	TOTAL 100.00%
10.065 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 0.92%	
10.065 lb. Dicamba (3,6-dichloro-b-anisic acid) per gallon or 0.39%	

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

TRIMEC® is a registered trademark of PBI/Gordon Corporation
U.S. Patent No. 3,184,186

Mfd. by  **pbi/gordon**
corporation
KANSAS CITY, KANSAS 64118



WEEDS

Beeswax
Black Medic
Buckhorn
Bull Thistle
Burdock
Buttercup
Chicory
Chickweed
Clover
Cockspur
Dandelion
Dock
Galinsoga
Ground Ivy
Hawkweed
Heal All
Henbit
Knotweed
Lamb's-quarter
Lespedeza
Mallow
Morning Glory
Mustard
Orals
Peppergrass
Pigweed
Plantain
Poison Ivy
Poison Oak
Purslane
Ragweed
Sheep Sorrel
Shepherd's Purse
Speedwell
Tansy
Thistle
Spurge
Vetch
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow
and many other broadleaf lawn weeds

GUARANTEED ANALYSIS

Total Nitrogen (N) 15.00%
15% Urea Nitrogen
Derived from Urea

No Spreading — No Complicated Mixing
Just attach to a garden hose & spray

Where to Use: Liquid Weed & Feed is especially formulated for use on sensitive southern grasses - St. Augustine, bentgrasses, bahia, bermuda, and centipede - as well as on bluegrass, fescue, rye or Zoysia lawns. In lawns containing sensitive grasses, use Liquid Weed & Feed at the 1 quart for 5,000 sq. ft. rate. For lawns with non-sensitive grasses, use Liquid Weed & Feed at the 1 quart for 3,500 sq. ft. rate. Liquid Weed & Feed is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheepsorrel and spurge. Liquid Weed & Feed also controls the easy-to-kill weeds such as chickweed, clover, dandelion, knotweed, and plantains. In addition, Liquid Weed & Feed also feeds your lawn with a 15% urea nitrogen. For best results, spraying should be done when weeds are young and tender. Older weeds are more difficult to kill and may require a retreatment in 30 days to achieve complete control.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or sods down in the bare spots.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE:

SHAKE WELL BEFORE USING.

Connect spray nozzle to Garden hose, remove safety cap from container, attach container to end of nozzle. Turn on water, place finger over hole on right side of nozzle and proper ratio of LIQUID WEED & FEED will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of this liquid has been used.)

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.

B13 1185
EPA REG. NO. 2217-716
EPA EST. NO. 2217-KS-1

BEST AVAILABLE COPY