TRIMEC® 901

LIQUID WEED & FEED For Sensitive Grasses

20-0-0

NET CONTENTS ONE-HALF GALLON (NET WEIGHT: 4 LBS. 12 OZ.)

-- KEEP FROM FREEZING --

ONE-HALF GALLON COVERS UP TO 5,000 SQ. FT.

CONTROLS: Dandelion, Chickweed, Knotweed, Henbit, Plantains. Clover, Spurge and many other broadleaf lawn weeds; some of which are shown on this label.



No Spreading -- No Complicated Mixing Just attach to a garden hose & spray

ACTIVE II	IGRED	IENTS:
------------------	-------	--------

ACTIVE INQUIEDIENTS.	
*Dimethylamine Salt of 2,4-dichlorophenoxyacetic	
acid	2.784%
"Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)	
propionic acid	1.336%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic	
acid)	. 0.312%
INERT INGREDIENTS	95.568%
TOTAL	100 000%

THIS PRODUCT CONTAINS:

- 10 220 lbs. 2,4-dichlorophenoxyacotic acid per gallon or 2 313%. 110 105 Rs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gatton or 1 104% 110 024 Rs. 3,6-dichloro-o-sinsic acid per gatton or 0 252%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention

See side panel for additional precautionary statements.

TRIMECS is a registered frademark of PSI/Gordon Corporation 2.2.2

703/887 EPA REG. NC 2217 725 EPA EST. NO. 2217-KS-1



MEASURING CUIDE

1,000 Sq. Ft

COVERAGE

3 000 Sa Ft

COVERAGE

5 000 Sq. Ft

COVERAGE

WHEN EMPTY

PRECAUTIONARY STATEMENTS

READ THE ENTIRE LABEL FIRST, DISSERVE ALL PRECAUTIONS AND

POLLOW INTECTIONS CAREFULLY.

Hazards to Humana and Demostic Animals: CAUTION; Harmful if swallowed. Avoid contact with eyes, skin or clothero. In case of contact, immediately flush cues or skin with plenty of water. If erritation develops get medical attention. Avoid contamination of food or feedstuffs

Environmental Mazards: Do not apply directly to water. Do not apply when weather conditions lavor drift from target area. Do not contaminate water by cleans; of equipment or disposal of wastes Do not contaminate domestic or irrigation waters

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE: Keep from freezing. Store in original container in a locked storage area maccessible to children. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near Other ferlifizers, insecticides, fungicides, or seeds PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in Irash. CONTAINER DIGPOSAL: Do not reuse empty container Rinse thoroughly before discarding in trask

OTHER USE PRECAMOORS:

Do not apply this product through any type of irrigation system. Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85 degrees F. Seed can be safely sown 3-4 weeks. after application. Do not use on Carpetgrass, Dichondra, St. Augustine, Centigede, closely-mowed Bentorass, nor on tawns or turt where desirable clovers are sessent. Do not spray roots of trees and ornamentals Do not exceed specified dosages for any area. Be particularly careful within drighne of tree and other prinamental species

WEED\$		
Bedstraw	Henbit	Sheepsorret
Black Medic	Knotweed	Shepherdspur
Buckhorn	Lambsquarler	Speegwell
Buli Thisne	Lespedeza	Tansy Mustard
Burdock	Mallow	Thistie
Buttercup	Morningglory	Spurge
Chicory	Mustard	Velvetteal
Chickweed	Oxaks	Wild Carrol
Clover	Peopergrass	Wild Gartic
Cocklebur	Pigweed	Wold Lettuce
Dandelron	Plantain	Wild Onion
Doc	Poison fry	Yarrow
Gannsoga	Porson Oak	and many
Ground Ivv	Perstane	other broadles
Hawkweed	Requeed	fawn weeds
Mesi All	,	34

29-9-9 **GUARANTEED AMALYSIS** Total Nitrogen (N) 20% Urea Nitroger Derived from Urea

20 00%

WHERE TO USE:

TRIMEC* 901 may be used on Bahia, Bermuda, Bluegrass, Foscue, Plys or Zoysia lawns, vacant lots, playgrounds and other similar areas where broadlest weeds are present. This product is especially effective on all hard-to-kill weeds ruch as Poison by Poison Dak. Sheepsorrel and Spurge in addition TRIMEC. \$81 controls most other broadlesf weeds, including sensitive weeds such as Dirickweed. Clover, Dandelion. Knotweed and Plantain

WHEN TO USE:

Spray at any time during the graning second when weeds are actively prowing. (I'very lawns, wait until the groot has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks heliore pulling new grass seed or stolens down in the bare spors. Poor weed control may result if apray is applied during drought or just before rain. For best results, mow 1 to 2 days before treatme. Do not water from within 24 hours after treatment

Shake Well Before Using.

(One of the following will be used depending on the hose and applicator used)

INSTRUCTIONS

- Connect spray nozzle to garden hose.
- Shake conterns of bottle well.
- Remove safety cap.
- Attach tube to nozzle applicator.
- Attack bottle to open and of spray nozzle. Turn bottle counter clockwise until threads mesh. Then turn the bottle cleckwise until tightly closed
- 6. Turn on water; place linger over hele on right side of nozzle and proper ratio of TRIMEC* 901 will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of this liquid has been used 1
- Connect spray nozzle to garden hose; remove safety cap from container, attach tube to applicator and insert tube section. Attach container to end of nozzle
- · Turn on water; place linger over hole on right side of nozzle and proper ratio of TRIMEC* 981 will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring made to see how much liquid has been used)
- Altach garden hose to connector en dispenser
- Insert screwdriver into slot in orange lock ring and twist, snapping lock ring in two
- Remove both parts of separated tock ring
- Turn arrow on activator switch to OFF position
- Pirsh activator switch securate down against top of sprayer, which will pierce the safety coal inside the dispenser
- To begin soraying, form on welfer and then move activator switch to the ON position
- 7. To stop dispensing the product, turn off water

- Attach spray nozzle to garden hose. Turn nozzle clockwise to firmly attach it to hose
- Remove can from bottle (directions on can) and screw to a into spray nozzle
- 3. Turn the on off swrich on top of the spray nozzle counter clockwise to the "OFF" position Then turn on the water at the faucet.
- 4. To begin spraying, turn the sprayer nozzle switch clockwise to the "ON" position. The water automatically threes with the product. Walk at a steady pace spraying in a side to side art. slightly overlapping freated areas. Refer to graduated scale on side of container to outde coverage
- 5. To stop spraying, turn the sprayer nozzle switch counter clockwise to the "OFF" position" and turn off water at the caucal
- This pation package is intended as a refill for the one-half gallon hose-end soraver approach; container of TRMMEC* 981. However, this product may be used in any hose-end sprayer in which herbicides have been used

Add TRHREC* 961 to the hose-and sprayer jar. DO NOT ADD WATER TO THE JAR. One-half gallon covers 5,000 square feet. One gallon covers up to 10,000 square feet

LIGHTED WARRANTY. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to latel directions. Because of underly varying use conditions it is impossible to elemente all risks even when label directions are followed

EXCLUSIONS OF OTHER WANNAITIES AND REMEDIES: Except where such disclaimers and exclusions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABIL-ITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFAC-TURER, and, woon Menulacturer's compliance with said firmled warranty. BUYER SHALL HAVE NO REMEDY AGAINST MARIUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS and IN NO EVENT SHALL MANUFACTURER DE LIABLE FOR CONSEQUENTIAL DAMAGE OR

feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1.000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area."

OPTION (B)

"Trimec 901 should be supplemented by Liquid Lawn Fertilizer every 3 weeks from March through September (or April through October depending upon the growing season in your area). Please consult your local extension office for questions regarding frequency of this product in your area."

