

Reg # 2217-727 PM-23 192

TRIMEC®

902

LIQUID WEED & FEED For Sensitive Grasses

20-0-0

NET CONTENTS ONE-HALF GALLON
(NET WEIGHT: 4 LBS. 12 OZ.)

— KEEP FROM FREEZING —

ONE-HALF GALLON COVERS UP TO 5,000 SQ. FT.

CONTROLS: Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds; some of which are shown on this label.

The fast, easy way to weed control
and supplement your lawn.

No Spreading — No Complicated Mixing
Just attach to a garden hose & spray

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	1.872%
**Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.555%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.229%
INERT INGREDIENTS	97.343%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

*0.1465 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 1.547%
**0.0425 lbs. 2,4-dichlorophenoxyacetic acid per gallon or 0.449%
***0.0180 lbs. 3,6-dichloro-o-anisic acid per gallon or 0.190%

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See side panel for additional precautionary statements.

TRIMEC® is a registered trademark of PBI Gordon Corporation

7J4 887

EPA REC. NO. 2217-727

EPA EST. NO. 2217-KS-1

Mfd. by  **pbi/gordon**
corporation
KANSAS CITY, KANSAS 64116

MEASURING
GUIDE

1,000 Sq. Ft.
COVERAGE

3,000 Sq. Ft.
COVERAGE

5,000 Sq. Ft.
COVERAGE
WHEN EMPTY



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals:

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Do not apply this product through any type of irrigation system. Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Seed can be safely sown 3-4 weeks after application. Do not use on Carpetgrass, Dichondra, or on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified usages. Be particularly careful within drip-line of trees. This product may be injurious to St. Augustine, Bent, and Centipede grasses under conditions of high temperature and moisture. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage may occur.

WEEDS

Bedstraw	Henbit	Sheep sorrel
Black Medic	Knotweed	Sheepspurge
Buckhorn	Lambsquarters	Speedwell
Bull Thistle	Lespedeza	Tansy Mustard
Burdock	Mallow	Thistle
Buttercup	Morningglory	Spurge
Chicory	Mustard	Velvetleaf
Chickweed	Oxalis	Wild Carrot
Clover	Peppergrass	Wild Garlic
Cocklebur	Pigweed	Wild Lettuce
Dandelion	Plantain	Wild Onion
Dock	Poison Ivy	Yarrow
Galinsoga	Poison Oak	and many
Ground Ivy	Purslane	other broadleaf
Hawkweed	Ragweed	lawn weeds
Heal-All		

20-0-0

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.00%
20% Urea Nitrogen	
Derived from Urea	

WHERE TO USE:

TRIMEC® 902 is especially formulated for sensitive southern grasses — St. Augustine, Bentgrass, Bahia, Bermuda, and Centipede grasses in lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. TRIMEC® 902 is especially effective on all hard-to-kill weeds such as Poison Ivy, Poison Oak, Sheep sorrel and Spurge. TRIMEC® 902 also controls the easy-to-kill weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantains. In addition TRIMEC® 902 also feeds your lawn with a 20% urea nitrogen. For best results, spraying should be done when weeds are young and tender. Older weeds are harder to kill.

WHEN TO USE:

Spray at any time during the growing season when weeds are actively growing. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results, mow 1 to 2 days before treating. Do not water lawn within 24 hours after treatment.

HOW TO USE:

Shake Well Before Using

(One of the following will be used depending on the hose end applicator used.)

(A)

INSTRUCTIONS

1. Connect spray nozzle to garden hose.
2. Shake contents of bottle well.
3. Remove safety cap.
4. Attach tube to nozzle applicator.
5. Attach bottle to open end of spray nozzle. Turn bottle counter clockwise until threads mesh. Then turn the bottle clockwise until tightly closed.
6. Turn on water, place finger over hole on right side of nozzle and proper ratio of TRIMEC® 902 will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of this liquid has been used.)

(B)

Connect spray nozzle to garden hose; remove safety cap from container; attach tube to applicator and insert tube section. Attach container to end of nozzle.

Turn on water, place finger over hole on right side of nozzle and proper ratio of TRIMEC® 902 will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much liquid has been used.)

(C)

1. Attach garden hose to connector on dispenser.
2. Insert screwdriver into slot in orange lock ring and twist, snapping lock ring in two.
3. Remove both parts of separated lock ring.
4. Turn arrow on activator switch to OFF position.
5. Push activator switch securely down against top of sprayer, which will pierce the safety seal inside the dispenser.
6. To begin spraying, turn on water and then move activator switch to the ON position.
7. To stop dispensing the product, turn off water.

(D)

1. Attach spray nozzle to garden hose. Turn nozzle clockwise to firmly attach it to hose.
2. Remove cap from bottle (directions on cap) and screw bottle into spray nozzle.
3. Turn the on/off switch on top of the spray nozzle counter clockwise to the "OFF" position. Then turn on the water at the faucet.
4. To begin spraying, turn the spray nozzle switch clockwise to the "ON" position. The water automatically mixes with the product. Walk at a steady pace spraying in a side to side arc, slightly overlapping treated areas. Refer to graduated scale on side of container to guide coverage.
5. To stop spraying, turn the spray nozzle switch counter clockwise to the "OFF" position and turn off water at the faucet.

(E)

This one gallon package is intended as a refill for the one-half gallon hose-end. Product may be used in any hose-end sprayer in which herbicides have been used.

Add TRIMEC® 902 to the hose-end sprayer jar. **DO NOT ADD WATER TO THE JAR.** One-half gallon covers 5,000 square feet. One gallon covers up to 10,000 square feet.


LIMITED WARRANTY. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions it is impossible to eliminate all risks even when label directions are followed.

EXCLUSIONS OF OTHER WARRANTIES AND REMEDIES: Except where such disclaimers and exclusions are specifically prohibited by applicable law, THE FOREGOING IS THE ONLY WARRANTY OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area."

OPTION (B)

"Trimec 902 should be supplemented by Liquid Lawn Fertilizer every 3 weeks from March through September (or April through October depending upon the growing season in your area). Please consult your local extension office for questions regarding frequency of this product in your area."



SEP 22 1987