

1004

PM 23

2217-729

MM885

For the control of broadleaf weeds in bluegrass, fescue, Bermudagrass and bentgrass turf.

--KEEP FROM FREEZING--

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-methyl-4-chlorophenoxyacetic acid....	38.28%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic....	25.20%
INERT INGREDIENTS.....	36.52%
	TOTAL 100.00%

ACCEPTED  
 AUG 13 1987  
 Federal Insect  
 2217-729

THIS PRODUCT CONTAINS:

\* 3.0 lbs 2-methyl-4-chlorophenoxyacetic acid per gallon or 31.25%  
 \*\* 2.0 lbs 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 20.83%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

IF IN EYES: In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.

IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air and apply respiration if indicated.

See additional precautionary information elsewhere on this label.

NET CONTENTS

Mfd by. FRI/GORDON CORPORATION  
 Kansas City, Kansas 66118

EPA REG NO. 2217-XXXX  
 EPA EST NO.

STOP!

READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.




PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals:

DIANGER: Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift away from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.



DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL


STORAGE: Store in original container in a locked storage area. Keep from freezing. To avoid cross-contamination, do not store near other pesticides, fertilizers, or seeds.

PRODUCT DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

USE PRECAUTIONS:

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near non-target plants. Do not use on warm season grasses, such as St. Augustine, Centipede, Dichondra, or on lawns and turf where desirable clovers are present. NOTE: Coarse sprays are less likely to wind-drift. Do not spray when air temperature exceeds 85 degrees F. Seed can be safely sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. After using this product, clean sprayer thoroughly with soap or detergent and water and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.



GENERAL INFORMATION AND INSTRUCTIONS:

This product is for the control of broadleaf weeds in bluegrass, fescue, bermudagrass, and bentgrass turf. Do not use this product on closely-mowed bentgrass or bermudagrass as on golf greens or tees. Weeds are best controlled when they are young and actively growing.

WEEDS CONTROLLED:

Susceptible -- Dandelion, clover, common chickweed, mouse ear chickweed, plantain, pigweed, ragweed and lambsquarter. Moderately Susceptible -- Knotweed, black medic, English daisy, dock, purslane, oxalis, mallow.

APPLICATION RATE:

For established turf, apply 1 oz. to 1 1/4 oz. for 1,000 sqft. (2 2/3 pints to 3 1/3 pints per acre) in 1 to 2 gallons of water per 1,000 sqft. (30 to 60 gallons per acre). Use higher rates when hard-to-kill weeds are present. Maximum control of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid applying during periods when turf is under stress from drought or extreme heat. Do not water for 24 hours after application. Do not apply to newly seeded turf until after the second or third mowing.

NOTE: Use application equipment that is capable of applying a uniform droplet over the entire area to be covered.

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

