

PM 23

2217-736

10471

MAR 26 1996

James A. Armbruster, Ph.D.
 PBI/Gordon Corporation
 P.O. Box 14090
 1217 West 12th Street
 Kansas City, MO 64101

Dear Dr. Armbruster:

Subject: Revised Labeling - Add WPS Exclusionary Statement;
 Update Precautionary Language and Environmental Hazards
 Trimec M 907 Liquid Weed and Feed
 EPA Registration No. 2217-736
 Your Submission Dated August 24, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

- 1) Add an oral statement to the Statement of Practical Treatment to read "IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting."
- 2) Revise the beginning of the Precautionary Statements to read "Causes moderate eye injury. Harmful if swallowed or absorbed through skin."
- 3) Add the statement "Wash thoroughly with soap and water after handling" to the Precautionary Statements.

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the referenced label, incorporating the above changes, before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller
 Product Manager (23)
 Fungicide-Herbicide Branch
 Registration Division (7505C)

Enclosure

DK: 305-7546-FAB/PM23							
CONCURRENTS							
SYMBOL ▶	7505C						
SURNAME ▶	D. KENNY						
DATE ▶	3/25/96						

2017

TRIMEC® M 907 LIQUID WEED & FEED FOR SENSITIVE GRASSES 20-0-0

KEEP FROM FREEZING

ONE-HALF GALLON COVERS UP TO 5,000 SQUARE FEET.

Controls Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds; some of which are shown on this label.

The fast, easy way to weed control and supplement your lawn.

No Spreading - No Complicated Mixing
Just attach to a garden hose and spray.

ACTIVE INGREDIENTS:

* Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	1.863%
** Dimethylamine Salt of 2-methyl-4-chlorophenoxyacetic acid.....	1.029%
*** Dimethylamine Salt of Dicamba: 3,6-dichloro-o-anisic acid.....	0.205%
INERT INGREDIENTS.....	96.903%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- *0.1465 lbs 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.54%.
- **0.0815 lbs. 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 0.84%.
- ***0.0163 lbs 3,6-dichloro-o-anisic acid equivalent per gallon or 0.17%.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water.

For eyes, flush with water for 15 minutes and get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

See back panel for additional Precautionary Statements

NET CONTENTS ONE-HALF GALLON (NET WEIGHT 4 LBS 12 OZ.)

706/ APXXXXXX
EPA REG NO 2217-736
EPA EST NO 2217-KS-1
Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 26 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
2217-736

397

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear long pants, long-sleeved shirts and shoes when applying this product. Wear goggles when applying this product. ~~In case of contact, wash skin with plenty of soap and water; for eyes, flush eyes with plenty of water and get medical attention.~~

Environmental Hazards: Drift or run-off may adversely affect nontarget plants. ~~For terrestrial uses, Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.~~ Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift from target area. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not reenter or permit others to reenter treated areas until sprays have dried or dusts have settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE & DISPOSAL

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. Avoid contamination of food or feedstuffs

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application. Do not use on carpetgrass, dichondra, or on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages. Be particularly careful within drip line of trees. This product may be injurious to St. Augustinegrass, bentgrass, and centipedegrass under conditions of high temperature and moisture. Under these conditions, or in cases of accidental overdose, some slight discoloration of the St. Augustinegrass may occur.

4/07/17

WEEDS CONTROLLED:		
BEDSTRAW	HEAL-ALL	RAGWEED
BLACK MEDIC	HENBIT	SHEEP SORREL
BUCKHORN	KNOTWEED	SHEPHERDSPURSE
BULL THISTLE	LAMBSQUARTERS	SPEEDWELL
BURDOCK	LESPEDEZA	SPURGE
BUTTERCUP	MALLOW	TANSY MUSTARD
CHICORY	MORNINGGLORY	THISTLE
CHICKWEED	MUSTARD	VELVETLEAF
CLOVER	OXALIS	WILD CARROT
COCKLEBUR	PEPPERGRASS	WILD GARLIC
DANDELION	PIGWEED	WILD LETTUCE
DOCK	PLANTAIN	WILD ONION
GALINSOGA	POISON IVY	YARROW
GROUND IVY	POISON OAK	and other broadleaf
HAWKWEED	PURSLANE	lawn weeds.

20-0-0 GUARANTEED ANALYSIS	
Total Nitrogen(N).....	20.00%
20% Urea Nitrogen	
Derived from Urea	

(OPTION 1 OR 2 TO BE USED DEPENDING ON THE SUBREGISTRATION)

OPTION 1

A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed, this product supplies 0.19 pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local Agricultural Extension Agent for the proper amount of nitrogen to be applied in your area.

OPTION 2

Trimec® M907 Liquid Weed and Feed should be supplemented by Liquid Lawn Fertilizer every 3 weeks from March through September (or April through October depending upon the growing season in your area). Please consult your local extension office concerning a regular fertilizer program.

5011

WHERE TO USE: Trimec® M907 Liquid Weed and Feed is especially formulated for sensitive southern grasses - St. Augustinegrass, Bentgrass, Bahia, Bermuda and Centipede grasses, but can also be used on Bluegrass, Fescue, and other northern grasses in lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as Poison Ivy, Poison Oak, Sheep sorrel and Spurge. Trimec® M907 Liquid Weed and Feed also controls the easy to kill weeds such as Chickweed, Clover, Dandelion, Knotweed, and Plantains. In addition, Trimec® M907 Liquid Weed and Feed also feeds your lawn with a 20% nitrogen. For best results, spraying should be done when weeds are young and tender. Older weeds are harder to kill.

WHEN TO USE:

1. Spray during the growing season when weeds are active growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks after application for seeding and sprigging.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE: Shake well before using.

(One of the following will be used depending on the hose-end applicator used).

OPTION (A)

INSTRUCTIONS:

1. Connect spray nozzle to garden hose.
2. Shake contents of bottle well
3. Remove safety cap
4. Attach tube to nozzle applicator
5. Attach bottle to open end of spray nozzle. Turn bottle counter-clockwise until threads mesh. Then turn bottle clockwise until tightly closed.
6. Turn on water; place finger over hole on right side of nozzle and proper ratio will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of this liquid has been used.)

OPTION (B)

Connect spray nozzle to garden hose, remove safety cap from container, attach tube to applicator and insert tube section. Attach container to end of nozzle

Turn on water, place finger over hole on right side of nozzle and proper ratio will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much liquid has been used.)

60217

OPTION (C)

1. Attach garden hose to connector on dispenser.
2. Insert screwdriver into slot in orange lock ring and twist, snapping lock ring in two
3. Remove both parts of separated lock ring
4. Turn arrow on activator switch to OFF position.
5. Push activator switch securely down against top of sprayer, which will pierce the safety seal inside the dispenser.
6. To begin spraying, turn on water and then move activator switch to the ON position
7. To stop dispensing the product, turn off water.

OPTION (D)

1. Attach spray nozzle to garden hose. Turn nozzle clockwise to firmly attach it to hose.
2. Remove cap from bottle (directions on cap) and screw bottle into spray nozzle.
3. Turn the ON/OFF switch on top of the spray nozzle counter-clockwise to the OFF position. Then turn on the water at the faucet.
4. To begin spraying, turn the sprayer nozzle switch clockwise to the ON position. The water automatically mixes with the product. Walk at a steady pace spraying in a side to side arc, slightly overlapping treated areas. Refer to graduated scale on side of container to guide coverage.
5. To stop spraying, turn the sprayer nozzle switch counter-clockwise to the OFF position and turn off water at the faucet

OPTION (E)

This gallon package is intended as a refill for the one-half gallon hose-end sprayer applicator/container of Trimec® M907 Liquid Weed and Feed. However, this product may be used in any hose-end sprayer in which herbicides have been used.

Add Trimec® M907 Liquid Weed and Feed to the hose-end sprayer jar. DO NOT ADD WATER TO THE JAR. One-half gallon covers 5,000 square feet. One gallon covers up to 10,000 square feet.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

7/10/17

MEASURING GUIDE

1,000 SQ FT

COVERAGE

3,000 SQ.FT.

COVERAGE

5,000 SQ.FT.

COVERAGE
WHEN EMPTY