



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

February 8, 2024

Eric D. Smith
Director, Regulatory Affairs
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, KS 66286

Subject: Registration Review Label Mitigation for Permethrin
Product Name: Gordon's Chemicals Permanone General Purpose Aqueous
Insecticide II
EPA Registration Number: 2217-781
Application Date: 1/22/2021
Decision Number: 569856

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Permethrin Interim Decision and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

Page 2 of 2
EPA Reg. No. 2217-781
Decision No. 569856

approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Carolyn Smith by phone at (202) 566-2273, or via email at smith.carolyn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Jaclyn Pyne". The signature is written in a cursive style and is centered below the word "Sincerely,".


Jaclyn Pyne, Team Leader
Risk Management and Implementation Branch 3
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

SUBLABEL 1: For Residential Users

GORDON'S CHEMICALS PERMANONE® GENERAL PURPOSE AQUEOUS INSECTICIDE II

EPA Reg. No. 2217-781



ACTIVE INGREDIENT:

*Permethrin	0.50%
INERT INGREDIENTS:	99.50%
TOTAL	100.00%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans. PERMANONE® is a registered trademark of Bayer AG.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

For both indoor and outdoor use. Application of product on pets (dogs) must only be done indoors.



Do not allow to enter indoor or outdoor drains. *No permita la entrada a desagües internos o externos.*



[Note to reviewer: Pictogram must at least 0.25 in.². unless this size is greater than 10% of the label size]

SHAKE WELL BEFORE USING

PRECAUTIONS AND USE RESTRICTIONS:

Entry Restrictions:

- Except when applying to pets (dogs) or livestock, do not enter or allow adults, children, or pets to enter until sprays have dried.

Application Restrictions:

- Except when applying to dogs and to livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to dogs and to livestock, only protected handlers may be in the area during application. Do not breathe dusts, vapors, or spray mist.

When Used In Dairy Barns Or Facilities:

- Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Do not apply as a broadcast or spot-treatment to indoor surfaces at residential sites, including nurseries, day care center, schools, hospitals, and nursing homes.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Applications to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhand, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces such as (such as the side of a man-made structure), directly above impervious surfaces (e.g., driveways, sidewalks, etc.) up to 2 feet above ground level.
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation, only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only. Spot treatments must not exceed two square feet in size (for example, 2ft. by 1ft. or 4 ft. by 0.5 ft.). Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and crack-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants. Do not apply when the wind speed is greater than 15 mph.

Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly into sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Do not apply or irrigate to the point of run-off. Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application. Not for use in outdoor residential misting systems. For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

INDOOR APPLICATION:

FOR HOUSEHOLD PESTS:

Not for use in commercial food/feed processing, preparation, serving or storage areas.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Cover or remove exposed food.

Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications.

Do not use this product in or on electrical equipment due to the possibility of shock hazard. For spot application only. Do not use as a space spray indoors. Do not apply to rooms occupied by the elderly or infirm. Do not apply to classrooms when in use.

SURFACE SPRAYING:

To control cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spider, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, drugstore beetles, mealworms, grain mites and cadelles, direct the spray into hiding places, cracks and crevices, and behind shelves and drawers.

For silverfish, spray bookcases.

For ants, spray trails, nests, and points of entry.

To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices, and infested areas of shelving. Repeat treatment as needed.

To control bedbugs, spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Allow mattresses and other treated articles to dry thoroughly before use. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control fleas (adults and larvae) and ticks (adults and larvae), thoroughly spray infested areas, pet beds (only for dogs), resting quarters, nearby cracks and crevices, along and behind baseboard, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal (dogs only) quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets (dogs) to contact treated areas.

To control clothes moths, remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus, and other storage directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To control lice and louse eggs, spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture, that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK:

To protect cattle (beef and dairy), goats, sheep, hogs, and horses from horn flies, face flies, house flies, and gnats, apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control stable flies, horse flies and deer flies, apply at a rate of 2 fluid ounces per adult animal, sufficient to wet the hair. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control poultry lice, spray roosts, walls and nests, or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control bedbugs and mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS:

To control fleas, ticks, and lice, start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use waterproof gloves or mitts when applying product. Repeat as necessary.

Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals (dogs only). Sensitivities may occur after using ANY pesticide product for pets (dogs). If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately or call this Animal Poison Control Hotline, 1-800-345-4735.

SPECIAL ANIMALS:

Consult a veterinarian or call this Animal Poison Control Hotline, 1-800-345-4735 before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

ADVERSE REACTION INFORMATION:

Sensitivities may occur after using ANY pesticide product for pets (dogs) . If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately, or call this Animal Poison Control Hotline, 1-800-345-4735.

OUTDOOR APPLICATION:

PERIMETER TREATMENT TO CONTROL TICKS (INCLUDING TICKS WHICH MAY CARRY LYME DISEASE):

Use Permanone General Purpose Aqueous Insecticide II as a residual treatment to control ticks including the deer tick (*Ixodes dammini*), lone star tick (*Amblyomma americanum*), brown dog tick (*Rhipicephalus sanguineus*), American dog tick (*Dermacentor variabilis*), and Western black-legged tick (*Ixodes pacificus*).

For each area listed below, apply until surface is wet (approximately 1 gallon per 750 to 1,000 square feet).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 7 feet wide around and the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 to 7 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS:

For use on roses, dahlias asters and other ornamentals to control Japanese beetles, whiteflies, aphids, exposed thrips, spiders, red mites and leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews, and zinnias.

Effective when used as directed against whiteflies, aphids, spider and red mites, exposed thrips, lace bug, Japanese beetle, pavement ants, clover mites, and armyworms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants, and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry area.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Do not pour or dispose down-the-drain or sewer.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

SUBLABEL 2: For Occupational Users

GORDON'S CHEMICALS PERMANONE® GENERAL PURPOSE AQUEOUS INSECTICIDE II

EPA Reg. No. 2217-781

Commercial application: For use in non-food areas indoors and outside premises of motels, commercial buildings, schools, warehouses, office buildings, kennels, apartments and condominium buildings, and non-food contact areas of food-feed processing establishments.

Household application: Indoors- along baseboards, around cupboards

Domestic Animals: Dogs to control fleas & ticks. Poultry to control lice, fleas & ticks. Livestock - protects beef and dairy cattle, goats, sheep and horses from horn flies, face flies, mosquitoes and gnats

ACTIVE INGREDIENT:

*Permethrin	0.50%
INERT INGREDIENTS:	99.50%
TOTAL	100.00%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans. PERMANONE® is a registered trademark of Bayer AG.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Waterproof gloves

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched (except as required by directions for use) or heavily contaminated with this product's concentrate. Do not reuse them.

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

Under some conditions, it may also have a potential for transport into surface water in runoff (primarily adsorbed to suspended soil particles), for several months or more after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, and areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

For both indoor and outdoor use. Application of product on pets (dogs) must only be done indoors.

Do not allow to enter indoor or outdoor drains. *No permita la entrada a desagües internos o externos.*



[Note to reviewer: Pictogram must at least 0.25 in.². unless this size is greater than 10% of the label size]

SHAKE WELL BEFORE USING

PRECAUTIONS AND USE RESTRICTIONS:

Entry Restrictions:

- Except when applying to pets (dogs) or livestock, do not enter or allow adults, children, or pets to enter until sprays have dried.

Application Restrictions:

- Except when applying to dogs and to livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to dogs and to livestock, only protected handlers may be in the area during application. Do not breathe dusts, vapors, or spray mist.

When Used In Dairy Barns Or Facilities:

- Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Do not apply as a broadcast or spot-treatment to indoor surfaces at residential sites, including nurseries, day care center, schools, hospitals, and nursing homes.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Applications to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhand, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces such as (such as the side of a man-made structure), directly above impervious surfaces (e.g., driveways, sidewalks, etc.) up to 2 feet above ground level .
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation, only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only. Spot treatments must not exceed two square feet in size (for example, 2ft. by 1ft. or 4 ft. by 0.5 ft.). Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and crack-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants. Do not apply when the wind speed is greater than 15 mph.

Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly into sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Do not apply or irrigate to the point of run-off. Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application. Not for use in outdoor residential misting systems. For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

INDOOR APPLICATION:

FOR HOUSEHOLD PESTS:

Not for use in commercial food/feed processing, preparation, serving or storage areas.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications.

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

For spot application only. Do not use as a space spray indoors. Do not apply to rooms occupied by the elderly or infirm. Do not apply to classrooms when in use.

SURFACE SPRAYING:

To control cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spider, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, drugstore beetles, mealworms, grain mites and cadelles, direct the spray into hiding places, cracks and crevices, and behind shelves and drawers.

For silverfish, spray bookcases.

For ants, spray trails, nests, and points of entry.

To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices, and infested areas of shelving. Repeat treatment as needed.

To control bedbugs, spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Allow mattresses and other treated articles to dry thoroughly before use. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control fleas (adults and larvae) and ticks (adults and larvae), thoroughly spray infested areas, pet beds (only for dogs), resting quarters, nearby cracks and crevices, along and behind baseboard, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal (dogs only) quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets (dogs) to contact treated areas.

To control clothes moths, remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus, and other storage directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To control lice and louse eggs, spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a

distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture, that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK:

To protect cattle (beef and dairy), goats, sheep, hogs, and horses from horn flies, face flies, house flies, and gnats, apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control stable flies, horse flies and deer flies, apply at a rate of 2 fluid ounces per adult animal, sufficient to wet the hair. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control poultry lice, spray roosts, walls and nests, or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control bedbugs and mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS:

To control fleas, ticks, and lice, start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use waterproof gloves or mitts when applying product. Repeat as necessary.

Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals (dogs only). Sensitivities may occur after using ANY pesticide product for pets (dogs). If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately or call this Animal Poison Control Hotline, 1-800-345-4735.

SPECIAL ANIMALS:

Consult a veterinarian or call this Animal Poison Control Hotline, 1-800-345-4735 before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

ADVERSE REACTION INFORMATION:

Sensitivities may occur after using ANY pesticide product for pets (dogs) . If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately, or call this Animal Poison Control Hotline, 1-800-345-4735.

OUTDOOR APPLICATION:

PERIMETER TREATMENT TO CONTROL TICKS (INCLUDING TICKS WHICH MAY CARRY LYME DISEASE):

Use Permanone General Purpose Aqueous Insecticide II as a residual treatment to control ticks including the deer tick (*Ixodes dammini*), lone star tick (*Amblyomma americanum*), brown dog tick (*Rhipicephalus sanguineus*), American dog tick (*Dermacentor variabilis*), and Western black-legged tick (*Ixodes pacificus*).

For each area listed below, apply until surface is wet (approximately 1 gallon per 750 to 1,000 square feet).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 7 feet wide around and the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 to 7 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not pour or dispose down-the-drain or sewer. Call your local solid waste Agency for local disposal options.

[Plastic Containers – Nonrefillable with capacities equal to 1 gallon or less than 5 gallons]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- For domestic use both indoors and outdoors
- A ready-to-use PERMANONE® stable emulsion insecticide
- For control of fleas, brown dog ticks, and lice on premises and on dogs
- Controls fleas and ticks on dogs for 35 days
- Aids in control of ticks on dogs for an additional 2 weeks
- For use on livestock and on premises to control pests listed on back panel.

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00781.20230808.amend-proposed-clean.doc

2. Reason for Issue: Permethrin Reg. Review