	U.S. ENVIRONMENTAL PROTECTION	EPA Reg. Number:	Date of Issuance:
UNITED STATES	AGENCY		Date of issuance.
	Office of Pesticide Programs	2217-793	
	Registration Division (7505P)		
VIAL PROTECT	Ariel Rios Building		NOV 10 2009
	1200 Pennsylvania Ave., NW		
	Washington, D.C. 20460		
	NOTICE OF PESTICIDE:	Term of Issuance:	
	Registration	Num Charlette Da	dura i
	<u>X</u> Reregistration	Name of Pesticide Pro	
	(under FIFRA, as amended)	Gordon's Tri-0	Guard 20-10-5
	ess of Registrant (include ZIP Code):		
BI/Gordon Co		•	
217 West 12 th	•		
	issouri 64101 in labeling differing in substance from that acce		nan in Shiring a contract of
	st be submitted to and accepted by the Registrat n any correspondence on this product always re	めいしい いししゅう やくねる ひや テレジバスタングッグ	Charles Charles The Charles and the State of
n the hasis of	information furnished by the registrant, the abo	ve named pesticide	is hereby
	istered under the Federal Insecticide, Fungicide	-	-
	construed as an endorsement or recommendation		
	health and the environment, the Administrator,		
	el the registration of a pesticide in accordance		
	tion with the registration of a product under this		
	right to exclusive use of the name or to its use i	f it has been covere	d by others.
e registrant a	right to exclusive use of the name or to its use i reregistered in accordance with FIFRA provide		d by others.
e registrant a m nis product is 1) S		d that you: ion/reregistration re	view of your
e registrant a m nis product is 1) S F	reregistered in accordance with FIFRA provide Submit and/or cite all data required for registrat	d that you: ion/reregistration re	view of your
e registrant a m nis product is 1) S F	reregistered in accordance with FIFRA provider Submit and/or cite all data required for registration product when the Agency requires all registrants	d that you: ion/reregistration re s of similar product	view of your
e registrant a m nis product is 1) S F gnature of Ap	reregistered in accordance with FIFRA provider Submit and/or cite all data required for registration product when the Agency requires all registrants	d that you: ion/reregistration re s of similar product	view of your
e registrant a m nis product is 1) S F gnature of Ap anne Miller	reregistered in accordance with FIFRA provider Submit and/or cite all data required for registration product when the Agency requires all registrants proving Official: $\int \int $	d that you: ion/reregistration re s of similar product	view of your
e registrant a m nis product is 1) S F	reregistered in accordance with FIFRA provider Submit and/or cite all data required for registration product when the Agency requires all registrants proving Official: $\int S \int$ er 23	d that you: ion/reregistration re s of similar product	view of your

.

Page 2 EPA Reg. 2217-793

2) Based on toxicity ranking per the acute toxicity review, the First Aid statements should be placed on the label in the following order:

"IF IN EYES:... IF ON SKIN OR CLOTHING:... IF INHALED:... IF SWALLOWED:..."

3) Per the acute toxicity review, the Hazards to Humans and Domestic Animals must be revised to read:

"CAUTION

Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust."

4) The first sentence of the Environmental Hazard section must be revised to read "This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants."

5) Delete "General" from "General Precautions".

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supercedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

GORDON'S TRI-GUARD 20-10-5

EPA Reg. No. 2217-793

ACTIVE INGREDIENT:	
2,4-dichlorophenoxyacetic acid	0.62%
(+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	
Dicamba (3,6-dichloro-o-anisic acid)	0.06%
INERT INGREDIENTS:	99.18%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

12.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.

2.8 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.

1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

TRIMEC® is a registered trademark of PBI/Gordon Corporation. Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- · long-sleeved shirt and long pants,
- shoes plus socks,
- chemical-resistant gloves

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry



1 of 6 002217-00793.20081105.proposed.doc

ACCEPTED with COMMENTS In EPA Letter Dated: NOV 1 0 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
	t container or label with you when calling a poison control center or doctor or going for additional information in case of emergency, call toll free 1-877-800-5556.

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not enter or allow people (or pets) to enter the treated area until dusts have settledoc
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- · For use on ornamental lawns and turf established in institutional, ornamental, and residential sites.
- · Grass seed crops and sod farms uses are excluded.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE:

Gordon's Tri-Guard is a Trimec® brand herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Gordon's Tri-Guard is intended to be applied by noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example, hospitals, nursing homes, schools, museums, libraries, sport facilities (including golfcourses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Gordon's Tri-Guard may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Gordon's Tri-Guard may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area. Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

Do not use this product on dichondra, carpetgrass, St. Augustinegrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with Gordon's Tri-Guard. A partial list of susceptible weeds is shown below:

Bedstraw	Clover	Knotweed	Purslane
Black medic	Dandelion	Lambsquarters	Ragweed
Buckhorn	Dock	Morningglory	Sheep sorrel
Burdock	Ground ivy	Peppergrass	Shepherdspurse
Chicory	Healall	Pigweed	Speedwell
Chickweed	Henbit	Plantain	Thistle

HOW MUCH TO APPLY:

For Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Bermudagrass, and Zoysiagrass: Apply 132.0 to 158.4 pounds (lbs.) of Gordon's Tri-Guard per acre of established turfgrass. Or the equivalent application rates are 3.0 to 3.6 pounds of product per 1,000 square feet (sq.ft.) of turfgrass.

The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application (0.98 Ibs 2,4-D ae, 0.22 Ib MCPP-p ae and 0.10 Ib dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. (1.96 Ibs 2,4-D ae, 0.42 Ib MCPP-p ae and 0.20 Ib dicamba ae per acre).

A sequential or follow-up application may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling.]

Bag Size,	Desired Rate	Desired Rate Per	Treated Area For This Bag Size	
Lbs.	Per Acre, Lbs.	1,000 Sq. Ft., Lbs.	Acres	Sq. Ft.
15	132.0	3.0		5,000
	140.5	3.2		4,700
	158.4	3.6		4,200
16	132.0	3.0		5,300
	140.5	3.2		5,000
	158.4	3.6		4,400
17	132.0	3.0		5,700
	140.5	3.2		5,300
	158.4	3.6		4,700
18	132.0	3.0		6,000
	140.5	3.2		5,600
	158.4	3.6	· · ·	5,000
20	132.0	3.0		6,700
	140.5	3.2	·	6,300
	158.4	3.6		5,600
30	132.0	3.0	0.23	10,000
	140.5	3.2	0.22	9,400
	158.4	3.6	0.19	8,300
35	132.0	3.0	0.27	11,700
	140.5	3.2	0.25	10,900
	158.4	3.6	. 0.22	9,700
36	132.0	3.0	0.28	12,000
	140.5	3.2	0.26	11,300
	158.4	3.6	0.23	10,000
40	132.0	3.0	0.31	13,300
	140.5	3.2	0.29	12,500
	158.4	3.6	0.25	11,100
45	132.0	3.0	0.34	15,000
	140.5	3.2	0.32	14,100
	158.4	3.6	0.29	12,500
48	132.0	3.0	0.37	16,000
	140.5	3.2	0.34	15,000
	158.4	3.6	0.31	13,300
50	132.0	3.0	0.38	16,700
	140.5	3.2	0.36	15,600
-	158.4	3.6	0.32	13,900
55	132.0	3.0	0.42	18,300
	140.5	3.2	0.39	17,200
	158.4	3.6	0.35	15,300
60	132.0	3.0	0.46	20,000
	140.5	3.2	0.43	18,800
	158.4	3.6	0.38	16,700

Optional: Equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels as requested by supplemental registrants.

APPLICATION SUGGESTIONS:

- 1. Spring and fall applications are recommended for the best control. However, Gordon's Tri-Guard may be applied any time broadleaf weeds are present and actively growing.
- 2. Mow the turfgrass 2 to 3 days before application.
- 3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
- 4. Schedule applications in the morning when dew is present. Or sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
- 5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

GUARANTEED ANALYSIS 20-10-5			
Total Nitrogen (N)	20%		
Water Insoluble Nitrogen (N)	5%		
Available Phosphate (P2O5)	10%		
Soluble Potash (K2O)	5%		

Fertilizer Labeling Requirements:

 Information regarding the contents and levels of metals in this product is available on the Internet at http://www.xxxxx.com.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Paper and Plastic Bags - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State and local authorities by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- · See next panel for additional Precautionary Statements and First Aid
- Net Weight: _____
- EPA Est. No. ____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

• Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5, and K2O include the following:

12-4-4	20-10-5	23-3-3	27-3-5
16-2-0	.21-3-5	23-3-4	28-3-7 ·
18-4-8	21-5-7	23-3-5	30-3-5
19-3-12	22-3-4	25-3-5	33-3-5

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

- Fifty pounds covers up to 16,700 square feet of turf.
- Twenty pounds covers up to 6,700 square feet.
- Provides excellent response to the fertilizer plus effective broadleaf weed control.
- Combines the advantages of fertilization plus a broad range of weed control.
- Provides controlled release nitrogen feeding from Poly-S(TM) Technology.
- Previously approved retail container package sizes include: 15, 16, 17, 18, 20, 30, 35, 36, 40, 45, 48, 50, 55, and 60 pounds.
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, and other broadleaf weeds.

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00793.20081105.proposed.doc

2. Reason for Issue: Various changes pertaining to re-registration. Re-format label.