2217-795	02-24			78
	United States nental Protection Washington, DC 20460	\	Registration Amendmen Other	
	Application for I	Pesticide - Section	on I	
1. Company/Product Humber 2217-795		2. EPA Product Man- Joanne I	ager	3. Proposed Classification
4. Company/Product (Name) Gordon's Triplex 24-	-4-8	PM# Product Manag	er—Team 23	None Restricted
5. Name and Address of Applicant (Include PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 Check if this is a new address		1	similar or identical in	rith FIFRA Section 3(c)(3) n composition and labeling to:
Check it this is a new address		tion - II		
Amendment - Explain below. Resubmission in response to Agency letter Notification - Explain below. Explanation: Use additional page(s) if necess NOTIFICATION: We are requesting to accordance with Pesticide Registrate the label—no other changes have been this notification is consistent with the provisions labeling or Confidential Statement of Formula of the EPA. I further understand that if this notificate FIFRA and I may be subject to enforcement actions.	sary. (For section I and Section delete a single ground ion Notice (PRN) 98-10 made to the use directly of PR Notice 98-10 and EPA this product. I understand the lon is not consistent with the	Agency letter of "Me Too" Appli Other - Explain of users from the otions, sites, or rates at it is a violation of 18 U.S terms of PR Notice 98-10	cation. below. labeling of Gore enclosed label. of application. 2.46, and no other ch	This is the only change to manges have been made to the ully make any false statement to
	Sec	tion - III	· · · · · · · · · · · · · · · · · · ·	the state of the s
1. Material This Product Will Be Packaged In: Child-Resistant Packaging Yes* No * Certification must be submitted 3. Location of Net Contents Information Label Container 6. Manner in Which Label is Affixed to Product	No. per If "Yes Packa" 4. Size(s) Retail Contai	ge wgt. container	2. Type of Contain Metal Plastic Glass Paper Other (Sp. Location of Label Dir On Label On Labeling according accordi	pecify)rections
	Paper glued Stenciled			
	Sect	ion - IV		
Contact Point (Complete items directly below for Name	or identification of individual	to be contacted, if necessa		pplication.)

both under applicable law.

2. Signature

3. Title

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or

Director of Regulatory Services

Manager of Regulatory Services

5. Date

February 4, 2004

816-460-6287

6. Date Application

(Stamped)

Hnceⁱved

Jim Kunstman

Craig Martens

GORDON'S TRIPLEX 24-4-8

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, and other broadleaf weeds.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	0.69%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.15%
Dicamba: 3,6-dichloro-o-anisic acid	0.06%
INERT INGREDIENTS	99.10%
TOTAL	100 00%

THIS PRODUCT CONTAINS:

13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton

3.0 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton

1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements.

NET WEIGHT ___ POUNDS

793 AP020404 EPA REG. NO. 2217-795 EPA EST. NO. 2217-KS-X MANUFACTURED BY:

Conponation

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missoun 64101

Telephone 800-821-7925



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

If in eyes:	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.	
	• Remove contact lenses, if present, after the first 5 minutes, then continue	
	rinsing eye.	
	Call a poison control center or doctor for treatment advice.	
If on skin or	Take off contaminated clothing.	
clothing:	Rinse skin immediately with plenty of water for 15 - 20 minutes.	
_	Call a poison control center or doctor for treatment advice.	
If swallowed:	Call a poison control center or doctor immediately for treatment advice.	
	Have person sip a glass of water if able to swallow.	
	Do not induce vomiting unless told to do so by a poison control center or doctor.	
	Do not give anything by mouth to an unconscious person.	
If inhaled:	Move person to fresh air.	
	• If person is not breathing, call 911 or an ambulance, then give artificial	
	respiration, preferably by mouth-to-mouth, if possible.	
	Call a poison control center or doctor for treatment advice.	
Have the product container or label with you when calling a poison control center or doctor or going		
for treatment.	You may also call toll-free, 1-877-800-5556 for emergency medical treatment	
information.		

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPP-p and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPP-p and 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

DISPOSAL:

If empty - Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled - Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

WHERE TO USE:

Gordon's Triplex is a Trimec[®] brand herbicide containing fertilizer, 2,4-D, mecoprop, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Gordon's Triplex is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example areas around hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Gordon's Triplex may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Gordon's Triplex may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 - 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (½ of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Do not use this product on dichondra, carpetgrass, St. Augustinegrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with Gordon's Triplex. A partial list of susceptible weeds is shown below:

Bedstraw	Dock	Pigweed	
Black medic	Ground ivy	Plantain	
Buckhorn	Healall	Purslane	
Burdock	Henbit	Ragweed	
Chicory	Knotweed	Sheep sorrel	
Chickweed	Lambsquarters	Shepherdspurse	
Clover	Morningglory	Speedwell	-
Dandelion	Peppergrass	Thistle	

HOW MUCH TO APPLY:

For Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Bermudagrass, and Zoysiagrass: Apply 140.5 to 174.2 pounds (lbs.) of Gordon's Triplex per acre of established turfgrass. Or the equivalent application rates are 3.2 to 4.0 pounds of product per 1,000 square feet (sq.ft.) of turfgrass.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling.

Bag Size,	Desired Rate	Desired Rate Per	Treated Area F	or This Bag Size
Lbs.	Per Acre, Lbs.	1,000 Sq.Ft. , Lbs.	Acres	Sq.Ft.
	140.5	3.2		4,700
15	158.4	3.6		4,200
	174.2	4.0	~ ~ ~ ~	3,800
	140.5	3.2	****	5,000
16	158.4	3.6		4,400
	174.2	4.0	47 48 10 46 48	4,000
	140.5	3.2	<u> </u>	5,300
17	158.4	3.6		4,700
	174.2	4.0		4,300
111111111111111111111111111111111111111	140.5	3.2		5,600
18	158.4	3.6		5,000
	174.2	4.0		4,500
	140.5	3.2		6,300
20	158.4	3.6		5,600
	174.2	4.0		5,000
	140.5	3.2	0.22	9,400
30	158.4	3.6	0.19	8,300
	174.2	4.0	0.17	7,500
	140.5	3.2	0.25	10,900
35	158.4	3.6	0.22	9,700
	174.2	4.0	0.20	8,800
, , , , , , , , , , , , , , , , , , , ,	140.5	3.2	0.26	11,300
36	158.4	3.6	0.23	10,000
	174.2	4.0	0.21	9,000
	140.5	3.2	0.29	12,500
40	158.4	3.6	0.29	11,100
	174.2	4.0	0.23	10,000
	140.5	3.2	0.32	14,100
45	158.4	3.6	0.29	12,500
	174.2	4.0	0.26	11,300
	140.5	3.2	0.36	15,600
50	158.4	3.6	0.32	13,900
	174.2	4.0	0.29	12,500
	140.5	3.2	0.39	17,200
55	158.4	3.6	0.35	15,300
	174.2	4.0	0.32	13,800
	140.5	3.2	0.43	18,800
60	158.4	3.6	0.38	16,700
	174.2	4.0	0.34	15,000

Optional:

Equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels as requested by supplemental registrants.

APPLICATION SUGGESTIONS:

- 1. Spring and fall applications are recommended for the best control. However, Gordon's Triplex can be applied any time broadleaf weeds are present and actively growing.
- 2. Mow the turfgrass 2 to 3 days before application.
- 3. Water the turfgrass thoroughly (>½ inch) at least 1 to 2 days before application.
- 4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<¼ inch) before application to provide maximum control.
- 5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

GUARANTEED ANALYSIS 24-4-8	
Total Nitrogen (N)	24%
Water Insoluble Nitrogen (N)	5%
Available Phosphate (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%

Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State
 Department of Agriculture regarding the components in this product is available on the internet at
 http://www.wa.gov/agr/."

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

I. Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

10-6-4	19-5-9	22-7-7	25-3-8	
12-5-5	20-4-6	22-10-10	26-3-3	
12-4-4	20-4-8	23-3-3	26-3-5	
15-5-10	20-10-0	23-7-7	26-4-6	
16-2-3	20-10-5	24-3-6	28-3-3	
16-3-5	21-5-7	24-4-8	28-3-7	
16-6-9	21-3-5	24-5-10	30-3-5	
18-3-5	22-3-6	25-3-3	30-3-10	
18-4-8	22-6-6	25-3-5	30-4-8	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

GUARANTEED ANALYSIS		
Total Nitrogen (N)	19%	
5.6% Water Insoluble Nitrogen		
9.0% Ammoniacal nitrogen		
1.4% Urea Nitrogen*		
3.0% Water Soluble Organic Nitrogen*		
Available Phosphate (P ₂ O ₃)	5%	
Soluble Potash (K ₂ O)	9%	
Total Magnesium (MG)	1.0%	
0.2% Water Soluble Magnesium (Mg)	Į	
Total Sulfur (S)	8.0%	
8.0% Combined Sulfur (S)		
Total Iron (Fe)	1.0%	
0.1% Soluble Iron (Fe)	1	
Total Manganese (Mn)	0.5%	
0.01% Water Soluble Manganese (Mn)		
*Contains 3.0% Slowly Available Nitrogen from Methylene Urea.	l	
Nutrient Sources: Urea, Methylene Urea, Ammonium Phos		
Ammonium Sulfate, Sulfate of Potash, Dolomite, Sulfate of Potash,		
Magnesia, Iron Sulfate, Manganese Sulfate, Manganese Oxide.	j	
Chlorine (CI) not more than	3.0%	

- II. Previously approved retail container package sizes include: 15, 16, 17, 18, 20, 30, 35, 36, 40, 45, 50, 55, and 60 pounds.
- III. Advertising Claims: Advertising claims presented on the container (bag) that may or may not be used by supplemental registrants:
 - Fifty pounds covers up to 15,600 square feet of turf.
 - Twenty pounds covers up to 6,300 square feet.
 - Provides excellent response to the fertilizer plus effective broadleaf weed control.
 - Combines the advantages of fertilization plus a broad range of weed control.
 - ◆ Provides controlled release nitrogen feeding from Poly-S™ Technology.
 - Homogeneous fertilizer granules for high quality turf.
 - Slow release nitrogen for long lasting feeding without burning when applied as directed.
 - Continual root development for a thick lush lawn.
 - This fertilizer contains iron which will stain concrete and painted surfaces.