

2217-798

06/21/2011

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr. James Kunstman
Director of Regulatory Affairs
PBI/Gordon Corporation
1217 West 12th Street
P.O. Box 014090
Kansas City, MO 64101-0090

JUN 21 2011

Subject: Label Amendment - Changes to Signal Word, Precautionary Statements, & Marketing Claims
Product Name: EH-1153 Weed and Feed 21-3-5
EPA Registration Number: 2217-798
Decision Number: 445105
Submission Date: January 28, 2010

Dear Mr. Kunstman:

The label amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable.

Please note that marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available, then those claims must be removed.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Products shipped after 18 months from the date on this notice or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of this product constitutes acceptance of these conditions. This label supersedes all previously accepted labels. You must submit one copy of the final printed label before the product is released for shipment. One copy of the label stamped "Accepted" is enclosed for your records. If you have any questions, please contact Dianne Morgan by phone at (703) 305-6217 or via email at morgan.dianne@epa.gov.

SEE NEXT PAGE

Page 2 of 2
Label Amendment
Product Name: EH-1153 Weed and Feed 21-3-5
EPA Registration Number: 2217-798
Decision Number: 445105

Sincerely,



Kathryn V. Montague *for*
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

ACCEPTED
 JUN 21 2011
 2217-798
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No

EH-1153 WEED AND FEED 21-3-5

EPA Reg. No. 2217-798

ACTIVE INGREDIENT:

MCPA acid	0.69%
Mecoprop-p acid	0.15%
Dicamba acid	0.06%
INERT INGREDIENTS	99.10%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 13.8 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per ton, or 0.69%.
- 3.0 lb (+)-R-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton, or 0.15%.
- 1.2 lb 3,6-dichloro-o-anisic acid equivalent per ton, or 0.06%.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt, long pants, socks, and shoes when applying this product.

First Aid	
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p>	



4/11

Environmental Hazards

This pesticide may adversely affect non-target plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

[LABEL OPTION 1]

WHERE TO USE:

For residential use only:

EH 1153 Weed and Feed (EPA Reg. No. 2217-798) is intended to be used outside and around residences/ homes that are occupied by individuals including lawns, yards, turfgrass, and the immediate surroundings associated with households/or home life.

EH-1153 Weed and Feed is a Trimec® brand herbicide containing fertilizer, MCPA, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

EH-1153 Weed and Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential/domestic areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, zoysiagrass, St. Augustinegrass, bahiagrass, and centipedegrass may be treated.

Do not apply this product to bentgrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

WHEN TO USE:

The maximum control of weeds will be obtained from spring or fall applications when the weeds are actively growing.

Dandelion, chickweed, plantain, and henbit are controlled with EH-1153 Weed and Feed. A partial list of susceptible weeds is shown below:

Bedstraw	Dollarweed	Morningglory	Sheep sorrel
Black medic	English daisy	Mustards	Shepherdspurse
Burclover	False dandelion	Peppergrass	Speedwell
Burdock	Filarees	Pigweed	Spurge
Carpetweed	Florida pusley	Plantain (buckhorn)	Spurweed
Chickweed (common, mouseear)	Ground ivy	Poison ivy	Thistle
Chicory	Healall	Poison oak	Wild carrot
Clover (red, white, yellow)	Heartleaf drymary	Puncturevine	Wild garlic
Cocklebur	Henbit	Purslane	Wild geranium
Dandelion	Knotweed	Ragweed	Wild onion
Dock	Lambsquarters	Red sorrel	Yarrow
	Mallow		

APPLICATION INSTRUCTIONS:

1. Spring and fall applications to established turfgrass are advised for the best control. However, EH-1153 Weed and Feed can be applied when broadleaf weeds are present and actively growing. Application of this product to newly seeded areas should be delayed until the third mowing.
2. Mow the turfgrass 2 to 3 days before application.
3. Water the turfgrass thoroughly (1/2 inch) at least 1 to 2 days before application.
4. Apply this product when plant leaves are moist by dew in the early morning or just **after** watering the lawn. Herbicide granules retained in water droplets on the leaves are readily absorbed. So, schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (1/4 inch) before application to provide maximum control.
5. Do not irrigate or water the turfgrass within 48 hours after application.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, and perennial ryegrass: Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass.

For bermudagrass, zoysiagrass, St. Augustinegrass, bahiagrass, and centipedegrass: Apply 3.0 to 3.2 pounds of product per 1,000 square feet of turfgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.2 lb MCPA ae, 0.26 lb MCPP-p ae and 0.10 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [2.40 lb MCPA ae, 0.52 lb MCPP-p ae and 0.20 lb dicamba ae per acre per year].

A uniform broadcast distribution of product on the treated area is essential. Choose a calm day to insure uniform coverage and to avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the specified application rate for any area.

Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product. Refer to the rate table for the appropriate bag sizes and the area to be treated.

SPREADER PATTERNS:

1. Apply two header strips.
2. Operate spreader the longest way of the lawn.
3. For drop spreaders, overlap wheels to insure complete coverage.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

Bag Size, Pounds	Desired Rate Per 1,000 Sq.Ft., Pounds	Treated Area For This Bag Size, Sq. Ft.	Bag Size, Pounds	Desired Rate Per 1,000 Sq.Ft., Pounds	Treated Area For This Bag Size, Sq. Ft.
15	3.0	5,000	35	4.0	8,800
	3.2	4,700			
	4.0	3,800			
16	3.0	5,300	36	4.0	9,000
	3.2	5,000			
	4.0	4,000			
17	3.0	5,700	40	4.0	10,000
	3.2	5,300			
	4.0	4,300			
18	3.0	6,000	45	4.0	11,300
	3.2	5,600			
	4.0	4,500			
20	3.0	6,700	50	4.0	12,500
	3.2	6,300			
	4.0	5,000			
30	4.0	7,500			

Labeling Requirements:

- Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

[LABEL OPTION 2: merged with the text of OPTION 1]

APPLICATION INSTRUCTIONS:

Apply this product when broadleaf weeds are present and actively growing in lawns comprised of the following northern and southern grasses:

- fescues
- Kentucky bluegrass
- perennial ryegrass
- bermudagrass
- centipedegrass
- St. Augustinegrass (except 'Floritam' in Florida)
- bahiagrass
- zoysiagrass

Do not apply this product to bentgrass, carpetgrass, dichondra, 'Floritam' St. Augustinegrass (in Florida), ornamentals, or lawns containing desirable clovers. Application on newly seeded areas should be delayed until the third mowing.

This 16 pound bag of covers 5,000 square feet of lawn when applied at the prescribed rate of 3.2 lb per 1,000 square feet. Measure your lawn. A 20 ft. by 50 ft. area equals 1,000 square feet.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.2 lb MCPA ae, 0.26 lb MCP-p ae and 0.10 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [2.40 lb MCPA ae, 0.52 lb MCP-p ae and 0.20 lb dicamba ae per acre per year].

PREPARATION FOR APPLICATION:

1. Mow lawn 2 to 3 days before application.
2. Water lawn thoroughly with approximately 1/2 inch 1 to 2 days before application to sustain moisture until next watering.
3. Apply this product when plant leaves are moist by dew in the early morning or just after watering the lawn. Herbicide granules retained in water droplets on the leaves are readily absorbed. So, schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (1/4 inch) before application to provide maximum control.

AFTER APPLICATION:

- Do not irrigate or water the turfgrass within 48 hours after application.
- Water lawn thoroughly 1 to 2 days after application to activate fertilizer.

REAPPLICATION:

A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. If needed, the second application of 3.2 lb of product per 1,000 square feet should be used 30 days after the first application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid

- Net Contents: ____ [15 to 50 pounds]
- EPA Est. No. ____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- a) [XXX] pounds covers up to [YYYY] square feet.
- b) Provides excellent response to the fertilizer plus effective broadleaf weed control.
- c) Combines the advantages of fertilization plus a broad range of weed control.
- d) Well maintained lawn benefits our environment.
 - Apply when weeds are young and actively growing.
 - Provides vital nutrients plus time-released nitrogen which encourages grass to fill in bare spots weeds leave behind.
 - Contains sulfur which helps maintain dark green color and encourages plant and root growth.
- e) Grass Cycling and Lawn Benefits: Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients will go back into your lawn.



Professional Lawn Care Association of America

- f) Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:
 - Controls erosion
 - Reduces glare
 - Traps dust, smoke particles and other pollutants
 - Reduces noise
 - Improves soil condition
 - Modifies temperature
 - Controls broadleaf weeds.
- Low phosphate formulation.
- g) Your lawn benefits.
 - Thick, green grass
 - Deep root development
 - Soil vitality
 - Kills dandelions plus (__) other weeds in 3 - 4 weeks.
 - Works on growing weeds.
 - Greens your lawn without excessive growth.
 - Quick green-up in 3 - 5 days after watering, continuing long lasting green for up to 3 months.
 - Thickens your lawn.
 - Promotes lush, even, deep-green turf.
- 4-way greening.
 - White granule - Quick release nitrogen starts greening in 3 - 5 days.
 - Yellow granule - Slow release nitrogen releases slowly over time (OR) Slow Release Nitrogen releases slowly greening your lawn over time.
 - Green granule - Stabilized nitrogen continues greening for up to 3 months, releasing as your lawn needs it. (OR) Stabilized nitrogen. Continuous greening for up to 2 months - available as your lawn needs it.

- Brown granule - Long term slow-released nitrogen with natural nutrient fertilizer components [OR Product Name] natural fertilizer components with slow release nitrogen.

Fertilizer Grade		
NITROGEN 28	PHOSPHATE 1	POTASH 5
Deep Green	Root Growth	Resilience
4 kinds of nitrogen promote deep greening without excessive growth. (OR) 4-way greening without excessive growth.	Low phosphate formulation supplements naturally occurring phosphorus in your soil. (OR) Low phosphate formulation.	Potassium source that strengthens grasses to withstand heat, cold and drought stresses.

- Principles of successful lawn maintenance: (OR) Successful lawn maintenance includes:
 - Spring and fall applications - (Product Name)
 - Good weather - Sun, 60° to 80°F
 - 1" water once a week including rainfall. A typical sprinkler applies 1" moisture in 3 to 4 hours.
 - Proper mowing height - Remove only 1/3 of grass height at each mowing.
 - Optimum thatch level of a lawn is 1/4" to 1/2"
- Annual soil aeration.

The fertilizer grade guaranteed analysis may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

21-3-5 GUARANTEED ANALYSIS	
Total Nitrogen (N)	21%
Available Phosphate (P2O5)	3%
Soluble Potash (K2O)	5%

- Controls [Kills] [desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge] and over [XXX] other [more] broadleaf weeds.
- Helps build a stronger, deeper root system.
- Feeds your lawn with time-released nitrogen for extended greening.
- Encourages early Spring green-up.
- Timed-release fertilizer for professional results
- Use [Apply] when [broadleaf] weeds are actively growing.
- Feeds for up to [time period as designated by subregistrant] [At 72°]
- Quick greening & extended feeding for up to 8 weeks.
- Nitrogen (N) contains timed-release for growing a thick green lawn.
- Phosphorous (P) improves root growth & plant storage capacity.
- Potassium (K) improves plant hardiness & drought resistance.
- For a thick, green lawn.
- Greens fast & lasts.
- Apply in Fall!
- Apply in Fall When Weeds are Growing!
- Kills [Listed] Fall Weeds [As Listed] and Greens the Lawn!
- Kills [Listed] Weeds & Greens the Lawn!
- [Before Photo] and [After Photo] for the Expert Gardener Weed & Feed
- [Before Photo] and [After Photo] for the Expert Gardener Winterizer Weed & Feed
- Kills [XXX][+] [Listed] Weeds [As Listed] While Providing Winter Lawn Protection and Early Spring Green-Up!

- Kills [XXX][+] [Listed] Weeds [As Listed] including Dandelions, Clover and Other Common Lawn Weeds
- Formulated with Scotts® Poly-S® Technology for Longer Lasting Greening and Feeding
- Do not use on Dichondra, Carpetgrass or Floratam, a type of St. Augustinegrass.
- Treats & Feeds [XXXX] sq.ft. [X/X acre = YYYY sq.ft.]
- Expert Gardener® Fertilizer Program (chart)
- For Best Results! Use the Expert Gardener® Fertilizer Program!
- Controls [XXX][+] [Listed] Weeds [As Listed] ! Including:
- Controls [Kills] [Over] [XXX][+] [Listed] [Broadleaf] Weeds [As Listed]
- When to Apply: Apply when broadleaf weeds are actively growing in the early fall.
- When to Apply: Apply when broadleaf weeds are actively growing in spring or early fall
- Spreading Tips: Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. For additional spreader settings, please call 1-800-543-TURF (8873). Calibrate your spreader before beginning your application and adjust settings if necessary.
- What to Expect: Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.
- When you choose Expert Gardener® products, you can be assured of high quality and performance in your lawn and garden. If for any reason you are not completely satisfied with this product, we will refund your money. Please mail your name, address and proof of purchase (dated receipt and package UPC/barcode) to the address below. Also, to ensure quality control, please write us a brief note explaining your experience with our product and reason for seeking a refund.
- Contact Us: Consumer Helpline 1-866-XXX-XXXX
- Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-sc.com>
- Nitrogen (N) For Deep Green Color
- Phosphorus (P) For Strong Roots
- Potassium (K) For Overall Plant Health
- Contains Poly-S® Slow-Release Technology for Longer-Lasting Greening
- Kills and [&] Controls over [XXX][+] Broadleaf Weeds, Including Dandelions, Clovers, Plantains (full list of broadleaf weeds on back panel).
- Optional Warranty Statements:
 - **NOTICE:** Buyer assumes all risks of use, storage or handling of this product [not in accordance with directions]
 - **GUARANTEE:** If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.
- Built-In Spreader.
- Kills [Listed] Weeds
- Builds Strong, Deep Roots
- Water Smart(TM) [graphic of water drop]
- Makes the Most of Every Drop(TM)
- Improves Lawn's Ability to Absorb Water & Nutrients
- Apply to a wet lawn
- Well-fed lawns are more efficient at using available water.
- Feeding with Scotts® [Water Smart(TM)] [products] help(s) grow a thick lawn with more roots to better absorb water and nutrients [than an unfed lawn]. This helps protect your lawn against heat, drought and other stresses.
- Feed regularly for your best lawn.
- [graphic of grass density] One Annual Feeding
- [graphic of grass density] 2-3 Annual Feedings
- [graphic of grass density] 4 Annual Feedings
- Scotts No-Quibble Guarantee® If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

11/11

- Get expert advice for your lawn NOW! Want information on what to use and when? Sign up today to get free expert advice specifically for your lawn
- [computer mouse icon] Visit our website: www.scotts.com
- [phone icon] Consumer Helpline: 1-800-543-TURF (8873)
- Easy To Apply
- Step 1: Flip: Remove seal under cap
- Step 2: Tilt: Tilt bottle as shown
- Step 3: Feed: To prevent over application, tilt bottle as shown when applying the product. Walk at a steady pace and shake bottle side-to-side using your wrist. Apply product uniformly in rows.
- Step 4: Clean-Up: To keep lawn products from reaching storm sewers, sweep or blow any stray particles from hard surfaces back onto the lawn.
- Tip: 1,000 sq.ft. (20x50 ft) is approximately half the size of a tennis court [tennis court image]
- Questions/Comments. [computer mouse icon] www.scotts.com 1-800-543-TURF (8873) [phone icon]
- Kills [XXX][+] [Listed] Weeds [As Listed]
- Promotes thick, green lawns
- Feeds for a thick, green lawn
- Grows strong roots
- Contains time-released nitrogen for prolonged, even feeding

	Early Spring	Mid/Late Spring	Summer	Fall	Grass Type
Northern	Crabgrass Preventer	Weed & Feed	Lawn Food	Lawn Food or Weed & Feed	Bluegrass, Rye, Fescues, Bermuda, Zoysia
Southern	Crabgrass Preventer	Weed & Feed	Lawn Food	Lawn Food or Weed & Feed	Bermuda, Fescues, Bahia
	Nothing until grass starts to green up	St. Augustine Weed & Feed or Lawn Food	Lawn Food	Lawn Food or St. Augustine Weed & Feed	St. Augustine, Centipede, Zoysia

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00798.20110620.amend-proposed-clean.doc
2. **Reason for Issue:** Change signal word, Precautionary Statements, and certain DFU. Ad claims, revised per EPA comments.