



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
2217-807

Date of Issuance:  
SEP 04 1996

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration

Term of Issuance:  
Conditional

Name of Pesticide Product:  
EH1312 Herbicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation  
P.O. Box 14090  
1217 West 12th Street  
Kansas City, MO 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes before you release the product for shipment:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 2217-807".

b. Add an inhalation statement to the Statement of Practical Treatment to read as follows:

"IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention."

c. Add a statement to the front panel to read "For Sale To and Use By Professional Turf or Landscape Personnel Only. Not For Sale To or Use By Homeowners."

Signature of Approving Official:

Date:

SEP 04 1996

page 2  
 EPA Reg. No. 2217-807

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller  
 Product Manager (23)  
 Fungicide-Herbicide Branch  
 Registration Division (7505C)

Enclosure

DK: 305-7546: FHB/PM23		CONCURRENCES						
SYMBOL ▶	7505C							
SURNAME ▶	D. KENNY							
DATE ▶	8/28/96							

3/8

# EH1312 HERBICIDE

## ACTIVE INGREDIENTS:

* Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid.....	18.43%
Diethanolamine Salt of 2,4-dichlorophenoxyacetic acid.....	9.14%
** Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	13.85%
*** Dimethylamine Salt of Dicamba: 3,6-dichloro-o-anisic acid.....	2.75%
INERT INGREDIENTS.....	<u>55.83%</u>
TOTAL 100.00%	

## THIS PRODUCT CONTAINS:

- \*2.03 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 21.54%
  - \*\*1.08 lbs. 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 11.45%
  - \*\*\*0.22 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.29%
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.  
Isomer Specific by AOAC Method 6.DOI-5.

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Hold eyelids open and flush with a steady stream of water for at least 15 minutes. Get medical attention.

**IF SWALLOWED:** Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

**IF ON SKIN:** Wash skin with plenty of soap and water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

See additional Precautionary Statements on the side panel.

NET CONTENTS 1, 2, 5, and 15 U.S. GALLONS

APXXXXXX  
 EPA REG. NO. 2217-XXX  
 EPA EST. NO. 2217-KS-X  
 Mfd. by PBI/GORDON CORPORATION  
 KANSAS CITY, MISSOURI 64101

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated**

SEP 04 1985

**Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended, for the pesticide  
 registered under EPA Reg. No.**

2217-807

EPA REC'D 5/20/96

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS  
AND FOLLOW DIRECTIONS CAREFULLY.

4/8

## PRECAUTIONARY STATEMENTS

### Hazards to Humans & Domestic Animals

**DANGER:** Corrosive. Causes irreversible eye damage. Do not get in eyes, on skin or on clothing. Harmful if swallowed, if absorbed through the skin, or if inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

### - FOR TURF SITES ONLY -

Clothing Requirement Statements: When mixing, loading, or applying this product, wear long-sleeved shirt, long pants, socks, shoes, chemical-resistant gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection.

Personal Hygiene Statements: After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

### - FOR INDUSTRIAL SITES ONLY -

Do not reenter or permit workers to reenter treated areas until sprays have dried or dusts have settled.

Clothing Requirement Statement: When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow and temple protection. For applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.

Personal Hygiene Statements: Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more: Do not open-pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

**ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area. Keep from freezing.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by other procedures approved by state and local authorities. For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

### OTHER USE PRECAUTIONS

#### OBSERVE CAREFULLY.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not use on St. Augustinegrass or dichondra nor on lawns or turf where desirable clovers are present. Note: Coarse sprays are less likely to wind-drift. Do not spray exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not spray while grass is emerging from dormancy. Do not apply to newly seeded lawns until after the second or third mowing. Do not broadcast apply when air temperature exceeds 85°F. Some injury may be expected with spot treatments when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. Be careful not to overdose centipedegrass. This species is sensitive to 2,4-D, a component of this product. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury. Do not apply this product through any type of irrigation system.

### HOW MUCH TO USE AND WHERE:

This product is intended for use on lawns, ornamental and residential turf, golf courses, parks, cemeteries and similar non-crop areas.

The maximum application rate to turf is 1.02 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

**Bahiagrass, Common Bermudagrass, Bluegrass, Fescue, Ryegrass, and Zoysiagrass:**

**Broadcast Treatments** - Use 3¼ to 4.0 pints of product per acre in 20 to 175 gallons of water. Or, apply 1.2 to 1.5 fluid ounces of product per 1,000 square feet with spray volumes of 0.5 to 4.0 gallons per 1,000 square feet. Do not use more than the recommended rate. Bermudagrass and zoysiagrass are moderately sensitive to 2,4-D.

**Spot Treatment** - Mix 1.2 to 1.5 fluid ounces of product in 1 gallon of water per 1,000 square feet. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stage.

**Controlled Droplet Applicator** - Add 2½ pints of product in 1 to 4 gallons of water per 33,000 square feet.

**Centipedegrass, Creeping Bentgrass and Hybrid Bermudagrass:**

**Broadcast Treatments** - Use 1.8 pints of product per acre in 40 to 175 gallons of water. Do not use more than the recommended rate. Bentgrass and hybrid bermudagrass are moderately sensitive to 2,4-D.

**Spot Treatment** - Mix 0.66 (2/3) fluid ounces of product in 1 gallon of water per 1,000 square feet.

**Controlled Droplet Applicator** - Add 1.4 pints of product in 1 to 4 gallons of water per 33,000 square feet.

**General** - Mix in 20 to 175 gallons of water per acre. Use higher application rates when weeds are dense or in flower. If hardy weeds persist, make a second application after three weeks. Apply when weeds are young and actively growing. Avoid applications during long excessively dry or hot periods. When grasses are under stress use reduced rates to reduce potential injury. Do not apply to newly seeded turf until after the second or third mowing. Seed can be sown 3 to 4 weeks after application at recommended rates.

- WEEDS CONTROLLED -		
Bedstraw	Innocence (Blue-eyed Mary)	Purple Cudweed
Black Medic	Knotweed	Purslane
Buckhorn	Lambquarters	Ragweed
Burdock	Lawn Burweed	Redweed
Chicory	Lespedeza	Sheep Sorrel
Chickweed	Mallow	Shepherdspurse
Clover	Morningglory	Speedwell
Corn Speedwell	Old World Diamond Flower	Spurge
Dandelion	Peppergrass	Wild Carrot
Dock	Pigweed	Wild Garlic
Ground Ivy	Plantains	Wild Lettuce
Healall	Poison Ivy	Wild Onion
Henbit	Poison Oak	Yarrow

- INDUSTRIAL SITES -

EH1312 Herbicide can be incorporated into industrial vegetation management programs as broadcast or spot treatments. EH1312 Herbicide may be useful in the following noncropland areas:

- Highway rights-of-way (principal, interstate, state, and country highways), interchange ramps, waysides, service areas, and unpaved roads.
- Municipal, state, and federal lands such as airports, military installations, schools/universities, libraries, and hospitals.
- Commercial/industrial areas including industrial parks, tank farms, plant sites, storage areas, fencerows, and utility rights-of-way.

EH1312 Herbicide may be applied to established cool season and warm season grasses established in noncropland sites. Cool season grasses include tall fescue, perennial ryegrass, smooth bromegrass, orchardgrass, and reed canarygrass. Warm season grasses include bermudagrass, zoysiagrass, and bahiagrass.

**USE PRECAUTIONS FOR INDUSTRIAL SITES:**

- Do not apply this product through any type of irrigation system.
- Avoid spray drift to cotton, soybeans, tomatoes, tobacco, grapes, fruit trees, flowers or garden crops, and all other hormone herbicide-sensitive desirable plants.
- Do not spray roots of desirable trees and shrubs.
- Do not apply herbicide when wind speed is sufficient to cause drift. Do not apply herbicides when a temperature air inversion exists. An air inversion may be detected by creating a smoke column and observing a layering effect.
- Do not apply when temperature exceeds 85°F and humidity is high.
- To reduce the potential of spray drift, use large spray droplets and low pressures. Do not use nozzles which produce fine spray droplets under high pressure. The use of thickening agents or anti-drift additives and drift-reducing equipment may be of value in preventing spray drift.
- Care should be taken not to make applications where runoff could contaminate bodies of water or spray drift could carry the chemical to food crops or grazing lands where cattle, sheep, goats, swine or poultry would be exposed.

- APPLICATION RATES AND SPRAY VOLUMES FOR INDUSTRIAL SITES -

**Broadcast Treatment** - Apply 3¼ to 4.0 pints of product per acre as a broadcast treatment to annual, biennial, and perennial broadleaf weeds. Biennial and perennial weeds may require a follow-up or sequential treatment. Spray volumes of 15 - 150 gallons per acre are recommended.

Generally, the lower application rates within the specified range will provide satisfactory control. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

**Spot Treatments for Mixed Brush** - Spraying can be effective throughout the growing season from full leaf to leaf drop for mixed brush. Full cover sprays should be applied during warm weather when brush and broadleaf weeds are young and actively growing. All leaves, stems, and shoots should be thoroughly wetted to the ground.

**For High Volume Foliar Applications** - Use 6.0 to 8.0 pints of product in 30 - 100 gallons of water (1.0 - 3.0% vol./vol) and apply as a full cover spray wetting all leaves, stems, and root collars of the woody plants. This spray concentration requires a spray volume of 100 - 300 gallons per acre depending on the height and density of the mixed brush.

BRUSH:		
Ash	Elms	Oak
Aspen	Gooseberry	Shortleaf Pine
Birches	Honey Locust	Thistle
Black Cherry	Kudzu	Willows
Brambles	Multiflora Rose	
WEEDS:		
Bedstraw	Innocence (Blue-eyed Mary)	Purple Cudweed
Black Medic	Knotweed	Purslane
Buckhorn	Lambsquarters	Ragweed
Burdock	Lawn Burweed	Redweed
Chicory	Lespedeza	Sheep Sorrel
Chickweed	Mallow	Shepherdspurse
Clover	Morningglory	Speedwell
Corn Speedwell	Old World Diamond Flower	Spurge
Dandelion	Peppergrass	Wild Carrot
Dock	Pigweed	Wild Garlic
Ground Ivy	Plantains	Wild Lettuce
Healall	Poison Ivy	Wild Onion
Henbit	Poison Oak	Yarrow

**LIMITED WARRANTY AND DISCLAIMER.**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.