

2217-808

NOTIFICATION 10/31/2002

1/7

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 02-28-95



United States Environmental Protection Agency Washington, DC 20460

- Registration
Amendment
Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number: 2217-808
2. EPA Product Manager: James A. Tompkins
3. Proposed Classification: [X] None [] Restricted
4. Company/Product (Name): EH795 Residential Herbicide
5. Name and Address of Applicant: PBI/Gordon Corporation, Acme Division, PO Box 014090, Kansas City, Missouri 64101
6. Expedited Review: In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name

Section - II

- Amendment - Explain below.
Resubmission in response to Agency letter dated
Notification - Explain below.
Final printed labels in response to Agency letter dated
'Me Too' Application.
Other - Explain below.

NOTIFICATION
OCT 31 2002

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION per PR Notice 98-10: We ask to present the following product features on the container labeling: 1) 'Controls crabgrass, yellow nutsedge & broadleaf weeds' 2) 'For selective control of grass and broadleaf weeds in bermudagrass, bluegrass, tall fescue, turf-type tall fescues, and zoysiagrass turf. 3) graphic display of water drop surrounded by phrase 'Concentrate, Mixes with water' One copy of the proposed labeling is enclosed for your review. This notification is consistent with the provisions of PRN 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PRN 98-10 and 40CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:
Child-Resistant Packaging: [] Yes* [] No
Unit Packaging: [] Yes [] No
Water Soluble Packaging: [] Yes [] No
2. Type of Container: [] Metal [] Plastic [] Glass [] Paper [] Other (Specify)
3. Location of Net Contents Information: [] Label [] Container
4. Size(s) Retail Container
5. Location of Label Directions: [] On Label [] On Labeling accompanying product
6. Manner in Which Label is Affixed to Product: [] Lithograph [] Paper glued [] Stenciled [] Other

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)
Name: Craig Martens Title: Manager of Regulatory Services Telephone No. (Include Area Code): 816/468-6287
Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.
2. Signature: Craig Martens
3. Title: Manager of Regulatory Services
4. Typed Name: Craig Martens
5. Date: October 24, 2002
6. Date Application Received (Stamped)

2/7

Official Name:

EH795 RESIDENTIAL HERBICIDE

Brand Name: **TRIMEC® 699 BROADLEAF PLUS CRABGRASS HERBICIDE**

Not for use on turf being grown for sale or other commercial use as sod,
or for commercial seed production, or for research purposes.

SHAKE WELL BEFORE USING

ACTIVE INGREDIENTS:

Monosodium acid methanearsonate	18.00%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	5.83%
Dimethylamine salt of (+)-(R)-2-(4-chloro-2-methylphenoxy) propanoic acid	2.93%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	1.46%
INERT INGREDIENTS:	<u>71.78%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

1.80 lbs. monosodium methanearsonate per gallon or 18.00%. Total Arsenic (as elemental) all in water soluble form 8.33%
 0.48 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.84%
 0.24 lb. (+)-(R)-2-(4-chloro-2-methylphenoxy) propanoic acid equivalent per gallon or 2.42%
 0.12 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.21%
 Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN

WARNING

NOTIFICATION
OCT 31 2002

See side panels for additional Precautionary Statements
and Statement of Practical Treatment.

NET CONTENTS: ONE GALLON

AP102402
 EPA REG NO 2217-808
 EPA EST. NO. 2217-KS-1
 MANUFACTURED BY:





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if absorbed through skin. Avoid breathing spray mist. Avoid contact with skin. When using this product wear long-sleeved shirt, long pants, socks, shoes, rubber gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more: Do not open-pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Statement of Practical Treatment

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from treated area.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

4/7

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

Storage & Disposal For Metal Drums:

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

WHERE TO USE:

EH795 Residential Herbicide is intended for residential and ornamental lawns with applications by homeowners only. This product contains four proven herbicides that are effective on hard-to-kill grass and broadleaf weeds. Spot treatments of this product are applied after the weeds have emerged with hand-operated sprayers (plunger or pump-up sprayers).

EH795 Residential Herbicide controls a wide range of broadleaf weeds including dandelion, spurge, oxalis, crabgrass, dallisgrass, and yellow nutsedge in bermudagrass, Kentucky bluegrass, perennial ryegrass, tall fescue, and zoysiagrass lawns. A partial list of weeds controlled is shown below.

PARTIAL LIST OF WEEDS CONTROLLED

Aster	Healall	Purslane
Bedstraw	Henbit	Ragweed
Black medic	Knotweed	Sandbur
Buckhorn	Lambsquarters	Sheep sorrel
Burdock	Lespedeza	Shepherdspurge
Chicory	Mallows	Speedwell
Chickweed	Morningglory	Spurge
Clovers	Nutsedge	Wild carrot
Crabgrass	Oxalis	Wild garlic
Dallisgrass	Peppergrass	Wild lettuce
Dandelion	Pigweeds	Wild onion
Docks	Plantain	Yarrow
Goosegrass	Poison ivy	Yellow wood sorrel
Ground ivy	Poison oak	

USE PRECAUTIONS:

- ◆ Do not use a hose end sprayer.
- ◆ Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
- ◆ Do not exceed the specified dosage for any area. Be particularly careful under the dripline of trees and ornamentals.
- ◆ Do not use this product on St. Augustinegrass, centipedegrass, carpetgrass, creeping fescue, dichondra, nor on lawns with desirable clovers.
- ◆ Keep children and domestic animals off treated area until this material has been watered into the soil or the grass has dried.
- ◆ Do not apply while grass is emerging from dormancy.

WHEN TO APPLY:

Apply when weeds are young and actively growing and the lawn is healthy, moist and not under stress from drought conditions. For bluegrass, perennial ryegrass, and tall fescue, apply this product when air temperatures are between 70°F and 80°F. For bermudagrass and zoysiagrass, apply this product when air temperatures are between 70°F and 90°F. Applications made when air temperatures exceed these conditions may result in injury to the lawn.

Do not apply this product to newly seeded lawns until after the fourth mowing. Seed can be sown 3 to 4 weeks after application.

HOW TO APPLY:

If the soil is dry, then water thoroughly 24 hours before application. Mow lawn to the proper height 1 - 3 days before application. Using a hand-operated sprayer (plunger or pump-up sprayer) spray weeds evenly until lightly covered. Over-spray can injure the turfgrass. Do not water lawns for 24 hours after application. Temporary discoloration may occur with the use of this product. After 24 hours, a thorough watering is recommended.

For Bermudagrass and Zoysiagrass: Mix 4 fluid ounces (8 tablespoons) of product with 2.0 gallons of water for treatment of 1,000 square feet of turfgrass. Do not apply when the air temperatures are above 90°F. A second application may be made in 14 days if necessary.

Table 1. Mixing instructions for bermudagrass and zoysiagrass with hand operated sprayers.

Area to Spray, Square Feet	Amount of Water, Gallons	Amount of Product,	
		Tablespoons	Fluid Ounces
500	1	4	2
1,000	2	8	4
1,500	3	12	6

For Kentucky Bluegrass, Perennial Ryegrass, and Tall Fescue: Mix 3 fluid ounces (6 tablespoons) of product with 2.0 gallons of water for treating 1,000 square feet of turfgrass. Do not apply when the air temperatures are above 80°F. A second application may be made in 14 days if necessary.

Table 2. Mixing instructions for Kentucky bluegrass, perennial ryegrass, and tall fescue with hand operated sprayers.

Area to Spray, Square Feet	Amount of Water, Gallons	Amount of Product,	
		Tablespoons	Fluid Ounces
500	1	3	1½
1,000	2	6	3
1,500	3	9	4½

67

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

7/7

APPENDIX

I. Advertising claims that may be presented on container labeling:

- ◆ Covers up to 5,000 square feet.
- ◆ Controls crabgrass, yellow nutsedge & broadleaf weeds
- ◆ For selective control of grass and broadleaf weeds in bermudagrass, bluegrass, tall fescue, turf-type tall fescues, and zoysiagrass turf.
- ◆ Controls crabgrass, yellow nutsedge & broadleaf weeds.
- ◆ For selective control of grass and broadleaf weeds in Bermudagrass, Bluegrass, Tall Fescue, Turf-Type Tall Fescues, and Zoysiagrass Turf.
- ◆

