



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 22, 2024

### **SENT BY EMAIL**

Eric Smith  
esmith@pbigordon.com  
PBI/GORDON CORPORATION

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Adding optional marketing statements and claims.  
Product Name: EH1328 WEED AND FEED  
Admin Number: 2217-811  
EPA Receipt Date: 08/23/2024  
Action Case Number: 00626653

Dear Eric Smith:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Elizabeth Danon via email at [danon.elizabeth@epa.gov](mailto:danon.elizabeth@epa.gov).

Sincerely,

*Kable Bo Davis*

Kable Bo Davis, Senior Advisor  
HB, RD  
Office of Pesticide Programs

# EH1328 WEED AND FEED

## EPA Reg. No. 2217-811

**ACCEPTED**

10/22/2024

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 2217-811

### ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt.....	0.78%
Mecoprop-p, dimethylamine salt .....	0.15%
Dicamba, dimethylamine salt.....	0.04%

OTHER INGREDIENTS: .....	99.03%
--------------------------	--------

TOTAL .....	100.00%
-------------	---------

### THIS PRODUCT CONTAINS:

13.00 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.65%  
2.60 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.13%  
0.60 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.03%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

**Net Contents:** \_\_\_\_\_

**EPA Est. No.** \_\_\_\_\_

### Company Name and Address:

PBI/Gordon Corporation  
P.O. Box 860350  
Shawnee, KS 66286



## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

<b>First Aid</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not give anything by mouth to an unconscious person. Do not induce vomiting unless told to by a poison control center or doctor.</li></ul>
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### **[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns until after the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

#### **WHERE TO USE:**

EH1328 Weed and Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others as listed, which are commonly found in home lawns. In addition, a [X-X-X] fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

EH1328 Weed and Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

#### **APPLICATION INSTRUCTIONS:**

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

<b>APPLICATION INSTRUCTIONS: (Optional Tabular Format)</b>	
<b>Application Schedule</b>	<b>Cultural Practices</b>
1 to 2 days before application	Mow lawn to normal height 1 to 2 days before application.
1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.
Day of application	Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material.
1 to 2 days after application	Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Do not apply to newly seeded lawns until after the third mowing. Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

## HOW MUCH TO APPLY:

**For Kentucky bluegrass, fescues, and perennial ryegrass, bermudagrass, zoysiagrass, St.**

**Augustinegrass and centipedegrass:** Apply 3.0 to 3.6 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.0 to 3.6 pounds of product per 1,000 square feet must be made at least 30 days after the first application. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application [(1.02 lb 2,4-D ae, 0.2 lb MCPP-p ae and 0.05 lb dicamba ae per acre per application)]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. [(2.04 lb 2,4-D ae, 0.40 lb MCPP-p ae and 0.10 lb dicamba ae per acre)].

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and to avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area.

## [Begin Optional Language]

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product.

## CALIBRATION INSTRUCTIONS:

**[Note to Supplemental Registrants:** Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.

[Coverage: To determine the square footage of the application area, multiply the length of the lawn by its width. This equals your total square footage. Deduct any non-lawn areas (driveway, sidewalk, house, etc.) One bag covers XXXX sq.ft. or XX pounds per 1,000 sq. ft.]

[sq.ft. calculation image Length x Width = Total Sq.Ft. -- Non-Lawn Area = Total Sq. Ft. Coverage you need]

[Spreader Settings [spreader images]

Calibrating spreader: Adjust spreader to apply X pound of product per XXX sq.ft.

Spreading patterns: First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to ensure even coverage [spreading pattern image(s)]

Bag Size, Pounds	Application Rate, Pounds/1,000 Sq.Ft.	Treated Area for This Bag Size, Square Feet	Bag Size, Pounds	Application Rate, Pounds/1,000 Sq.Ft.	Treated Area for This Bag Size, Square Feet
15	3.0	5,000	18	3.0	6,000
	3.2	4,700		3.2	5,600
	3.4	4,400		3.4	5,300
	3.6	4,200		3.6	5,000
16	3.0	5,300	20	3.0	6,600
	3.2	5,000		3.2	6,300
	3.4	4,700		3.4	5,900
	3.6	4,400		3.6	5,600
17	3.0	5,600	36	3.0	12,000
	3.2	5,300		3.2	11,200
	3.4	5,000		3.4	10,600
	3.6	4,700		3.6	10,000

#### SPREADER PATTERNS AND PRODUCT DISTRIBUTION:

1. Apply two header strips.
2. Operate spreader the longest way of the lawn.
3. For drop spreaders, overlap wheels to ensure complete coverage.
4. Close spreader openings when turning.

#### SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to ensure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

**Coverage** - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

[End optional language]

BROADLEAF WEEDS [207]			
Annual yellow sweetclover	Cockle	Musk thistle	Spiny amaranth
Aster	Cocklebur	Mustard	Spiny cocklebur
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny sowthistle
Bedstraw	Creeping jenny	Narrowleaf vetch	Spotted catsear
Beggarticks	Cudweed	Nettle	Spotted spurge
Betony, Florida	Curly dock	Orange hawkweed	Spurweed
Bindweed, field	Daisy, English	Oriental cocklebur	Stinging nettle
Bird vetch	Daisy fleabane	Oxalis	Strawberry, India mock
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall nettle
Bittercress, hairy	Dandelion	Parsnip	Tall vervain
Bitterweed	Dichondra	Pearlwort	Tansy ragwort
Black-eyed Susan	Dogbane	Pennycress	Tansy mustard
Black medic	Dogfennel	Pennywort	Tanweed
Black mustard	Dollarweed	Peppergrass	Thistle
Blackseed plantain	Elderberry	Pepperweed	Trailing crownvetch
Blessed thistle	False dandelion	Pigweed	Tumble mustard
Bloodflower milkweed	Falseflax	Pineywoods bedstraw	Tumble pigweed
Blue lettuce	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue vervain	Fiddleneck	Plantain	Venice mallow
Bracted plantain	Florida pusley	Poison ivy	Virginia buttonweed
Brassbuttons	Frenchweed	Poison oak	Virginia creeper
Bristly oxtongue	Galinsoga	Pokeweed	Virginia pepperweed

BROADLEAF WEEDS [207]			
Broadleaf dock	Goathead	Poorjoe	Water pennywort
Broadleaf plantain	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broomweed	Ground ivy	Prickly lettuce	Western clematis
Buckhorn	Gumweed	Prickly sida	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild aster
Bull thistle	Healall	Prostrate spurge	Wild buckwheat
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild carrot
Burclover	Heathaster	Puncturevine	Wild four-o'clock
Burdock	Hedge bindweed	Purslane, common	Wild garlic
Burning nettle	Hedge mustard	Ragweed	Wild geranium
Burweed	Hemp	Red sorrel	Wild lettuce
Buttercup	Henbit	Redroot pigweed	Wild marigold
Buttonweed	Hoary cress	Redstem filaree	Wild mustard
Canada thistle	Hoary plantain	Rough cinquefoil	Wild onion
Carolina geranium	Hoary vervain	Rough fleabane	Wild parsnip
Carpenterweed	Horsenettle	Russian pigweed	Wild radish
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild rape
Catnip	Knawel	Scarlet pimpernel	Wild strawberry
Catsear	Knotweed	Scotch thistle	Wild sweet potato
Chickweed, common	Kochia	Sheep sorrel	Wild vetch
Chickweed, mouseear	Lambsquarters	Shepherdspurse	Woodsorrel
Chicory	Lespedeza	Slender plantain	Woolly croton
Cinquefoil	Mallow	Smallflower galinsoga	Woolly morningglory
Clover, crimson	Matchweed	Smooth dock	Woolly plantain
Clover, hop	Mexicanweed	Smooth pigweed	Wormseed
Clover, red	Milk vetch	Sorrel	Yarrow
Clover, strawberry	Morningglory	Sowthistle	Yellow rocket
Clover, sweet	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, white	Mugwort	Speedwell	

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.



# APPENDIX

## 1. Advertising claims / label statements that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- 3-way active ingredient for optimum weed control
- [Apply] [Use] when weeds are [actively] growing
- Weed control works through foliage contact
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and other broadleaf weeds.
- Controls dandelions, thistle, clover, and over [XXX] other listed weeds (Note to Reviewer: claim not to exceed 207 total listed weeds)
- Controls dandelion and over [XXX] broadleaf weeds (Note to Reviewer: claim not to exceed 207 total listed weeds)
- Controls [over] [XXX] [broadleaf] weeds (Note to Reviewer: claim not to exceed 207 total listed weeds)
- 2 in 1 Kills Weeds\* Greens Your Lawn \*[as] listed
- Kills listed weeds to the root
- Contains three (3) proven weed killers
- [Kills][Controls] most common weeds\* \*[as] listed
- Contains slow release nitrogen for long lasting feeding without burning, when applied as directed.
- [Contains] Slow release nitrogen for a longer lasting green lawn
- [Controlled] [Slow] release nitrogen for extended feeding
- Fortified with Milorganite® [for use only on formulations containing milorganite]
- Nitrogen [XX] Promotes thick, green turf
- Phosphorus free [formula] [for use on formulation containing zero phosphate]
- Phosphorus 0 No phosphate runoff into lakes and streams [for use on formulations containing zero phosphate]
- Potassium [XX] Strengthens grass against disease
- Feeds and thickens your lawn
- Fast [quick] greening [formula]
- Long-lasting feeding [greening]
- Contains [X%] Iron for deep greening
- Contains Iron for deep greening
- Deep greening formula
- Prepares your lawn for winter
- Formulated to promote root building and winter hardiness
- Nourishes all winter for a quick spring green-up
- Better green-up next spring
- Continual root development for a thick, lush lawn
- Promotes a thick, lush lawn
- Builds strong roots
- Stimulates root growth
- Apply when lawn is wet/moist
- Labeled for use on Bermudagrass, Zoysiagrass, St. Augustinegrass, [and] Centipedegrass
- For use on established cool season or warm season turfgrasses
- Feeds northern or southern grasses
- Apply late spring/early summer [when weeds are actively growing]
- Consistent blend/consistent feeding
- Continues to feed for [XX] weeks [months]
- Healthy lawns require less water
- A granular fertilizer containing Trimec® herbicide which controls: dandelion, chickweed, knotweed, plantains, henbit, spurge and many other broadleaf weeds [some of which are listed on the back panel]
- A Granular Fertilizer Containing Trimec® Herbicide Which Controls: [weeds]



- (Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Answers to FAQ's on the internet at [www.pbigordon.com](http://www.pbigordon.com)
- Lilly Miller Ultragreen [product] feeds and protects all year long!
- Broad spectrum weed control & fertilizes in one application
- formulated to control weeds and prepare your lawn for Winter dormancy
- Broadleaf weed control plus provides a full feeding of fertilizer for a lush, healthier lawn
- Greens lawns and develops deep roots while controlling weeds
- Improves color, density, and uniformity to turf
- (Product name) contains a balanced herbicide blend that controls a wide variety of existing broadleaf weeds plus provides a full feeding of fertilizer. This product will continue to feed your lawn for up to 6 weeks thereby improving the color, density and overall uniformity of the turf grass.
- Answers to FAQ's on the internet at [www.ecgrowproducts.com](http://www.ecgrowproducts.com) [website and/or computer image]
- Control listed weeds and fertilize in one application
- **Maximize your lawn**
- Covers 10,000 Sq. Ft. [Covers XXX Sq. Ft.]
- Calculating total lawn area: [graphic] length x width = total sq. ft. then subtract non-lawn areas (house, garden, driveway)
- 100% Satisfaction Guarantee Or Your Money Back
- Satisfaction guaranteed
- Stein's Policy, 100% Satisfaction Guaranteed, If for any reason you are not satisfied with the results after using this premium product, you are entitled to replacement bags of fertilizer or your money back. Simply bring proof of purchase to any Stein Garden Center.
- [Brand Name] 100% Satisfaction Guarantee
- Manufactured exclusively for [company name]
- [www.steingg.com](http://www.steingg.com)
- FAQ's – Product Info – Tips [www.ecgrowproducts.com](http://www.ecgrowproducts.com)
- If after using this product, you are dissatisfied with the result, simply mail the sales receipt, a brief description of the problem and your contact information to the address listed below for a refund of the purchase price of the product.
- For more information on this and other products, visit [website address].

Referrals to other products used sequentially:

1<sup>st</sup> Application

2<sup>nd</sup> Application

3<sup>rd</sup> Application

4<sup>th</sup> Application


- **Stein Lawn Care Program**  
[If you have weeds, use Stein Lawn Care Program]  
Application 1  
Crabgrass Preventer plus Lawn Food  
Application 2  
Weed Control Plus Lawn Food  
Application 3  
Lawn Food  
Application 4  
Winterizer
- **Forevergreen [Lawn Builder] [4-Step] [Lawn Care] Program**  
[If you have weeds, use Forevergreen [Lawn Builder] [4-Step] [Lawn Care] Program]  
[Allow six weeks between applications]  
Forevergreen Crabgrass Preventer
  - First application/ Step 1

- Mar to early May [Early Spring]
  - Season-long crabgrass control
  - Greens your lawn with slow release nitrogen
  - Slow release nitrogen for extended feeding
- Forevergreen Weed & Feed
- Second application
  - [Apply] Mid-April to July
  - Apply Late Spring/Early Summer
  - Kills dandelions, clover, plantains, and over 150 other broadleaf weeds
  - Thickens and greens your lawn with slow release nitrogen
  - Controls dandelion, clover and over 150 other listed weeds
  - Greens lawns and develops roots while controlling weeds
- Forevergreen Turf Food
- Third application
  - June to September
  - Thickens and promotes a lush green lawn
  - Greens with Iron and Sulfur
  - Slow release nitrogen for long-lasting feeding
  - Contains iron for deep greening
- Forevergreen Fall Turf Food
- Fourth application
  - Late Aug to Nov
  - Slow release nitrogen for long-lasting feeding
  - Enhances root building and prepares your lawn for winter dormancy

#### Optional/Example Graphics

The following are representative examples of graphics that could be used in the marketed label. If the graphic is used, minor changes to the exact color, shapes or orientation of items depicted, or the amount or size of graphic may occur based on marketing decisions and the available space on the label.





## 2. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “3.1 State Restrictions”]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N) .....	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	X%
Soluble Potash (K <sub>2</sub> O) .....	X%
Sulfur (S) .....	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Idaho: “Apply only as directed”
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the

associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the "Soil Erosion and Sediment Control Act," P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots."

- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: "In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency."
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program."

## **DOCUMENT CONTROL INFORMATION**

**1. Unique Label Identifier:** 002217-00811.20240822.notif-proposed-highlighted

**2. Reason for Issue:** Ad Claims

<b>Internal Usage Notes</b>
Associated Container Label: Supplemental Distributors