

# EH1335 HERBICIDE

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.


SHAKE WELL BEFORE USING

ACTIVE INGREDIENTS:

*Monosodium acid methanearsonate .....	0.36%
**Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid.....	0.12%
***Dimethylamine Salt of (+)-(R)-2-(4-chloro-2-methylphenoxy)propanoic acid ....	0.06%
****Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid).....	0.02%
INERT INGREDIENTS: .....	99.44%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- \*0.030 lbs. monosodium methanearsonate per gallon or 0.36%.
  - Total Arsenic (as elemental) all in water soluble form 0.17%
  - \*\*0.008 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 0.10%
  - \*\*\*0.004 lbs. (+)-(R)-2-(4-chloro-2-methylphenoxy) propanoic acid equivalent per gallon or 0.05%
  - \*\*\*\*0.002 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.02%
- Isomer Specific by AOAC Method.



KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional Precautionary Statements and First Aid.

NET CONTENTS 24 and 32 FLUID OUNCES

AP092899  
 EPA Reg. No. 2217-815  
 EPA Est. No. 2217-KS-X  
 MANUFACTURED BY:





**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards To Humans & Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. When using this product wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

*Containers over 1 gallon and less than 5 gallons:* Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

**First Aid**

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**ENVIRONMENTAL HAZARDS:**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

**STORAGE & DISPOSAL**

**STORAGE:** Store in original container in a locked storage area. Keep from freezing.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse container. Rinse thoroughly before discarding in trash.

**WHERE TO USE:**

EH1335 Herbicide is intended for residential and ornamental lawns with applications by homeowners only. This product contains four proven herbicides that are effective on hard-to-kill grass and broadleaf weeds. EH1335 Herbicide is recommended for sparse weed infestations and for follow-up treatments of broadcast herbicide applications. This ready-to-use herbicide is intended for spot treatments only. Weeds that have not emerged at the time of application will not be controlled.

EH1335 Herbicide controls a wide range of broadleaf weeds including dandelion, spurge, oxalis, crabgrass, dallisgrass, and yellow nutsedge in bermudagrass, Kentucky bluegrass, tall fescue, and zoysiagrass lawns. A partial list of weeds controlled is shown below.

**- PARTIAL LIST OF WEEDS CONTROLLED -**

Aster	Healall	Purslane
Bedstraw	Henbit	Ragweed
Black medic	Knotweed	Sandbur
Buckhorn	Lambsquarters	Sheep sorrel
Burdock	Lespedeza	Shepherdspurse
Chicory	Mallows	Speedwell
Chickweed	Morningglory	Spurge
Clovers	Nutsedge, yellow	Wild carrot
Crabgrass (smooth, large)	Oxalis	Wild garlic
Dallisgrass	Peppergrass	Wild lettuce
Dandelion	Pigweed	Wild onion
Docks	Plantain	Yarrow
Goosegrass	Poison ivy	Yellow woodsorrel
Ground ivy	Poison oak	

**HOW TO USE:**

**For Containers with Triggers Sprayers:** EH1335 Herbicide is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Usually, two (2) pumps of this trigger sprayer will provide adequate coverage of individual weeds. Excessive spraying can injure some lawns.

**For Refill Containers:** EH1335 Herbicide is ready-to-use and is applied directly from the container. Remove the cap from the container and attach the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Excessive spraying can injure some lawns.

**USE PRECAUTIONS:**

- ◆ Do not use a hose end sprayer.
- ◆ Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
- ◆ Do not use this product on St. Augustinegrass, centipedegrass, carpetgrass, creeping fescue, dichondra, nor on lawns with desirable clovers.
- ◆ Keep children and domestic animals off treated area until this material has been watered into the soil or the grass has dried.
- ◆ Do not apply while grass is emerging from dormancy.

**WHEN TO APPLY:**

Apply when weeds are young and actively growing for the best results. Schedule the application to match the conditions of adequate soil moisture and these air temperatures. For Kentucky bluegrass and tall fescue, apply this product when air temperatures are between 70°F and 80°F. For bermudagrass and zoysiagrass, apply this product when air temperatures are between 70°F and 90°F.

Applications made when air temperatures exceed these conditions may result in injury to the lawn. Temporary discoloration may occur with the use of this product.

Weed regrowth may occur because of the growth from stems and rhizomes, environmental conditions, or misapplication. Also, weeds can germinate from seed and emerge throughout the season. Two or more applications at 10 - 14 day intervals may be required.

Weed control may be reduced when applications are made during periods of inadequate soil moisture. If the soil is dry, then water thoroughly 24 hours before application. Do not water lawns for 24 hours after application.

Newly seeded lawns may be treated after the fourth mowing. Seed can be sown 3 to 4 weeks after application.

**LIMITED WARRANTY AND DISCLAIMER.**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Gordon's® and Trimec® are registered trademarks of PBI/Gordon Corporation.  
Day-Glo® is a registered trademark of Day-Glo Color Corporation.