

2217-815

10-3-2002

1/9

ACCEPTED

OCT 3 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2217-815

AMAZING!
Kills crabgrass, nutsedge, dandelion
and other lawn weeds too...
without injury to lawn grasses!

GORDON'S TRIMEC®



READY TO USE
RTU

Contains
**COLOR
MARKER**

**JUST
2 SQUIRTS
DOES
THE JOB!**

*Shake Well
Before Using*

**KEEP OUT OF
REACH OF
CHILDREN**

CAUTION

See attached booklet on
back of container for
Precautionary Statements
and First Aid.

2,4-dichlorophenoxyacetic acid	0.17%
Dimethylamine salt of (+)-(R)-2-(4-chloro-2-methylphenoxy)propanoic acid	0.05%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.02%
INERT INGREDIENTS:	99.76%

THIS PRODUCT CONTAINS: TOTAL 100.00%

*0.030 lbs. monosodium methanearsonate per gallon or 0.36%. Total Arsenic (as elemental) all in water soluble form 0.17%. **0.008 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 0.10%. ***0.004 lbs. (+)-(R)-2-(4-chloro-2-methylphenoxy)propanoic acid equivalent per gallon or 0.05%. ****0.002 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.02%. Isomer Specific by AOAC Method.

738/3-2003

NET CONTENTS: 32 FLUID OUNCES

2/9




GORDON'S

TRIMEZ PLUS



READY TO USE
RTU




AMAZING!

*Kills crabgrass, nutsedge,
dandelion and other lawn
weeds too . . . without injury to lawn grasses!*

**SHAKE WELL
BEFORE USING!**
*Peel open for Precautionary
Statements, First Aid and
Full Directions for Use.*

738/3-2002 AP031202
EPA REG. NO. 2217-815
EPA EST. NO. 2217-KS-1

Manufactured By
 **pbi / GORDON**
CORPORATION
An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
www.pbigordon.com



3/9

GORDON'S®
TRIMEC®
PLUS



READY TO USE!
RTU

ACTIVE INGREDIENTS:

* Monosodium acid methanearsonate	0.36%
** Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.12%
*** Dimethylamine salt of (+)-(R)- 2-(4-chloro-2-methylphenoxy) propanoic acid	0.06%
**** Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.02%

INERT INGREDIENTS:	<u>99.44%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

*0.030 lbs. monosodium methanearsonate per gallon or 0.36%.
 Total Arsenic (as elemental) all in water soluble form 0.17%.
 **0.008 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon
 or 0.10%. ***0.004 lbs. (+)-(R)-2-(4-chloro-2-methylphenoxy)
 propanoic acid equivalent per gallon or 0.05%. ****0.002 lbs.
 3,6-dichloro-o-anisic acid equivalent per gallon or 0.02%. Isomer
 Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN
CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE:

GORDON'S® TRIMEC® PLUS Ready-To-Use is intended for residential and ornamental lawns with applications by homeowners only. This product contains four proven herbicides that are effective on hard-to-kill grass and broadleaf weeds. GORDON'S® TRIMEC® PLUS Ready-To-Use is recommended for sparse weed infestations and for follow-up treatments of broadcast herbicide applications. This ready-to-use herbicide is intended for spot treatments only. Weeds that have not emerged at the time of application will not be controlled.

GORDON'S® TRIMEC® PLUS Ready-To-Use controls a wide range of broadleaf weeds including dandelion, spurge, oxalis, crabgrass, dallisgrass, and yellow nutsedge in bermudagrass, Kentucky bluegrass, tall fescue, and

zoysiagrass lawns. A partial list of weeds controlled is shown below.

PARTIAL LIST OF WEEDS CONTROLLED:

Aster; Bedstraw; Black Medic; Buckhorn; Burdock; Chicory; Chickweed; Clovers; Crabgrass (Smooth, Large); Dallisgrass; Dandelion; Docks; Goosegrass; Ground Ivy; Healall; Henbit; Knotweed; Lambsquarters; Lespedeza; Mallows; Morningglory; Nutsedge, Yellow; Oxalis; Peppergrass; Pigweeds; Plantains; Poison Ivy; Poison Oak; Purslane; Ragweed; Sandbur; Sheep Sorrel; Shepherdspurse; Speedwell; Spurge; Wild Carrot; Wild Garlic; Wild Lettuce; Wild Onion; Yarrow; Yellow Woodsorrel.

HOW TO USE:

For Containers with Triggers Sprayers:

GORDON'S® TRIMEC® PLUS Ready-To-Use is applied directly from the container fitted with the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Usually, two (2) pumps of this trigger sprayer will provide adequate coverage of individual weeds. Excessive spraying can injure some lawns.

USE PRECAUTIONS:

- Do not use a hose end sprayer.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
- Do not use this product on St. Augustinegrass, centipedegrass, carpetgrass, creeping fescue,

- dichondra, nor on lawns with desirable clovers.
- Keep children and domestic animals off treated area until this material has been watered into the soil or the grass has dried.
- Do not apply when grass is emerging from dormancy.

WHEN TO APPLY:

Apply when weeds are young and actively growing for the best results. Schedule the application to match the conditions of adequate soil moisture and these air temperatures. For Kentucky bluegrass and tall fescue, apply this product when air temperatures are between 70°F and 80°F. For bermudagrass and zoysiagrass, apply this product when air temperatures are between 70° and 90°F.

Applications made when air temperatures exceed these conditions may result in injury to the lawn. Temporary discoloration may occur with the use of this product.

Weed regrowth may occur because of the growth from stems and rhizomes, environmental conditions, or misapplication. Also, weeds can germinate from seed and emerge throughout the season. Two or more applications at 10-14 day intervals may be required.

Weed control may be reduced when applications are made during periods of inadequate soil moisture. If the soil is dry, then water thoroughly 24 hours before application. Do not water lawns for 24 hours after application.

Newly seeded lawns may be treated after the fourth mowing. Seed can be sown 3 to 4 weeks after application.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area. Keep from freezing.

CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. When using this product wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

First Aid

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of

the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

GORDON'S® and TRIMEC® are registered trademarks of PBI/Gordon Corporation.

738/3-2002 AP031202
EPA REG. NO. 2217-815
EPA EST. NO. 2217-KS-1

Manufactured By



An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com