117



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

AUG 30 2001

Craig Martens PBI/Gordon Corporation P.O. Box 014090 1217 West 17th Street Kansas City, MO 64101

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Dear Mr. Martens:

Subject: Conversion from MCPP to MCPP-p

EH 1346 Weed and Feed

EPA Registration Number: 2217-816 Your Submission Dated June 6, 2001

The Agency is conditionally approving an amendment to the registration of the above-referenced product under the authority of section 3(c)(7)(A) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). Under this amendment the Agency is approving the conversion of the active ingredient mecoprop from its racemic form [2-(2-methyl-4-chlorophenoxy) propionic acid, or applicable salt thereof] to its single isomer form [(+)-R-2-(2-methyl-4-chlorophenoxy) propionic acid, or applicable salt thereof]. The revised labeling and Confidential Statements of Formula submitted in support of the conversion are acceptable with the following provisions:

- 1. This acceptance is based on your certification that no changes have been made to the Confidential Statements of Formula (CSF) or the product labeling other than those specified within the body of your certification statement. No other revisions are being reviewed or considered with this action.
- 2. You may sell or distribute product containing the racemic form of mecoprop, or applicable salt thereof, and bearing the previously approved labeling for 18 months from the date of this letter.

Copies of the revised Confidential Statements of Formula Basic and Alternate Formulations, dated June 6, 2001 has been placed in the Agency's file for the subject product. These CSFs supersede all previously submitted CSFs for this product.

A stamped copy of the labeling is enclosed for your records. Submit one copy of your final printed labeling before you release the product for shipment.

Sincerely yours,

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505C)

Enclosure

EH1346 HERBICIDE

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	42.8%	
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid.		
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)		
INERT INGREDIENTS:	42.7%	
TOTAL	100.0%	

THIS PRODUCT CONTAINS:

3.51 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 35.7% 0.88 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 9.0% 0.29 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 3.0% Contains the single isomer form of Mecoprop-p. Isomer Specific by AOAC Methods

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for Precautionary Statements and Statement of Practical Treatment.

NET CONTENTS: 2.5, 5, 15, 30, and 55 GALLONS

AP EPA REG. NO. 2217-816 Amend EPA Est. No. 2217-KS-X MANUFACTURED BY:

G pbl/gordon
An Employee-Owned Company
1217 West 12th Street
Karress City, Miscoun 64101

ACCEPTED
with COMMENTS
In EPA Letter Dated

AUG 30 2001

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling. When mixing, loading, or applying this product, wear long-sleeved shirt, long pants, socks, shoes, rubber gloves, and eye protection (goggles, face shield, or safety glasses). It is recommended that safety glasses include front, brow, and temple protection.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons in capacity: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more in capacity: Do not open pour from this container. A mechanical system (probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a nonrefillable pesticide container are emptied, the probe must be rinsed before removal.

Statement of Practical Treatment

IF IN EYES: Hold eyelids open and flush with a steady stream of water for at least 15 minutes. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash skin plenty of soap and water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

- SITES -

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is intended to be applied by lawn care operators, maintenance applicators, and service technicians for use in ornamental turf established in institutional, ornamental, and residential/domestic sites. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses, and office buildings. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings. Finally, residential/domestic sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

- APPLICATION SCHEDULES -

Apply this product to emerged broadleaf weeds that are actively growing. Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. If required, the treatment interval between the first and second application should be 14 to 28 days.

- APPLICATION RATES -

Dosage rates and spray volume recommendations of this product are presented in Table 1.

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

The maximum application rate to turf is 1.0 pound 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Spray Volume Amount of Product Species Pints/Acre Fl.oz./1000sq.ft. Gallons/Acre Gallons/1000sq.ft. Cool Season Turf Kentucky Bluegrass 2.0 - 2.250.73 - 0.8320 to 175 0.5 to 4.0 Perennial Ryegrass 2.0 - 2.25 0.73 - 0.8320 to 175 0.5 to 4.0 20 to 175 2.0 - 2.25 0.73 - 0.830.5 to 4.0 Tall Fescue Fine leaf Fescues 2.0 - 2.250.73 - 0.8320 to 175 0.5 to 4.0 Hard Fescues Sheeps Fescue Red Fescue (Chewings Fescue, Slender Creeping Fescue, Spreading Fescue) Creeping Bentgrass 1.0 - 1.25 0.37 - 0.4640 to 175 1.0 to 4.0 Warm Season Turf Common Bermudagrass 2.0 -2.25 0.73 - 0.8320 to 175 0.5 to 4.0 Hybrid Bermudagrass 1.5 - 1.75 0.55 - 0.6420 to 175 0.5 to 4.0 2.0 - 2.25 0.73 - 0.8320 to 175 0.5 to 4.0 Zoysiagrass 1.5 - 1.75 20 to 175 St. Augustinegrass 0.55 - 0.640.5 to 4.0

Table 1. Rate Recommendations for Ornamental Turfgrass.

- WEED SPECTRUM -

This product controls a wide variety of problem weed species including the following:

- WEEDS -			
Bedstraw	Henbit	Purslane	
Black medic	Knotweed	Ragweed	
Buckhorn	Lambsquarters	Sheep sorrel	
Burdock	Lespedeza	Shepherdspurse	
Chicory	Mallow	Speedwell	
Chickweed	Morningglory	Spurge	
Clover	Peppergrass	Wild carrot	
Dandelion	Pigweed	Wild garlic	
Dock	Plantain	Wild lettuce	
Ground ivy	Poison ivy	Wild onion	
Healali	Poison oak	Yarrow	

- USE PRECAUTIONS AND LIMITATIONS -

- Do not apply this product through any type of irrigation system.
- Avoid drift of spray mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants.
 Do not pour spray solutions near these plants.
- Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present.
- Coarse sprays are less likely to drift. Avoid fine mists. Do not spray exposed roots of ornamentals and trees
- Do not apply to newly seeded grasses until well established.
- ◆ Do not broadcast apply when air temperatures exceed 85°F; some injury may be expected with spot treatments when air temperatures exceed 85°F.
- Seed can be sown 3 to 4 weeks after application at recommended rates.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing. Product may crystallize at low temperatures.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.