

002217-00817-103098

Printed: 15:03:13 Tuesday, 25 May, 1999 # 24 / 3096

Systems Integration Group, Inc.

PM 23 Please read instructions on reverse before completing form.	-817	10/30/98 /// Form Ap	proved. OMB No. 2	2070-0060. Approval Expires (- 7)5-31-98
SEPA Environmental	ed States Protection ton, DC 20460	Agency	Registra Amenda Other		umber
Appl	ication for	Pesticide - Sect	tion I		
Company/Product Number 2217-817		2. EPA Product Ma Joanne	anager t. Miller	3. Proposed Classificati	on
Company/Product (Name) EH1350 Weed and Feed		PM# (PM	1 23)	X None ☐ F	Restricted
5. Name and Address of Applicant (Include ZIP Coo PBI/Gordon Corporation PO Box 014090 Kansas City, Missouri 64101 Check if this is a new address	de)	(b)(i), my product i to: EPA Reg. No.		nce with FIFRA Section 3(tical in composition and la	, , ,
Oncok ii kiio io a new adarese		Product Name			
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.	56	Final printed Agency letter "Me Too" Ap	plication.	NOTIFICATION OCT 3 0 1998	
NOTIFICATION of Minor revisions to Directions effective on June 11, 1998. Fertilizer products sol the Fertilizer Regulation Act. We ask to add the requirements of the Department of Agriculture is consistent with the provisions of PR Notice 95-2 labeling or the confidential statement of formula of any false statement to the EPA. I further understated 152.46, this product may be in violation of FIFRATERICA.	d and distribute required statem n Washington. and EPA regulf this product. Ind that if this n	ed in Washington dur ents to comply with p Please refer to pag lations at 40CFR 152 understand that it is a otification is not consi	ing 1999 must be provisions of the ge 4 of the encloaded. 46, and no other a violation of 18 istent with the te	pear the required text as defective Regulation Accorded labeling. This notified the changes have been made U.S.C.Sec. 1001 to willfurns of PR Notice 95-2 and	efined in t and the ication is de to the ally make d 40CFR
	Sec	tion - III			
De Submitted	per If "Ye	age wgt. container	2. Type of C Metro Plas Glas Pap Othe 5. Location of Lab	al ss er er (Specify)	
6. Manner in Which Label is Affixed to Product	Lithograph Paper glued	Other	On Labeling a	accompanying product	
	Stenciled	tion - IV			
Contact Point (Complete items directly below for identification)			ry, to process this a	application.)	
Name Craig Martens	Title	anager of Regulator		Telephone No. (include Area 816/460-6287	•
I certify that the statements I have made on this form a I acknowledge that any knowingly false or misleading s both under applicable law.	tatement may be			6. Date Application Received Sampe	• • • • • •
2. Signature han tens_		ager of Regulatory	Services		
4. Typed Name Craig Martens	5. Date	October 19, 1998	 8		

EH1350 WEED AND FEED

29-3-4

NOTIFICATION OCT 3 0 1998

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.67%
Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	0.34%
Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.06%
INERT INGREDIENTS	98.93%
TOTAL	100,00%

THIS PRODUCT CONTAINS:

11.25 lbs. or 0.56% 2, 4-dichlorophenoxyacetic acid equivalent per ton.
5.60 lbs. or 0.28% 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
0.95 lb. or 0.05% 3, 6-dichloro-o-anisic acid equivalent per ton.
Isomer Specific by AOAC Methods.
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and Statement of Practical Treatment.

NET WEIGHT (15, 16, 17, 18, or 20) POUNDS

AP101998
EPA Reg. No. 2217-817
EPA EST. NO. 2217-KS-1
MANUFACTURED BY:
pbi/scrdon
corporation



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes (moderate) eye injury (irritation). Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

Statement of Practical Treatment

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or area where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

USE PRECAUTIONS:

Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns until after the third mowing.

WHERE TO USE:

EH1350 Weed and Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 29-3-4 fertilizer provides a balanced supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

EH1350 Weed and Feed controls broadleaf weeds commonly found in cool season and warm ceason turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

THE THE POST OF THE PROPERTY O

APPLICATION INSTRUCTIONS:

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and growing actively, preferably in the moming when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

APPLICATION INSTRUCTIONS: (Optional Tabular Format)

Application Schedule	Cultural Practices
1 - 2 days before application	Mow lawn to normal height 1 to 2 days before application.
1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.
Day of application	Apply when weeds are young and growing actively, preferably in the moming when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
1 - 2 days after application	Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, and perennial ryegrass, bermudagrass, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.0 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.

The maximum application rate to turf is 1.0 pound 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to insure uniform coverage and to avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product.

CALIBRATION INSTRUCTIONS:

Refer to the rate table (in the appendix) for the appropriate bag sizes and the area to be treated. For example: (Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) pounds covers up to 5,000 square feet.

- ◆ For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Equipment settings of drop spreaders, rotary spreaders, and other granular equipment may be presented on the container labeling as requested by supplemental registrants.]

Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State
 Department of Agriculture regarding the components in this product is available on the internet at
 http://www.wa.gov/agr/"

	- WEEDS -	
Bedstraw	Knotweed	Ragweed
Black medic	Lambsquarters	Sheep sorrel
Buckhorn	Lespedeza	Shepherdspurse
Burdock	Mallow	Speedwell
Chicory	Morningglory	Spurge
Chickweed	Oxalis	Thistle
Clover	Peppergrass	Wild carrot
Dandelion	Pigweed	Wild garlic
Dock	Plantain	Wild lettuce
Ground ivy	Poison ivy	Wild onion
Healall	Poison oak	Yarrow
Henbit	Purslane	

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

I. Container Packaging and Size:

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size,	Application Rate,	Treated Area for This Bag Size,
Pounds	Pounds/1,000 Sq.Ft.	Square Feet
	3.0	5,000
15	3.2	4,700
	3.4	4,400
	3.6	4,200
	4.0	3,750
	3.0	5,300
16	3.2	5,000
	3.4	4,700
	3.6	4,400
	4.0	4,000
	3.0	5,600
17	3.2	5,300
	3.4	5,000
	3.6	4,700
	4.0	4,250
	3.0	6,000
18	3.2	5,600
	3.4	5,300
ļ	3.6	5,000
	4.0	4,500
	3.0	6,600
20	3.2	6,300
	3.4	5,900
}	3.6	5,600
	4.0	5,000

SPREADER PATTERNS:

- 1. Apply two header strips.
- 2. Operate spreader the longest way of the lawn.
- 3. For drop spreaders, overlap wheels to insure complete coverage.
- 4. Close spreader openings when turning.

II. Previously Approved Fertilizer Grades:

- Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following: 23-3-7 and 29-3-4
- The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	23%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	3%
Soluble Potash (K ₂ O)	7%

III. Advertising claims appropriate for the bag size:

- (Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and other broadleaf weeds.