



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 23, 2017

Mr. Eric Smith
Federal Registration Manager
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, MO 64101

Subject: Notification per PRN 98-10 – Added weeds, alternative warranty statement, marketing claims
Product Name: EH 1350 Weed and Feed
EPA Registration Number: 2217-817
Application Date: November 11, 2016
Decision Number: 523677

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at montague.kathryn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

2217-817

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

01/23/2017

EH1350 WEED AND FEED

EPA Reg. No. 2217-817

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-D (CAS 2008-39-1)	0.67%
Dimethylamine salt of Mecoprop-p (CAS 66423-09-04)	0.17%
Dimethylamine salt of Dicamba (CAS 2300-66-5)	0.06%

INERT OTHER INGREDIENTS	99.10%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 11.2 lb 2, 4-dichlorophenoxyacetic acid equivalent per ton or 0.56%. (CAS 2008-39-1)
- 2.8 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.14%. (CAS 66423-09-04)
- 1.0 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.05%. (CAS 2300-66-5)

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

Guaranteed Analysis: See appendix for example.

Net Contents: _____
EPA Est. No. _____

Company Name and Address:
PBI/Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks,
- chemical-resistant gloves

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns until after the third mowing.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

WHERE TO USE:

EH1350 Weed and Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many other listed weeds commonly found in home lawns, others commonly found in home lawns. In addition, a 29-3-4 X-X-X fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

EH1350 Weed and Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

APPLICATION INSTRUCTIONS:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

APPLICATION INSTRUCTIONS: [Optional Tabular Format]

Application Schedule	Cultural Practices
1 to 2 days before application	Mow lawn to normal height 1 to 2 days before application.

1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.
Day of application	Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
1 to 2 days after application	Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, and perennial ryegrass, bermudagrass, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.0 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

Do not apply more than 4 pounds of product per 1000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 8 pounds of product per 1000 square feet per year. [note to reviewer: the following information may be omitted from the final printed label: 4 pounds of this product contains 0.98 lb 2,4-D (ae), 0.24 lb mecoprop-p (ae), and 0.09 lb dicamba (ae).]

[Alternate optional maximum rate statement:

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [0.98 lb 2,4-D ae, 0.24 lb MCPP-p ae and 0.09 lb. dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [1.95 lb 2,4-D ae, 0.49 lb MCPP-p ae and 0.17 lb dicamba ae per acre].]

[Note to Reviewer: This was moved up from the appendix]

A uniform distribution of product on the treated area is essential. Choose a calm day to insure uniform coverage and to avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area. Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product.

[Note to Supplemental Registrants: A subset of the following list may be used in place of the full list.]

Broadleaf Weeds Controlled [248 Weeds, 233 Weeds CA]			
Amaranth, purple	Creeping Beggarweed*	Morningglory, mexican	Speedwell, bi-lobed
Amaranth, slender	Creeping Jenny	Morningglory, red	Speedwell, common
Amaranth, spiny	(creeping charlie)	Morningglory, tall	Speedwell, corn
Aster, heath	Cupid shaving Brush*	Morningglory, wolly	Speedwell, ivyleaf
Aster, white heath	Cudweed	Mugwort	Speedwell, Persian
Aster, purple	Daisy, English	Mustard, ball	Speedwell, purslane
Aster, white prairie	Daisy, oxeye	Mustard, black	Speedwell, slender
Aster, wild	Dandelion, common	Mustard, blue	Speedwell, snow
Artichoke*	Day flower*	Mustard, hedge	Speedwell, thymeleaf
Austrian fieldcress	Deadnettle	Mustard, Indian,	Spurge, spotted
Bedstraw, Catchweed	Dichondra	Mustard, tansy	Spurge, prostrate
Bedstraw, smooth	Dock, broadleaf	Mustard, tumble	Star of Bethlehem*
Bedstraw, Northern	Dock, curly	Mustard, white	Strawberry, India mock
Bedstraw, pineywoods	Dock, smooth	Mustard, wormseed	Tansy ragwort
Beggarticks, hairy	Dogbane	Nettle, stinging (burning)	Tanweed
Betony, Florida	Dogfennel	Nettle, tall	Thistle, blessed
Bindweed, Field	Dollarweed	Nettle, wood	Thistle, bull
Bindweed, Hedge	(Pennywort)	Parsley-piert	Thistle, Canada
Bitter Wintercress	Dove weed	Parsnip	Thistle, flodman (prairie)

Bittercress, hairy	Elderberry	Pearlwort	Thistle, musk
Bitterweed	Falseflax	Pennycress (frenchweed)	Thistle, platte
Black-eyed Susan	False sunflower	Pepperweed, perennial	Thistle, plumeless
Black Medic	Fiddleneck	Pepperweed, Virginia	Thistle, Russian
Blood flower milkweed	Filaree, redstem	(Peppergrass)	Thistle, scotch
Blue lettuce	Filaree, whitestem	Pigweed, prostrate	Thistle, wavyleaf
Brassbuttons	Fleabane, daisy	Pigweed, redroot	Thistle, yellow (horrible)
Bristly oxtongue	(annual)	Pigweed, Russian	Trailing crownvetch
Broomweed	Fleabane, hairy	Pigweed, smooth	Velvetleaf
Bullnettle	Fleabane, rough	Pigweed, tumble	Vervain, blue
Burclover	Galinsoga, hairy	Pineapple weed	Vervain, hoary
Burdock	Galinsoga, smallflower	Plains coreopsis (tickseed)	Vervain, prostrate
Burweed, lawn (spurweed)	Goldenrod	Plantain, blackseed	Vervain, tall
Buttercup, bulbous	Ground ivy	Plantain, bracted	Vetch, bird
Buttercup, creeping	Groundsel*	Plantain, broadleaf	Vetch, hairy
Buttonweed, common	Gumweed	Plantain, buckhorn	Vetch, milk
(poorjoe)	Hawkweed, meadow	Plantain, hoary	Vetch, narrowleaf
Buttonweed, Virginia	Hawkweed, mouseear	Plantain, narrowleaf	Vetch, wild
Carolina geranium (wild)	Hawkweed, orange	Plantain, slender	Virginia creeper
Carpetweed	Hawkweed, yellow	Plantain, woolly	Western clematis
Catnip	Healall	Poison ivy	Western salsify
Catsear, spotted (false	Heartleaf drymary	Poison oak	Wild buckweat
dandelion)	Hemp	Pokeweed	Wild carrot
Chamber bitter*	Henbit	Prairie Sunflower	Wild four-o'clock
Chickweed, common	Hoary cress	Prickly lettuce (compass plant)	Wild garlic
Chickweed, mouseear	Horsenettle	Prickly sida	Wild lettuce
Chickweed, sticky	Horseweed*	Puncturevine	Wild marigold
Chicory	Jimsonweed	Purslane, common	Wild mustard (yellow rocket)
Cinquefoil	Johnny-Jumpup violet*	Pusley, Brazil	Wild onion
Cinquefoil, rough	Knawel	Pusley, Florida	Wild parsnip
Clover, alyce	Knotweed, prostrate	Pusley, large flower	Wild radish
Clover, crimson	Kochia	Ragweed, bur	Wild rape
Clover, hop	Lambsquarters	Ragweed, common	Wild strawberry
Clover, rabbit foot	Lespedeza	Ragweed, lanceleaf	Wild sweet potato
Clover, red	Mallow, alkali	Ragweed, western	Wild violet
Clover, strawberry	Mallow, bristly	Scarlet pimpernel	Woodsorrel, creeping
Clover, yellow sweet	Mallow, common	Shepard's purse	Woodsorrel, violet
Clover, white sweet	Mallow, dwarf	Smartweed, ladythumb*	Woodsorrel, yellow (Oxalis)
Clover, white	Mallow, venice	Smartweed, pale*	Woolly croton
Cockle	Marcela*	Smartweed, Pennsylvania*	Wormseed
Cocklebur	Matchweed	Smooth chaff-flower	Yarrow, common
Cocklebur, oriental	Mexicanweed	Sorrel, red (sheep)	Yarrow, western
Cocklebur, spiny	Moneywort*	Sowthistle, annual	Yarrow, yellow
Common mullein	Morningglory, bigroot	Sowthistle, spiny	
Corn Spurry*	Morningglory, ivyleaf	Spanish needles	

*Not for use in California.

[Note to Supplemental Registrants: The following language on equipment settings/spreader settings is optional – it may be presented on the container at your discretion.]

SPREADER PATTERNS:

1. Apply two header strips.
2. Operate spreader the longest way of the lawn.
3. For drop spreaders, overlap wheels to insure complete coverage.
4. Close spreader openings when turning.

[Note to Reviewer: This was moved up from the appendix]

CALIBRATION INSTRUCTIONS:

Refer to the rate table (in the appendix) for the appropriate bag sizes and the area to be treated. For example: (Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) pounds covers up to 5,000 square feet.

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Equipment settings of drop spreaders, rotary spreaders, and other granular equipment may be presented on the container labeling as requested by supplemental registrants.]

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling.]

Bag Size, Pounds	Application Rate, Pounds/1,000 Sq.Ft.	Treated Area for This Bag Size, Square Feet	Bag Size, Pounds	Application Rate, Pounds/1,000 Sq.Ft.	Treated Area for This Bag Size, Square Feet
15	3.0	5,000	18	3.0	6,000
	3.2	4,700		3.2	5,600
	3.4	4,400		3.4	5,300
	3.6	4,200		3.6	5,000
	4.0	3,750		4.0	4,500
16	3.0	5,300	20	3.0	6,600
	3.2	5,000		3.2	6,300
	3.4	4,700		3.4	5,900
	3.6	4,400		3.6	5,600
	4.0	4,000		4.0	5,000
17	3.0	5,600			
	3.2	5,300			
	3.4	5,000			
	3.6	4,700			
	4.0	4,250			

[Note to Reviewer: This was moved up from the appendix]

[End Optional Text]

Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- Information regarding the contents and levels of metals in this product is available on the internet at: "<http://www.aapfco.org/metals.htm>."

WEEDS

Bedstraw	Ground ivy	Peppergress	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chicory	Lambsquarters	Poison oak	Wild garlic
Chickweed	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Deck	Oxalis	Shopherspurse	

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- (Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and other broadleaf weeds.
- A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000sq.ft. per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 sq.ft.. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.
- Plus...Kills Dandelions & other broadleaf weeds as listed
- Plus... extended feeding

- Plus... Iron, a key nutrient
- Promotes thick grass
- Contains Trimec® Broadleaf Herbicide
- Stabilized nitrogen
- Dark green grass
- Nitrogen, for a lush green lawn
- Phosphate, improves root growth
- Potash, vital for growth & drought resistance
- Note: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. nitrogen per 1,000 sq.fr.
- This XX lb. bag should cover up to 12,000 sq.ft.
- Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers may cause discoloration.
- Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn. Apply the product to the lawn. Refer to the path charts shown [optional charts for centrifugal/rotary/drop spreaders]
- Close the spreader opening when turning to change directions or when stopping.
- If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.
- This product helps thicken new and existing lawns and stimulates root growth. The granules are specially sized for even and easy application.
- [Contains] Trimec Broadleaf Herbicide
- www.tandninc.com
- Step 4 Fall
- Apply in Fall, Sept/Oct/Nov/Dec
- [Brand] Turf Management
 - Step 1 Crabgrass Preventer Early Spring Feb/Mar/Apr
 - Step 2 Weed and Feed Late Spring Apr/May/June
 - Step 3 Lawn Food Summer Jun/Jul/Aug
 - Step 4 Winterizer Plus Fall Sept/Oct/Nov/Dec

3. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “State Restrictions”]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and

micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: "Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>"
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
 - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida: "Apply only to actively growing turf."
- Florida: "Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use."
- Idaho: "Apply only as directed"
- Maryland: "This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area."
- New Jersey: "In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the "Soil Erosion and Sediment Control Act," P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots."
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: "In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency."
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

- [Advisories for fertilizers containing iron:] "This product may stain concrete and stone, where spilling occurs, sweep off immediately, then flush with water." OR "Sweep all residues from concrete surfaces immediately after application. The iron content in this product will stain sidewalks, patios, driveways, swimming pools and deck coatings if not removed before watering. Neither seller nor distributor assumes any responsibility for damage incurred to any surface other than product's intended use."
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet [and herbicide to control listed weeds]. Please consult your local extension office for questions regarding a regular fertilizer program [suitable for your geographic region]."

3. Container Packaging and Size:

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

Bag Size, Pounds	Application Rate, Pounds/1,000 Sq-Ft.	Treated Area for This Bag Size, Square Feet	Bag Size, Pounds	Application Rate, Pounds/1,000 Sq-Ft.	Treated Area for This Bag Size, Square Feet
15	3.0	5,000	18	3.0	6,000
	3.2	4,700		3.2	5,600
	3.4	4,400		3.4	5,300
	3.6	4,200		3.6	5,000
	4.0	3,750		4.0	4,500
16	3.0	5,300	20	3.0	6,600
	3.2	5,000		3.2	6,300
	3.4	4,700		3.4	5,900
	3.6	4,400		3.6	5,600
17	4.0	4,000		4.0	5,000
	3.0	5,600			
	3.2	5,300			
	3.4	5,000			
	3.6	4,700			
	4.0	4,250			

4. Spreader Information

• **SPREADER PATTERNS:**

1. Apply two header strips.
2. Operate spreader the longest way of the lawn.
3. For drop spreaders, overlap wheels to insure complete coverage.
4. Close spreader openings when turning.

5. Previously Approved Fertilizer Grades:

- Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following: 10-6-4, 23-3-7, 25-3-3 and 29-3-4
- The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	23%
Water Insoluble Nitrogen	5%
Available Phosphate (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	7%

6. Alternate optional maximum rate statement:

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (0.98 lb 2,4-D ae, 0.24 lb MCPP-p ae and 0.09 lb. dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (1.95 lb 2,4-D ae, 0.49 lb MCPP-p ae and 0.17 lb dicamba ae per acre).

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00817.20161111.notif-proposed-highlighted

2. Reason for Issue: ad claims; updated weeds list; state fertilizer language; formatting; correction of typos – inert to other, move CAS numbers

Internal Usage Notes
Associated Container Label: Supplemental Distributors