2217-819 Please read instructions on re	everse before complete	ing form	0711-1	CAT.	Form A	Proved S	3-200	/ 2-0060. Approval Expires	1/9
≎ EPA	Environme	United Sta	ection A	Agend		F	legistration Amendmen	OPP Identifier N	
		Applicati	on for P	estic	de - Sec	tion I			
Company/Product Num	ber 2217-819			2. EP#	N Product Ma Joanne	anager e I. Mille	r	3. Proposed Classificati	on
4. Company/Product (Nam EH1	ne) 352 Weed & Feed	<u> </u>		PM#	<u>(PN</u>	/I 23)		X None R	estricted
 Name and Address of A PBI/Gordon Corporati P.O. Box 014090 Kansas City, Missouri Check if thi 	ion	P Code)		(b)(i), r EPA F	ny product is Reg. No	s similar	or identical ir	ith FIFRA Section 3(c) n composition and labe	
Oneck ii gii	3 IS a new address		Sect	ion - I					
Amendment - Explain	below.				Final printed Agency lette		response to	MOTIFICATION	1
Resubmission in responsible Notification - Explain b		lated			"Me Too" Ap	•		AUG 2 3 2001	
Explanation: Use addition			1 1 0 1						
application schedule, appl Appendix Section, page 7 This notification is consist been made to the labeling willfully make any false st 10 and 40 CFR 152.46, this 12 and 14 of FIFRA.	of the enclosed labe tent with the provision or the confidential satement to the EPA	cling. ons of PR Notatement of I further up	otice 98-10 formula of nderstand th	and EF this pro	'A regulation duct. I unde is notificatio	ns at 40C rstand tha on is not c	FR 152.46, a at it is a viola consistent wi	and no other changes hation of 18 U.S.C.Sec.	1001 to cice 98-
			Secti	on - l					
Material This Product Will Child-Resistant Packaging Yes* No * Certification must	Be Packaged In: Unit Packaging Yes No If "Yes"	No. per	Water S))	Packaging No. per	2.	Type of Contain Metal Plastic Glass	ner	
be submitted	Unit Packaging wgt.	container	Packag		container		Paper Other (Sp	ecify)	1
3. Location of Net Contents In Label Conta		4. Size(s) Re	etail Contain	er			on of Label Dir abel		
6. Manner in Which Label is A	ffixed to Product	Lithogr Paper Stencil	glued ed		Other_				
1.0-1-12-140	man allowed to be a long of the	al a m bld: 22	Secti					t 1 /	
Contact Point (Complete its Name Craic	ims directly below for it. Martens	aenutication o	Title		tacted, if nece Regulatory		Tele	pplication.) phone No. (Include Area of 816-460-6287)	Code)
l certify that the statemer I acknowledge that any k both under applicable lav	nts I have made on this nowingly false or misle		ation attachments ent may be p	thereto a	are true, accu	rate and c	omplete.	6. Date Application Received (Stamped	• • • • •
2. Signaturen Nanz Nan	tous_			er of R	egulatory S	Services			.: .*
4. Typed Name Craig Martens 5.			5. Date	August 10, 2001					

EH1352 WEED & FEED

ACTIVE INGREDIENTS:

2,4-D		0.55 to	0.84%
Mecoprop		0.27 to	0.41%
Dicamba		0.03 to	0.04%
OTHER INGREDIENTS:	*************	99.15 to	98.71%
	TOTAL	16	00.00%

THIS PRODUCT CONTAINS:

11.00 to 16.80 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
5.40 to 8.20 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
0.60 to 0.80 lbs. dicamba (3,6-dichloro-o-anisic acid) equivalent per ton.
Isomer Specific by AOAC Methods

CAS Registry Numbers: 2,4-D CAS# 94-75-7 Mecoprop CAS# 7085-19-0 Dicamba CAS#1918-00-9

IKIMEC.	is a registered	tragemark of	PBI/Gordon (Corporation	Π.

FERTILIZER ANALYSIS	{
Total Nitrogen (N)	xx%
Water Insoluble Nitrogen	xx%
Available Phosphate (P2O5)	x%
Soluble Potash (K ₂ O)	x%

NOTE: Fertilizer grades with nutrient contents expressed as the percentages of N, P_2O_5 and K_2O may include the following: 23-3-7, 29-3-4, and x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State
 Department of Agriculture regarding the components in this product is available on the internet at
 http://www.wa.gov/agr/."

KEEP OUT OF REACH OF CHILDREN

NOTIFICATION

CAUTION

AUG 2 3 2001

See back panel for additional Precautionary Statements and First Aid.

NET WEIGHT: 15, 16, 17, 18, or 20 POUNDS

AP081001 EPA Reg. No. 2217-819 EPA EST. NO. 2217-KS-X MANUFACTURED BY:





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. **If person** is unconscious, do not give anything by mouth and do not induce vomiting.

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

WHERE TO USE:

EH1352 Weed & Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, the fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

EH1352 Weed & Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

WHERE TO USE:

EH1352 Weed & Feed controls a wide range of broadleaf weeds in home lawns. In addition, the fertilizer provides a supply of plant food.

For Use On:

Kentucky bluegrass

Perennial ryegrass

Centipede

Fescue Buffalograss Bermuda

St. Augustine Bahia

falograss Zoysia Ba

USE PRECAUTIONS:

Note to supplemental registrants: For loading rates of 24 - 28 pounds of MUP/ton of fertilizer:

• Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

Note to supplemental registrants: For loading rates of 32 - 37 pounds of MUP/ton of fertilizer:

- ♦ Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
- Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals. [or: Do not exceed the application rate for any area. When applying under trees in an open lawn, walk past trees in a straight path. Circling trees will cause you to pass over some areas twice so that too much material is applied.]
- If overseeding wait 4 weeks after grass seeds sprout before applying this product.

APPLICATION INSTRUCTIONS:

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

An advisory for fertilizers containing iron:

5. This product will stain sidewalks and driveways. Sweep off any product which may have drifted onto these areas.

(Tabular Format)

Application Schedule	Cultural Practices			
1 - 2 days before application	Mow lawn to normal height 1 to 2 days before application.			
1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.			
Day of application	Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.			
1 - 2 days after application	Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.			

HOW MUCH TO APPLY:

Note to supplemental registrants: For loading rates of 24 - 28 pounds of MUP/ton of fertilizer:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

Note to supplemental registrants: For loading rates of 32 - 37 pounds of MUP/ton of fertilizer:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, and centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.

The maximum application rate to turf is xxx (xxx = 1.0, 1.1, 1.3, 1.4, or 1.5) pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

NOTE TO SUPPLEMENTAL REGISTRANTS: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Pounds	Application Rate, Pounds/1000 Sq.Ft.	Treated Area for This Bag Size, Square Feet		
	3.2	4,700		
15	3.6	4,200		
1	4.0	3,700		
	3.2	5,000		
16	3.6	4,400		
	4.0	4,000		
	3.2	5,300		
17	3.6	4,700		
1	4.0	4,200		
	3.2	5,600		
18	3.6	5,000		
	4.0	4,500		
	3.2	6,400		
20	3.6	5,600		
	4.0	5,000		

SPREADERS AND CALIBRATION [or SPREADER CALIBRATIONS AND PATTERNS]:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

Choose one of these application or distribution patterns:

Single Pass - Operate the equipment at intervals corresponding to the width or swath of the spreader.
 For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.

[OR Single Pass - Always apply EH1352 Weed & Feed with a properly calibrated rotary or drop type fertilizer spreader. NEVER APPLY BY HAND. To avoid gaps or streaks, first spread border of lawn. Fill in center with uniform, parallel passes. Spreader settings are approximate. Be sure to check spreader before beginning, adjusting if necessary.]

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.


Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers ___ sq.ft.

- 2. Double Pass Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions or when stopping.
- 3. Cross Hatching Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks or the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

CALIBRATION INSTRUCTIONS:

- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Spreader Settings			
Scott	XX		
Sears	XX		
Lawn Beauty	XX		
Jackson	XX		



LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

- I. Advertising claims appropriate for the bag size:
- (Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) Pounds Covers Up To X,000 Square Feet.
- Controls (Kills) <u>(desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge)</u> and other broadleaf weeds.
- Controls (Kills) <u>(desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge)</u> and over <u>(number)</u> other (more) broadleaf weeds.
- Controls (Kills) over (<u>number</u>) varieties (types) of lawn weeds including (<u>desired weed species from attached listing may be inserted here</u>.).
- Polyon[®] Timed Release Fertilizer.
- Timed-Release for professional results
- Contains Polyon® timed-release fertilizer as used on championship golf courses around the country.
- Use where weeds are actively growing.
- Feeds for up to (time period as designated by subregistrant). (At 72°)
- Quick greening & extended feeding for up to 8 weeks.
- Nitrogen (N) contains timed-release for growing a thick green lawn.
- Phosphorous (P) improves root growth & plant storage capacity.
- Potassium (K) improves plant hardiness & drought resistance.
- (Brand Name) products with Polyon® timed-release nitrogen work differently from other slow-release products. Polyon® timed-released nitrogen releases in response to soil temperature rather than moisture level. Other slow release products release according to the amount of moisture present in the soil...the more moisture the quicker the nutrient release...putting your grass at risk from burning. Because (Brand Name) products contain Polyon® timed-release nitrogen, your grass is fed for up to (time period as designated by subregistrant). (At 72°)
- Fertilizer advisory: Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per
 growing season. When used as directed this product will supplement your regular fertilizer program by
 providing ___ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for
 questions regarding a fertilizer program suitable to your geographical region.
- Helps build a stronger, deeper root system.
- Feeds your lawn with time-released nitrogen for extended greening.
- Encourages early Spring green-up.
- · Guaranteed to banish weeds and green lawns.
- Controls (Kills) weeds and feeds lawns in one step.
- Kills (Controls) all major broadleaf weed types
- Promotes a deep green lawn with stronger roots
- Kills (Controls) weeds Feeds lawns
- Covers (number) sa.ft.
- Try our other (brand name) quality lawn and garden products.
- Satisfaction Guaranteed. If you apply (<u>brand rame</u>) as directed and are not completely satisfied, send
 proof of purchase from this bag and the cash register receipt to the address below. We will promptly
 send your money back.
- ◆ [Alternate Spreader Listing and Graphics] <u>Rotary Spreaders:</u> Ames -see spreader instructions: EV-N-Spread - XX; Ortho - XX; Precision; - XX; Quaker - XX; Republic EZ - XX; Republic EZ Grow - XX; Scotts EasyGreen - XX; Scotts SpeedyGreen - XX; Scotts Speedy Green 1000 - XX; Scotts Speedy Green 2000 - XX; Scotts SpeedyGreen 3000 -XX; True Temper - XX; Vigoro - XX.

<u>Drop Spreaders:</u> Arnes -see spreader instructions, Bandini - XX; Ortho - XX; Precision - XX; Republic EZ - XX; Scotts (PF) - XX; Scotts AccuGreen 1000 - XX; Scotts EvenGreen - XX; Vigoro - XX. . . .

• Controls (Number) weeds! (Listing of weeds may include the following.)

	-	WEEDS -	
Annual yellow sweetclover	Cockle	Musk thistle	Spiny cocklebur
Aster	Cocklebur	Mustard	Spiny sowthistle
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spotted catsear
Bedstraw	Creeping jenny	Narrowleaf vetch	Spotted spurge
Beggarticks	Cudweed	Nettle	Spurweed
Betony, Florida	Curly dock	Orange hawkweed	Stinging nettle
Bindweed, field	Daisy, English	Oriental cocklebur	Strawberry, India mock
Bird vetch	Daisy fleabane	Oxalis	Tall nettle
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall vervain
Bittercress, hairy	Dandelion	Parsnip	Tansy ragwort
Bitterweed	Dichondra	Pearlwort	Tansy mustard
Black-eyed Susan	Dogbane	Pennycress	Tanweed
Black medic	Dogfennel	Pennywort	Thistle
Black mustard	Dollarweed	Pepperweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pigweed	Tumble mustard
Blessed thistle	False dandelion	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	Falseflax	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	False sunflower	Plantain	Venice mallow
Blue vervain	Fiddleneck	Poison ivy	Virginia buttonweed
Bracted plantain	Florida pusley	Poison oak	Virginia creeper
Brassbuttons	Frenchweed	Pokeweed	Virginia pepperweed
Bristly oxtongue	Galinsoga	Poorjoe	Water pennywort
Broadleaf dock	Goathead	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly lettuce	Western clematis
Broomweed	Ground ivy	Prickly sida	Western salsify
Buckhorn	Gumweed	Prostrate knotweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild buckwheat
Bull thistle	Healall	Prostrate vervain	Wild carrot
Bullnettle	Heartleaf drymary	Puncturevine	Wild four-o'clock
Burclover	Heath aster	Purslane, common	Wild garlic
Burdock	Hedge bindweed	Ragweed	Wild geranium
Burning nettle	Hedge mustard	Red sorrel	Wild lettuce
Burweed	Hemp	Redroot pigweed	Wild marigold
Buttercup	Henbit	Redstem filaree	Wild mustard
Buttonweed	Hoary cress	Rough cinquefoil	Wild onion
Canada thistle	Hoary plantain	Rough fleabane	Wild parsnip
Carolina geranium	Hoary vervain	Russian pigweed	Wild radish
Carpetweed	Horsenettle	Russian thistle	Wild rape
Catchweed bedstraw	Jimsonweed	Scarlet pimpernel	Wild strawberry
Catnip	Knawel	Scotch thistle	Wild sweet potato
Catsear	Knotweed	Sheep sorrel	Wild vetch
Chickweed, common	Kochia	Shepherdspurse	Woodsorrel
Chickweed, mouseear	Lambsquarters	Slender plantain	Woolly croton
Chicory	Lespedeza	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Mallow	Smooth dock	Woolly plantain
Clover, crimson	Matchweed	Smooth pigweed	Wormseed
Clover, hop	Mexicanweed	Sorrel	Yarrow
Clover, red	Milk vetch	Sowthistle	Yellow rocket
Clover, strawberry	Morningglory	Spanishneedles	Yellowflower pepperweed :
Clover, sweet	Mouseear hawkweed	Speedwell	and other
Clover, sweet Clover, white	Mugwort	Spiny amaranth	broadleaf weeds
Clovel, wille	MIGEWOIT	Spiny amarahin	broadlear weeds