

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Dr. James L. Kunstman, Ph. D. Product Registration PBI/Gordon, Corporation P.O. Box 014090 Kansas City, Missouri 64101

JUN 12 2008

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request General Label Change (Marketing Claims)

EPA Reg. No. 2217-819

Application Dated May 19, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 05/19/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

Please read instructions on reverse before completing form.	Fo	rm Approved	. OMB No. 207	0-0060.	Approval Expires 2-28-95
United States Environmental Protecti Washington, DC 204			Registration Amendme Other		OPP Identifier Number
Application f	or Pesticide -	Section		 -	<u> </u>
				72.0-	an and Oleverification
Company/Product Number 2217-819	2. EPA Produ	anne I. Mi		3. Pro	oposed Classification
4. Company/Product (Name)	PM#	MINIE 1. IVII		╛┍╾	,
EH-1352 Weed and Feed	101#	(PM 23)			None Restricted
5. Name and Address of Applicant (Include ZIP Code)	6 Evnedited		accordance s	with FIE	FRA Section 3(c)(3)
PBI/Gordon Corporation	i -				position and labeling to:
Post Office Box 014090	EPA Reg. No				ATION
Kansas City, Missouri 64101					
Check if this is a new address	Product Nam	ie	4UL	12	2008
	Section - II				
		orinted labels	in response to		
Amendment - Explain below.		y letter dated			
Resubmission in response to Agency letter dated	"Me To	oo" Applicatio	on.		
Notification Evaluin holow	Other	- Explain belo	^ \W		
Notification - Explain below.					
Explanation: Use additional page(s) if necessary. (For section I and NOTIFICATION: Labeling Notification per PRN 98-10. We ask to approved. These claims may be presented on container labels, advertise the highlighted text on pages 9-12 of the appendix of the labeling. One (1) copy of the proposed labeling is enclosed for your review. This notification is consistent with the provisions of PR Notice 98-10 and EPA	to add advertising clain sement brochures, and regulations at 40 <u>CFR</u> 15	other marke	ting/sales prome	otional r e been m	materials. Please refer to
confidential statement of formula of this product. I understand that it is a violation that if this notification is not consistent with the terms of <u>PR Notice</u> 98-10 and of the product					
action and penalties under Section 12 and 14 of FIFRA.	40 <u>CFK</u> 132.40, mis prod	uct may be in	VIOLATION OF THE	A allu 11	may be subject to enforcement
	ection - III				
Material This Product Will Be Packaged In:	ection - in				
	Vater Soluble Packagir	ng	2. Type of Cont	ainer	·
Yes* Yes	Yes		Metal		,
	No		Plastic		•
			Glass		
	f "Yes" No. p Package wgt. conta		Paper		
be submitted Unit Packaging wgt. container F	ackage wgt. come	11 (C)	Other (S	pecify)_	·
3. Location of Net Contents Information 4. Size(s) Retail C	ontainer	5. Loc	ation of Label D	irections	S
Label Container	•	님	On Label		•
			On Labeling acco	mpanyi	ng product
6. Manner in Which Label is Affixed to Product Lithograph		Other			 .
Paper glued					
Stenciled					
	ection - IV	,			€ 6 (.
Contact Point (Complete items directly below for identification of indivi-		necessary, t			
James L. Kunstman, Ph.D.	: Director of Regul	atory Serv			No. (Include Area Code) 8¥6-460-6292
Certification					3. Date Application
I certify that the statements I have made on this form and all attache I acknowledge that any knowingly false or misleading statement ma both under applicable law.				v.	Received (Stamped)
2. Signature 3. Til	tle				Charles
	Director of Regulate	ory Service	es		1.4.6
	·		·		
4. Typed Name James L. Kunstman, Ph.D. 5. Da	ate May 19, 20	008			2.5 St.

FPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

White - EPA File Copy (original) Yellow - Applicant Copy



1217 WEST 12th STREET
P.O. BOX 014090
KANSAS CITY, MISSOURI 64101-0090

816-460-6232 • FAX: 816-421-2731

May 16, 2008

Document Processing Desk (DIST)
Office of Pesticide Programs (7504C)
US Environmental Protection Agency
Ariel Rios Bldg., 1200 Pennsylvania Ave., NW
Washington, DC 20460-0001

SUBJECT: NOTICE OF SUPPLEMENTAL REGISTRATION OF DISTRIBUTOR

Distributor: Spectrum Group, Division of United Industries Corp., Distributor Company 8845.

Dear Sir or Madam:

Enclosed please find the current Notice of Supplemental Distribution of a Registered Pesticide Product form for the following product(s):

EPA Reg. No.	Distributor No.	Distributor Product Name
2217-819	8845	Sta-Green Phosphorus-Free Winterizer Weed & Feed2 26-0-11
2217-819	8845	Vigoro Ultra Turf Phosphorus-Free Winterizer Weed & Feed2 26-0-11

The forms have been signed by both the registrant, PBI/Gordon Corporation, and the distributor. Copies of the forms have been retained for our files. Please forward notification of acceptance to the distributor.

If you have questions or concerns, please contact me at 816/460-6292 or email: jkunstman@pbigordon.com.

Sincerely,

James L. Kunstman, Ph.D.

Director, Regulatory Services

1/a_ IZ

Electronic Labeling Record

LABELINFORMATION

Record Id#: ELL002477

Attachment Name: 002217-00819.20080519.proposed.pdf

Product Active

Status:

Registration #: 002217-00819

Product: EH1352 WEED & FEED **PC Code:** 029801; 030001; 129046

RM Team: RM 23

Submission 05/19/2008

Date:

Receipt Date:

Version: Initial

CD Enclosed?: Yes

Comments: proposed

Application Correspondence:

Attachment:

002217-00819.20080519.proposed.pdf

SUBMISSION INFORMATION

Status: Pending

Submisssi 830175

on BC:

Action 332

Code:

OPPIN Receipt Date: 05/19/2008

Action Comments: Notification for marketing claims

Reviewer Comments:

User Reviews & Response Letters

Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL				
EPA Registration #	Date Submitted to EPA	Electronic file name		
2217-819	5/19/2008	002217-00819.20080519.proposed		

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Jan 7 Janta	05/19/2008
Signature	Date
James L. Kunstman, Ph.D. Name (typed)	
Director, Regulatory Services	, .
Title	

EH1352 WEED & FEED

ACTIVE INGREDIENTS:

2,4-D		0.55 to	0.84%
Mecoprop-p		0.14 to	0.21%
Dicamba		0.03 to	0.04%
OTHER INGREDIENTS:	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	99.28 to	98.91%
	TOTAL	1.	00 000/

TOTAL

100.00%

THIS PRODUCT CONTAINS:

11.00 to 16.80 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
2.80 to 4.20 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.

0.60 to 0.80 lbs. dicamba (3,6-dichloro-o-anisic acid) equivalent per ton.

Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

CAS Registry Numbers:	
2,4-D CAS# 94-75-7	
Maconron n CAS# 16484	•

Mecoprop-p CAS# 16484-77-Dicamba CAS#1918-00-9

FERTILIZER ANALYSIS	
Total Nitrogen (N)	xx%
Water Insoluble Nitrogen	xx%
Available Phosphate (P ₂ O ₅)	x%
Soluble Potash (K ₂ O)	x%

NOTE: Fertilizer grades with nutrient contents expressed as the percentages of N, P_2O_5 and K_2O may include the following: 23-3-7, 29-3-4, and x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at http://www.wa.gov/agr/."

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid.

NET WEIGHT: 14, 15, 16, 17, 18, or 20 POUNDS

AP080807 EPA Reg. No. 2217-819 EPA EST NO. 2217-KS

EPA EST. NO. 2217-KS-2 MANUFACTURED BY:

An Employer-Owned Company
1217 Wost 1216 Street
Herosic City, Moserie (MSH)
1-800-821-7925

NOTIFICATION

JUN 12 2008

Page 1 of 12



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

	First Aid		
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 		
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. 		
 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 			
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.			

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until dusts have settled.

STORAGE AND DISPOSAL

[Note to reviewer: Text within brackets may be optional.]

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number [(1-800-XXX-XXXX)] for

Page 2 of 12

disposal instructions. Never place unused product down any indoor or outdoor drain.

WHERE TO USE:

EH1352 Weed & Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, the fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

EH1352 Weed & Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

Alternate format for "WHERE TO USE" Instructions:

WHERE TO USE:

EH1352 Weed & Feed controls a wide range of broadleaf weeds in home lawns. In addition, the fertilizer provides a supply of plant food.

For Use On:

Kentucky bluegrass

Perennial ryegrass

Centipedegrass

Fescue

Bermudagrass

St. Augustinegrass

Buffalograss

Zoysiagrass

Bahiagrass

USE PRECAUTIONS:

Note to supplemental registrants: For loading rates of 24 - 28 pounds of MUP/ton of fertilizer:

 Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

Note to supplemental registrants: For loading rates of 32 - 37 pounds of MUP/ton of fertilizer:

- ♦ Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
- Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals. [or: Do not exceed the application rate for any area. When applying under trees in an open lawn, walk past trees in a straight path. Circling trees will cause you to pass over some areas twice so that too much material is applied.] If overseeding wait 4 weeks after grass seeds sprout before applying this product.

APPLICATION INSTRUCTIONS:

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

An advisory for fertilizers containing iron:

5. This product will stain sidewalks and driveways. Sweep off any product which may have drifted onto these areas.

(Tabular Format)

Application Schedule	Cultural Practices
1 - 2 days before application	Mow lawn to normal height 1 to 2 days before application.
1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.
Day of application	Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
1 - 2 days after application	Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

Controls (Number) weeds! (Listing of weeds may include the following.)

- BROADLEAF WEEDS -			
Annual yellow sweetclover	Cockle	Musk thistle	Spiny cocklebur
Aster '	Cocklebur	Mustard	Spiny sowthistle
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spotted catsear
Bedstraw	Creeping jenny	Narrowleaf vetch	Spotted spurge
Beggarticks	Cudweed	Nettle	Spurweed
Betony, Florida	Curly dock	Orange hawkweed	Stinging nettle
Bindweed, field	Daisy, English	Oriental cocklebur	Strawberry, India mock
Bird vetch	Daisy fleabane	Oxalis	Tall nettle
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall vervain
Bittercress, hairy	Dandelion	Parsnip -	Tansy ragwort
Bitterweed	Dichondra	Pearlwort	Tansy mustard
Black-eyed Susan	Dogfennel	Pennycress	Tanweed
Black medic	Dollarweed	Pepperweed	Thistle
Black mustard	False dandelion	Pigweed	Trailing crownvetch
Blackseed plantain	Falseflax	Pineywoods bedstraw	Tumble mustard
Blessed thistle	False sunflower	Plains coreopsis	Tumble pigweed
Bloodflower milkweed	Fiddleneck	Plantain	Velvetleaf
Blue lettuce	Florida pusley	Poison ivy	Venice mallow
Blue vervain	Frenchweed	Poison oak	Virginia buttonweed
Bracted plantain	Galinsoga	Pokeweed	Virginia creeper
Bristly oxtongue	Goldenrod	Poorjoe	Virginia pepperweed
Broadleaf dock	Ground ivy	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Gumweed	Prickly lettuce	Western clematis
Broomweed	Hairy fleabane	Prickly sida	Western salsify
Buckhorn	Hawkweed	Prostrate knotweed	White mustard
Buckhorn plantain	Healall	Prostrate pigweed	Wild aster
Bulbous buttercup	Heartleaf drymary	Prostrate spurge	Wild buckwheat
Bull thistle	Heath aster	Prostrate vervain	Wild carrot
Bullnettle	Hedge bindweed	Puncturevine	Wild four-o'clock
Burclover	Hedge mustard	Purslane, common	Wild garlic
Burdock	Hemp	Ragweed	Wild geranium
Burning nettle	Henbit	Red sorrel	Wild lettuce
Burweed	Hoary cress	Redroot pigweed	Wild marigold
Buttercup	Hoary plantain	Redstem filaree	Wild mustard
Buttonweed	Hoary vervain	Rough cinquefoil	Wild onion
Canada thistle	Horsenettle	Rough fleabane	Wild parsnip
Carolina geranium	Jimsonweed	Russian pigweed	Wild radish
Carpetweed	Knawel	Russian thistle	Wild rape
Catchweed bedstraw	Knotweed	Scarlet pimpernel	Wild strawberry
Catnip	Kochia	Scotch thistle	Wild sweet potato
Catsear	Lambsquarters	Sheep sorrel	Wild vetch
Chickweed, common	Lespedeza	Shepherdspurse	Woodsorrel
Chickweed, mouseear	Mallow	Slender plantain	Woolly croton
Chicory	Matchweed	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth dock	Woolly plantain
Clover, crimson	Milk vetch	Smooth pigweed	Wormseed
Clover, hop .	Morningglory	Sorrel	Yarrow
Clover, red	Mouseear hawkweed	Sowthistle	Yellow rocket
Clover, strawberry	Mugwort	Spanishneedles	
Clover, strawberry	Milliamorr	Speedwell	Yellowflower pepperweed
Clover, sweet	•	•	and other
Ciover, willte		Spiny amaranth	broadleaf weeds or

HOW MUCH TO APPLY:

Note to supplemental registrants: For loading rates of 24 - 28 pounds of MUP/ton of fertilizer:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 2.8 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 2.8 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.

Note to supplemental registrants: For loading rates of 32 - 37 pounds of MUP/ton of fertilizer:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, and centipedegrass: Apply 2.8 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 2.8 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.

The maximum application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.5 lbs 2,4-D acid equivalent per acre per application).

The maximum number of broadcast applications is limited to 2 per year.

The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (3.0 lbs 2,4-D acid equivalent per acre).

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

NOTE TO SUPPLEMENTAL REGISTRANTS: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Pounds	Application Rate, Pounds/1,000 Sq.Ft.	Treated Area for This Bag Size, Square Feet
	2.8	5,000
Ì	3.2	4,400
14	3.6	3,800
	4.0	3,500
	3.2	4,700
15	3.6	4,200
	4.0	3,700
-	3.2	5,000
16	3.6	4,400
	4.0	4,000
	3.2	5,300
17	3.6	4,700
	4.0	4,200

Page 6 of 12

	3.2	5,600
18	3.6	5,000
	4.0	4,500
	3.2	6,400
20	3.6	5,600
	4.0	5,000

SPREADERS AND CALIBRATION [or SPREADER CALIBRATIONS AND PATTERNS]:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

[Note to Supplemental Registrants: Choose one or more of these application or distribution patterns]

 Single Pass - Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.

[OR Single Pass - Always apply EH1352 Weed & Feed with a properly calibrated rotary or drop type fertilizer spreader. NEVER APPLY BY HAND. To avoid gaps or streaks, first spread border of lawn. Fill in center with uniform, parallel passes. Spreader settings are approximate. Be sure to check spreader before beginning, adjusting if necessary.]

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

- 2. **Double Pass** Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions or when stopping.
- 3. **Cross Hatching** Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks or the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

[Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers ____ sq.ft.]

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

CALIBRATION INSTRUCTIONS:

- For 2.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 360 square feet and adjust to apply at the rate of 1 pound of product per 360 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Spreader Settings	
Scott	XX
Sears	XX
Lawn Beauty	XX
Jackson	XX

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED EXCEPT AS OTHERWISE PROVIDE BY LAW. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

I. Advertising claims appropriate for the bag size:

- (Fourteen, Fifteen, Sixteen, Seventeen, Eighteen, or Twenty) Pounds Covers Up To X,000 Square Feet.
- Controls (Kills) <u>(desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge)</u> and other broadleaf weeds [and many other listed broadleaf weeds].
- Controls (Kills) <u>(desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge)</u> and over <u>(number)</u> other (more) broadleaf weeds.
- Controls (Kills) over (<u>number</u>) varieties (types) of lawn weeds including (<u>desired weed species from attached listing may be inserted here.</u>).
- Polyon® Timed Release Fertilizer.
- Timed-Release for professional results
- Contains Polyon[®] timed-release fertilizer as used on championship golf courses around the country.
- · Use where weeds are actively growing.
- Feeds for up to (time period as designated by subregistrant). (At 72°)
- Quick greening & extended feeding for up to 8 weeks.
- Nitrogen (N) contains timed-release for growing a thick green lawn.
- Phosphorous (P) improves root growth & plant storage capacity.
- Potassium (K) improves plant hardiness & drought resistance.
- (Brand Name) products with Polyon® timed-release nitrogen work differently from other slow-release products. Polyon® timed-released nitrogen releases in response to soil temperature rather than moisture level. Other slow release products release according to the amount of moisture present in the soil...the more moisture the quicker the nutrient release...putting your grass at risk from burning. Because (Brand Name) products contain Polyon® timed-release nitrogen, your grass is fed for up to (time period as designated by subregistrant). (At 72°F)
- Fertilizer advisory: Typically a lawn will utilize 1 to 4 pounds [lbs.] of nitrogen per 1,000 square feet per growing season. When used [applied] as directed this product will supplement your regular fertilizer program by providing [this product provides] ____ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.
- · Helps build a stronger, deeper root system.
- Feeds your lawn with time-released nitrogen for extended greening.
- Encourages early Spring green-up.
- Guaranteed to banish weeds and green lawns.
- Controls (Kills) weeds and feeds lawns in one step.
- Kills (Controls) all major broadleaf weed types
- Promotes a deep green lawn with stronger roots
- · Kills (Controls) weeds Feeds lawns
- Covers (number) sq.ft. [Approximately xx acre]
- Try our other (brand name) quality lawn and garden products.
- Satisfaction Guaranteed. If you apply (<u>brand name</u>) as directed and are not completely satisfied, send
 proof of purchase from this bag and the cash register receipt to the address below. We will promptly
 send your money back
- [Alternate Spreader Listing and Graphics]
 Rotary Spreaders: Ames -see spreader instructions; EV-N-Spread XX; Ortho XX; Precision XX;
 Quaker XX; Republic EZ XX; Republic EZ Grow XX; Scotts EasyGreen XX; Scotts SpeedyGreen XX; Scotts Speedy Green 1000 XX; Scotts Speedy Green 2000 XX; Scotts SpeedyGreen 3000 XX; True Temper XX; Vigoro XX.
 - <u>Drop Spreaders:</u> Ames -see spreader instructions; Bandini XX; Ortho XX; Precision XX; Republic EZ XX; Scotts (PF) XX; Scotts AccuGreen 1000 XX; Scotts EvenGreen XX; Vigoro XX.

- Greens in 72 Hours
- Kills over 200 Weeds
- Timed-Release for Extended Feeding
- Promotes Thick, Green Growth and Controls Dandelion, White Clover, Plantain, Chickweed, Spurge & Many Other* Broadleaf Weeds (*See back panel for a complete list of weeds Controlled)
- "To Do" List ✓ Mow Lawn ✓ Measure Lawn ✓ Apply to wet grass at recommended rate
- **CTR, or Composite Timed-Release, is your guarantee that a QUICK & LONG LASTING FEEDING will be systematically delivered to your lawn.
- GENERAL TIPS
 - Determine size of area to be treated measure length and width in feet, then multiply to determine coverage area in sq. ft. (length x width). Next, subtract non-lawn areas.
 - Use the spreader setting chart to determine the appropriate setting:
 - ♦ Treat as directed thorough coverage is important.
- Visit us on the Internet or call our Customer Satisfaction Hotline 1-800-XXX-XXXX
- Delivers a long lasting feeding without burning, for a stronger, greener lawn.
- · Kills weeds while feeding your lawn.
- Questions: for further information on this product, call 1-800-XXX-XXXX
- Why does your lawn need (Brand) Weed & Feed? (Brand) Weed & Feed is specially formulated to kill over xxx (number) lawn weeds and give your grass the nutrients it needs for thick growth. Our timedrelease formula keeps your grass greener longer and helps build strong, hardy root systems.
- · Promotes quick spring green-up.
- Controls fall lawn weeds & protects turf from the stress of winter.
- · Enjoy a Beautiful Weed-Free Lawn
- TURF NOTES: Proper mowing prevents weeds. The best way to control weeds is to mow the lawn properly. Close mowing or scalping encourages weeds. Mow your lawn as tall as recommended for the variety or grass type.
- Encourages Quick Greening and Provides Extended Feeding.
- Controls over 150 weeds including Dandelion, Chickweed, Knotweed, Plantain, Henbit, Thistle and many more.
- Apply Anytime When Broadleaf Weeds Are Actively Growing.
- Apply Anytime Weeds Are Growing
- Controls over 175 weeds [as listed]
- Formulated with Nutralene® Enriches Soil—Feeds the Lawn
- Kills Fall Lawn Weeds!
- · Controls fall lawn weeds and prepares your lawn for the stress of winter
- · Promotes a strong, deep root system in the fall for a quicker green-up in the spring
- Time-releasesd nitrogen feeds and enriches the soil resulting in a thick, green, strong lawn
- · Feeds and enriches the soil resulting in a thick, green, strong lawn
- · Consistent even feeding for up to 3 months and guaranteed not to burn
- Apply in the Fall
- · Feeds up to 3 Months
- If you apply Schultz Supreme Winterizer Weed and Feed as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below.
 We will refund the purchase price you paid.

Why use ACE Green Turf® Phosphorus FREE Weed & Feed?

ACE Green Turf Phosphorus FREE Weed & Feed is specially formulated to complement the
phosphorus already present in some soils and give your turf the additional primary nutrients
needed for a good turf feeding. A soil test can help you determine if your soil has an adequate
supply of phosphorus and if this is the best product for your soil type. Proper watering and mowing
practices along with regular feedings will help promote a beautiful green lawn.

TIPS FOR REDUCING NUTRIENT RUNOFF

- Avoid blowing leaves or grass clippings into the street or driveway. The leaves, grass, and other plant material, which are carried by storm water to our lakes eventually break down and release nutrients that may harm our waterways.
- Cut grass where it borders the street or any paved surface so the clippings are blown back into the yard, or bagged for compost.
- Do not pile grass clippings and leaves in street gutters.
- · Sweep or rake up organic materials found in the street.
- Leave a buffer of natural vegetation near shorelines to prevent erosion and retain nutrients.
- Plant vegetation on bare slopes to help prevent soil erosion.
- Contains Composite Timed-Release (CTR)**Nitrogen for Extended Feeding
- APPLY ANYTIME WEEDS ARE ACTIVELY GROWING
- Fertilizer advisory: A lawn typically uses 1 to 4 pounds [lbs.] of nitrogen per 1,000 square feet per growing season. When applied as directed, this product provides _____ lbs. of nitrogen per 1,000 sq. ft. Consult your local Extension Service for questions regarding a fertilizer program suitable to your geographical region.
 - PROJECT GUIDE SPRING...FALL
- o Early Spring—BAHIA WEED & FEED
- o Late Spring—BAHIA WEED & FEED
- o Summer—SUMMER FERTILIZER
- o Fall-TURF FERTILIZER
- Vigoro offers complete annual lawn care programs for all areas. See your local store associate or visit Vigoro.com for more information about a customized program that is right for your area and turf type.
- · Apply anytime broadleaf weeds are actively growing
- Apply in Spring and Fall
- Enjoy a Beautiful Weed-Free Lawn
- Special guidelines may apply to the specific timing of fall fertilizer applications to Centipedegrass and St. Augustinegrass lawns. Contact your local Extension Service for information specific to your geographic region.
- Typically a lawn utilizes 1-4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed this product provides 0.79 lb of nitrogen per 1,000 sq ft."
- Zero phosphate fertilizer specially formulated for lawns located near lakes & streams
- Promotes Quick Spring Green-Up & a Thick, Dense Lawn
- Fertilzer grades: 26-0-11
- Made in the USA
- Money Back Guarantee
- Turf Note

Apply Early For Best Results!

Many winter weeds such as chickweed, henbit, clover and dandelion germinate from seed in early fall while temperatures are still warm. Weed & Feed applications made in October and early November will control these weeds while they are young and tender."

• The Sta-Green Promise

Your Satisfaction Guaranteed or Your Money Back!

If you use Sta-Green Phosphorus-Free Winterizer Weed & Feed2 26-0-11 with slow-release nitrogen as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below. We will refund the purchase price you paid:

Project Guide

For a thicker, greener more beautiful lawn, follow Vigoro's Annual Lawn Care Program. It's easy to follow and can be started at any time by simply applying the product that corresponds with the current season! Vigoro offers complete annual lawn care programs for all areas. See your local store associate or visit Vigoro com for more information about a customized program that is right for your area and turf type.