

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

James L. Kunstman, Ph.D. Director of Regulatory Services PBI/Gordon Corporation P.O. Box 014090 Kansas City, MO 64101

OCT 1 6 2008

SUBJECT: Application for Pesticide Notification – Add Alternate Warranty EH1369 Weed and Feed EPA Reg. No. 2217-827 Application Dated October 29, 2007

Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Terri Stowe of my staff at 703-305-6117.

Sincerely,

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before com	2/11
United States Environmental Protec Washington, DC 20	tion Agency Amendment
Application	for Pesticide - Section I
1. Company/Product Number 2217-827	2. EPA Product Manager 3. Proposed Classification Joanne I. Miller
4. Company/Product (Name) EH1369 Weed and Feed	PM# Product Manager—Team 23
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No
Check if this is a new address	OCT 1 \$ 2008
· · · · · · · · · · · · · · · · · · ·	Section - II
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.	Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below.
This addition is presented as highlighted text on p This notification is consistent with the provisions of <u>PR Notice 98-10</u> labeling or Confidential Statement of Formula of this product. I under	which may be presented on the container label of a supplemental distributor. page 9 of the enclosed label. and EPA regulations at 40 <u>CFR</u> 152.46, and no other changes have been made to the rstand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to t with the terms of <u>PR Notice 98-10</u> and 40 <u>CFR</u> 152.46, this product may be in violation of
1. Material This Product Will Be Packaged In:	······································
Child-Resistant Packaging Yes* No * Certification must Unit Packaging Yes No No If "Yes" No. per	Water Soluble Packaging 2. Type of Container Yes Metal No Plastic If "Yes" Glass If "Yes" Paper
be submitted Unit Packaging wgt. container	Package wgt. container Other (Specify)
3. Location of Net Contents Information 4. Size(s) Retain 1 Label Container 6. Manner in Which Label is Affixed to Product Lithograp Paper glu Paper glu	Ail Container 5. Location of Label Directions On Label On Label On Labeling accompanying product Other ued
	Section - IV
	Individual to be contacted, if necessary, to process this application.
James L. Kunstman, Ph.D.	Title Telephone No. (Include Area Code) Director of Regulatory Services 81-5-460-6292
Certificati I certify that the statements I have made on this form and all atta I acknowledge that any knowingly false or misleading statement both under applicable law.	ion 6. Date Application tachments thereto are true, accurate and complete. 8. Date Application it may be punishable by fine or imprisonment or 8. Date Application Stamped; 8. Date Application
2. Signature	3. Title Director of Regulatory Services
4. Typed Name 5 James L. Kunstman, Ph.D.	5. Date October 29, 2007

		Previous	are	obsòlete.

•

OCT 1 6 2008

EH1369 WEED & FEED

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	0.723%
Mecoprop-p, dimethylamine salt	0.148%
Dichlorprop-P, dimethylamine salt	0.088%
INERT INGREDIENTS:	<u>99.041%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS:

2,4-dichlorophenoxyacetic acid equivalent, 0.601%.

(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent, 0.122% (+)-(R)-2-(2,4-dichlorophenoxy) propionic acid equivalent, 0.074%.

Isomer Specific by AOAC Methods

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

FERTILIZER ANALYSIS	
Total Nitrogen (N)	xx%
Water Insoluble Nitrogen	xx%
Available Phosphate (P ₂ O ₅)	x%
Soluble Potash (K ₂ O)	x%

NOTE: Fertilizer grades with nutrient contents expressed as the percentages of N, P,O, and K,O may include the following: 23-3-7, 29-3-4, and X-X-X. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

Fertilizer Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- Information regarding the contents and levels of metals in this product is available on the internet at http://aapfco.org/metals.htm [or http://www.regulatory-info-sc.com]

For state registrations in Kentucky, add the following:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of Lbs.) pounds of nitrogen per 1,000 square feet (Optional P_2O_5 and K_2O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid.

NET WEIGHT: 15, 16, 17, 18, 20, 32 or 48 POUNDS

AP102907 EPA Reg. No. 2217-827 EPA EST. NO. 2217-KS-X MANUFACTURED BY: pbl/coador conponation wned Comnan

1217 West 12th Street Konsas City, Missouri 64101 Telephone: 800-821-7925

002217-00827.20071029.notif

Page 1 of 9 EPA Reg. No. 2217-827

100.000% CAS Registry Numbers:

2.4-D, DMA salt CAS#2008-39-1 MCPP-p, DMA salt CAS#66423-09-4 2,4-DP-p, DMA salt CAS#104786-87-0



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if inhaled, or if absorbed through skin, or if swallowed. Avoid breathing dust. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid				
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 			
lf on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. 			
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 			
	ct container or label with you when calling a poison control center or doctor or going OPTIONAL: You may also contact 1-800-xxx-xxxx for emergency medical treatment			

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-XXX-XXXX for disposal instructions. Never place unused product down any indoor or outdoor drain.

WHERE TO USE:

EH1369 Weed & Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, the fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests, or State Extension Service recommendations or professional advice.

EH1369 Weed & Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

Page 3 of 9 EPA Reg. No. 2217-827 Controls (Number) weeds! (Listing of weeds may include the following.)

- BROADLEAF WEEDS -

-		
	Annual yellow sweetclover	Cockle
	Aster	Cockleb
	Austrian fieldcress	Commo
1	Bedstraw	Creepin
1	Beggarticks	Cudwee
	Betony, Florida	Curly d
	Bindweed, field	Daisy, I
	Bird vetch	Daisy fl
	Bitter wintercress	Daisy, o
1	Bittercress, hairy	Dandeli
	Bitterweed	Dichon
	Black-eyed Susan	Dogban
	Black medic	Dogfen
	Black mustard	Dollarw
	Blackseed plantain	Elderbe
	Blessed thistle	False da
	Bloodflower milkweed	Falsefla
	Blue lettuce	False su
1	Blue vervain	Fiddlen
	Bracted plantain	Florida
	Brassbuttons	Frenchy
	Bristly oxtongue	Galinso
	Broadleaf dock	Goathea
	Broadleaf plantain	Golden
	Broomweed	Ground
	Buckhorn	Gumwe
	Buckhorn plantain	Hairy fl
-	Bulbous buttercup	Hawkw
	Bull thistle	Healall
	Bullnettle	Heartle
	Burclover	Heath a
	Burdock	Hedge l
	Burning nettle	Hedge
	Burweed	Hemp
	Buttercup	Henbit
	Buttonweed	Hoary c
	Canada thistle	Hoary p
	Carolina geranium	Hoary v
	Carpetweed	Horsen
1	Catchweed bedstraw	Jimsony
	Catnip	Knawel
1	Catsear	Knotwe
	Chickweed, common	Kochia
	Chickweed, mouseear	Lambsc
	Chicory	Lesped
ļ	Cinquefoil	Mallow
	Clover, crimson	Matchw
	Clover, hop	Mexica
Ì		Milk ve
	Clover, red Clover, strawberry	Mornin
	Clover, sweet	Mousee
	Clover, white	Mugwo

bur on mullein ng jenny ed lock English fleabane oxeye lion dra ne nel veed erry andelion ax unflower leck pusley weed oga ad irod ivy eed leabane reed af drymary aster bindweed mustard cress plantain vervain lettle weed eed quarters leza weed inweed etch igglory ear hawkweed ort

Musk thistle Mustard Narrowleaf plantain Narrowleaf vetch Nettle Orange hawkweed Oriental cocklebur Oxalis Parsley-piert Parsnip Pearlwort Pennycress Pennywort Pepperweed Pigweed Pineywoods bedstraw Plains coreopsis (tickseed) Plantain Poison ivy Poison oak Pokeweed Poorioe Prairie sunflower Prickly lettuce Prickly sida Prostrate knotweed Prostrate pigweed Prostrate spurge Prostrate vervain Puncturevine Purslane, common Ragweed Red sorrel Redroot pigweed Redstem filaree Rough cinquefoil Rough fleabane Russian pigweed Russian thistle Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smooth dock Smooth pigweed Sorrel Sowthistle Spanishneedles Speedwell Spiny amaranth

Spiny cocklebur Spiny sowthistle Spotted catsear Spotted spurge Spurweed Stinging nettle Strawberry, India mock Tall nettle Tall vervain Tansy ragwort Tansy mustard Tanweed Thistle Trailing crownvetch Tumble mustard Tumble pigweed Velvetleaf Venice mallow Virginia buttonweed Virginia creeper Virginia pepperweed Wavyleaf bullthistle Western clematis Western salsify White mustard Wild aster Wild buckwheat Wild carrot Wild four-o'clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild mustard Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild vetch Woodsorrel Woolly croton Woolly morningglory Woolly plantain Wormseed Yarrow Yellow rocket Yellowflower pepperweed and other broadleaf weeds

6/11

002217-00827.20071029.notif

Page 4 of 9 EPA Reg. No. 2217-827

USE PRECAUTIONS:

- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
- Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.
- If overseeding wait 4 weeks after grass seeds sprout before applying this product.

APPLICATION INSTRUCTIONS:

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

An advisory for fertilizers containing iron:

5. This product will stain sidewalks and driveways. Sweep off any product which may have drifted onto these areas.

(Tabular Format)

Application Schedule	Cultural Practices
1 - 2 days before application	Mow lawn to normal height 1 to 2 days before application.
1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.
Day of application	Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
1 - 2 days after application	Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Cultivars vary in tolerance to this product. Do not apply this product to 'Floratam' St. Augustinegrass.

The maximum application rate to turf is 1.1 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

DISTRIBUTION AND COVERAGE:

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

002217-00827.20071029.notif

Page 5 of 9 EPA Reg. No. 2217-827 Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.).

NOTE TO SUPPLEMENTAL REGISTRANTS: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

	length
x	width
=	Total Sq.Ft.
-	Non-Lawn Areas
=	Total Sq. Ft. To Be Treated

ILLUSTRATION

Bag Size, Pounds	Application Rate, Pounds/1000 Sq.Ft.	Treated Area for This Bag Size, Square Feet
	3.2	4,700
	3.4	4,400
15	3.6	4,200
	4.0	3,700
	3.2	5,000
	3.4	4,700
16	3.6	4,400
	4.0	4,000
	3.2	5,300
	3.4	5,000
17	3.6	4,700
	4.0	4,200
	3.2	5,600
	3.4	5,300
18	3.6	5,000
	4.0	4,500
	3.2	6,400
	3.4	5,900
20	3.6	5,600
	4.0	5,000

SPREADERS AND CALIBRATION:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product. *Choose one of these application or distribution patterns:*

- 1. Single Pass Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.
- Double Pass Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions or when stopping.
- 3. **Cross Hatching** Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks or the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

Page 6 of 9 EPA Reg. No. 2217-827

CALIBRATION INSTRUCTIONS:

Calibrate (check) your spreader before beginning your application and adjust the settings if necessary.

- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 295 square feet and adjust to apply at the rate of 1 pound of product per 295 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Spreader Settings		
Scott	XX	
Sears	XX	
Lawn Beauty	XX	
Jackson	XX	

IMPORTANT: It is important to follow directions as indicated because over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

[Alternate Warranty Statement for use by Supplemental Distributor]

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

002217-00827.20071029.notif

Page 7 of 9 EPA Reg. No. 2217-827

APPENDIX

- I. Advertising claims appropriate for the bag size:
 - (Fifteen, Sixteen, Seventeen, Eighteen, Twenty, Thirty Two, or Forty Eight) Pounds Covers Up To X,000 Square Feet.
 - Controls (Kills) (desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge) and over (number) other (more) broadleaf weeds.
 - Helps build a stronger, deeper root system.
 - Feeds your lawn with time-released nitrogen for extended greening.
 - Encourages early Spring green-up.
 - Timed-release fertilizer for professional results
 - Use when broadleaf weeds are actively growing.
 - Feeds for up to (time period as designated by subregistrant). (At 72°)
 - Quick greening & extended feeding for up to 8 weeks.
 - Nitrogen (N) contains timed-release for growing a thick green lawn.
 - Phosphorous (P) improves root growth & plant storage capacity.
 - Potassium (K) improves plant hardiness & drought resistance.
 - For a thick, green weed-free lawn.1
 - Greens fast & lasts.
 - ♦ Apply in Fall!
 - Apply in Fall When Weeds are Growing!
 - Kills Fall Weeds and Greens the Lawn!
 - Apply Anytime Weeds are Growing!
 - Kills Weeds & Greens the Lawn!
 - (Before Photo) and (After Photo) for the Expert Gardener Weed & Feed
 - (Before Photo) and (After Photo) for the Expert Gardener Winterizer Weed & Feed
 - Kills 200+ Weeds While Providing Winter Lawn Protection and Early Spring Green-Up!
 - Kills 200+ Weeds including Dandelions, Clover and Other Common Lawn Weeds
 - ◆ Formulated with Scotts[®] Poly-S[®] Technology for Longer Lasting Greening and Feeding
 - Do not use on Dichondra, Carpetgrass or Floratam, a type of St. Augustinegrass.
 - Treats & Feeds 5,000 sq ft (1/8 acre = 5,445 sq ft)
 - Expert Gardener[®] Fertilizer Program (chart)
 - For Best Results! Use the Expert Gardener[®] Fertilizer Program!
 - Controls 200+ Weeds! Including:
 - When to Apply: Apply when broadleaf weeds are actively growing in the early fall.
 - When to Apply: Apply when broadleaf weeds are actively growing in spring or early fall
 - Spreading Tips: Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts[®] spreader. For additional spreader settings, please call 1-800-543-TURF (8873). Calibrate your spreader before beginning your application and adjust settings if necessary.
 - What to Expect: Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.
 - Expert Gardener Guarantee
 - When you choose Expert Gardener[®] products, you can be assured of high quality and performance in your lawn and garden. If for any reason you are not completely satisfied with this product, we will refund your money. Please mail your name, address and proof of purchase (dated receipt and package UPC/barcode) to the address below. Also, to ensure quality control, please write us a brief note explaining your experience with our product and reason for seeking a refund.

Page 8 of 9 EPA Reg. No. 2217-827

- Contact Us: Consumer Helpline 1-866-369-2533
- Information regarding the contents and levels of metals in this product is available on the Internet at: <u>http://www.regulatory-info-sc.com</u>

(.

- Nitrogen (N) For Deep Green Color
- Phosphorus (P) For Strong Roots
- Potassium (K) For Overall Plant Health
- Application: Anytime Weeds Are Growing
- ◆ Contains Poly-S[®] Slow-Release Technology for Longer-Lasting Greening
- Kills and [&] Controls over 200 Broadleaf Weeds, Including Dandelions, Clovers, Plantains (full list of broadleaf weeds on back panel).
- Non-Burning Formulation (when used as directed)
- Contains:
 - 0.723% 2,4-D, DMA Salt
 - ♦ 0.148% MCPP-p, DMA Salt
 - 0.088 2,4-DP-p, DMA Salt
- Optional Warranty Statements:
 - NOTICE: Buyer assumes all risks of use, storage or handling of this product.
 - **GUARANTEE:** If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

 ζ

Page 9 of 9 EPA Reg. No. 2217-827