

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENHON PENHODES AND TOXIC SUBSTANCES

JAN 3 0 2009

James L. Kunstman, Ph. D. Product Registration PBI/Gordon, Corporation P.O. Box 014090 Kansas City, Missouri 64101

SUBJECT:

Application for Pesticide Notification (PRN 98-10 and 2008-1)

Request Marketing Claims, Directions for Use & Fertilizer Grades

EPA Reg. No. 2217-827

Application Dated August 20, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 08/20/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader
Registration Division (7505P)

Registration Division (7505P)

Office of Pesticide Programs

Please read instructions on re	verse before completing form.		Form A	oproved. OMB No. 2	070-0060. Approval Expires 2-28-95
≎ EPA	Environmental P	States Protection An, DC 20460	Agency	Registration Amenda	
	Applia	ation for P	esticide - Sec	<u> </u>	
Company/Product Numb			2. EPA Product M		3. Proposed Classification
4. Company/Product (Nam			PM#		None Restricted
PBI/Gordon Corporati Post Office Box 0140 Kansas City, Missouri	90		Product Manager—Team 23 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or ide NOTHE GASTO Med labeling to: EPA Reg. No. Product Name JAN 3 0 2009		
		Sect	ion – II		
Amendment – Explain Resubmission in respo	onse to Agency letter dated		Final printe Agency lett "Me Too" A	•	to
We ask to add a fertilize We ask to add referral s Addition of advisory sta This notification is consistent wi have been made to the labeling of statement to the EPA. I further u	r the confidential statement of for	r to page 9 of the ed sequentially. It is ticide Registrat 10 and PR Notice mula of this products not consistent with RA and I may be s	e labeling. Please refer to page 10 ion Notice (PRN) 200 (2008-1 and EPA regulant I understand that it is the terms of PR Notice 10 independent of PR Notice 10 independent in its the terms of PR Notice 10 independent in its independent in it	08.1: Please refer to pations at 40 CFR 156.15 a violation of 18 U.S. ce 98-10 and PR Notice	0. 156.80. 152.46 and no other changes C.Sec. 1001 to willfully make any false e 2008-1 and the requirements of 40 CFR
Material This Product Will	Re Packaged In:	Secti	1011 – 111		
Child-Resistant Packaging Yes* No * Certification must be submitted	Unit Packaging Yes No If "Yes" No. per Unit Packaging wgt. contain	Y If "Yes			il cic s
3. Location of Net Contents In Label Conta		(s) Retail Contair	ner	5. Location of Labo On Label	
6. Manner in Which Label is A	P	ithograph aper glued tenciled	Other		
			ion – IV		
	ms directly below for identificat		o be contacted, if nec	essary, to process th	
Name James L. K	(unstman, Ph.D.	Title Dire	ector of Regulator	v Services	Telephone No. (Include Area Code) 816-460-6292
I certify that the statemer	Cer ats I have made on this form an nowingly false or misleading st	tification d all attachments	thereto are true, acc	urate and complete.	6. Date Application Received (Stamped)
		Direct	tor of Regulatory	Services	
4. Typed Name James L. Kunstman, Ph.D.			August 20, 200	8	
EPA Form 8570-1 (Rev. 3-94)	Previous editions are obsolete		Whi	te - EPA File Copy ((original) Yellow - Applicant Copy

30813

Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL					
me	EPA Registration # Date Submitted to EPA Electron				
30820	8/20/2008	2217-827			
30		2217-827			

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Ja I Zata	08/20/2008
Signature	Date
James L. Kunstman, Ph.D. Name (typed)	
Director, Regulatory Services	• .
Title	

EH1369 WEED & FEED

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	0.723%
Mecoprop-p, dimethylamine salt	0.148%
Dichlorprop-P, dimethylamine salt	0.088%
INERT INGREDIENTS:	99.041%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

2.4-dichlorophenoxyacetic acid equivalent, 0.601%. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent, 0.122% (+)-(R)-2-(2.4-dichlorophenoxy) propionic acid equivalent, 0.074%. Isomer Specific by AOAC Methods
TRIMEC* is a registered trademark of PBI/Gordon Corporation.

CAS Registry Numbers:

2.4-D, DMA salt CAS#2008-39-1 MCPP-p, DMA salt CAS#66423-09-4 2.4-DP-p, DMA salt CAS#104786-87-0

NOTHICATION

JAN 3 0 2009

FERTILIZER ANALYSIS	
Total Nitrogen (N)	xx%
Water Insoluble Nitrogen	xx%
Available Phosphate (P ₂ O ₅)	x%
Soluble Potash (K ₂ O)	x%

NOTE: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅ and K₂O may include the following: 23-3-7, 29-3-4, and X-X-X. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

Fertilizer Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement:
 "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- Information regarding the contents and levels of metals in this product is available on the internet at http://www.negultuory-info-se.com

For state registrations in Kentucky, add the following:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing *(No. of Lbs.) pounds of nitrogen per 1,000 square feet (Optional P₂O₅ and K₂O levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid.

AP082008 EPA Reg. No. 2217-827 EPA EST. NO. 2217-KS-X MANUFACTURED BY: Componition Ast Employee Option Componition Telephone: 800-821-7925 Page 1 of 10 002217-00827.20080820.proposed Page 1 of 10 EPA Reg. No. 2217-827





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if inhaled, or if absorbed through skin, or if swallowed. Avoid breathing dust. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants; socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

	First Aid				
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 				
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. 				
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 				
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [OPTIONAL: You may also contact 1-800-xxx-xxxx for emergency medical treatment information.]					

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. [To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems. [Do not contaminate water when disposing of equipment washwater.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if availables of

If partly filled: Call your local solid waste agency or 1-800-XXX-XXXX for disposal instructions.

60f13

Never place unused product down any indoor or outdoor drain.

WHERE TO USE:

EH1369 Weed & Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, the fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests, or State Extension Service recommendations or professional advice.

EH1369 Weed & Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass, and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

Controls (Number) weeds! (Listing of weeds may include the following.)

	- BROAD	LEAF WEEDS -	
Annual yellow sweetclover	Cockle	Musk thistle	Spiny cocklebur
Aster	Cocklebur	Mustard	Spiny sowthistle
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spotted catsear
Bedstraw	Creeping jenny	Narrowleaf vetch	Spotted spurge
Beggarticks	Cudweed	Nettle	Spurweed ·
Betony, Florida	Curly dock	Orange hawkweed	Stinging nettle
Bindweed, field	Daisy, English	Oriental cocklebur	Strawberry, India mock
Bird vetch	Daisy fleabane	Oxalis	Tall nettle
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall vervain
Bittercress, hairy	Dandelion	Parsnip	Tansy ragwort
Bitterweed	Dichondra	Pearlwort	Tansy mustard
Black-eyed Susan	Dogbane	Pennycress	Tanweed
Black medic	Dogfennel	Pennywort	Thistle
Black mustard	Dollarweed	Pepperweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pigweed	Tumble mustard
Blessed thistle	False dandelion	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	Falseflax	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	False sunflower	Plantain	Venice mallow
Blue vervain	Fiddleneck	Poison ivy	Virginia buttonweed
Bracted plantain	Florida pusley	Poison oak	Virginia creeper
Brassbuttons	Frenchweed	Pokeweed .	Virginia pepperweed
Bristly oxtongue	Galinsoga	Poorjoe	Wavyleaf bullthistle
Broadleaf dock	Goathead ·	Prairie sunflower	Western clematis
Broadleaf plantain	Goldenrod	Prickly lettuce	Western salsify
Broomweed	Ground ivy	Prickly sida	White mustard
Buckhorn	Gumweed	Prostrate knotweed	Wild aster
Buckhorn plantain	Hairy fleabane	Prostrate pigweed .	Wild buckwheat
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild carrot
Bull thistle	Healall	Prostrate vervain	Wild four-o'clock
Bullnettle	Heartleaf drymary	Puncturevine	Wild garlic
Burclover	Heath aster	Purslane, common	Wild geranium
Burdock	Hedge bindweed	Ragweed	Wild lettuce
Burning nettle	Hedge mustard	Red sorrel	Wild marigold
Burweed	Hemp	Redroot pigweed	Wild mustard
Buttercup	Henbit	Redstem filaree	Wild onion
Buttonweed	Hoary cress	Rough cinquefoil	Wild parsnip
Canada thistle	Hoary plantain	Rough fleabane	Wild radish
Carolina geranium	Hoary vervain	Russian pigweed	Wild rape
Carpetweed	Horsenettle	Russian thistle	Wild strawberry
Catchweed bedstraw	Jimsonweed	Scarlet pimpernel	Wild sweet potato
Catnip	Knawel	Scotch thistle	Wild vetch
Catsear	Knotweed	Sheep sorrel	Woodsorrel
Chickweed, common	Kochia	Shepherdspurse	Woolly croton
Chickweed, mouseear	Lambsquarters	Slender plantain	Woolly morningglory
Chicory	Lespedeza	Smallflower galinsoga	Woolly plantain
Cinquefoil	Mallow	Smooth dock	Wormseed
Clover, crimson	Matchweed	Smooth pigweed	Yarrow
Clover, hop	Mexicanweed	Sorrel	Yellow rocket ''
Clover, red	Milk vetch	Sowthistle	Yellow Egwer pepperweed
Clover, strawberry	Morningglory	Spanishneedles	and other control
Clover, sweet	Mouseear hawkweed	Speedwell	broadleaftweeds & co
Clover, white	Mugwort	Spiny amaranth	((((



USE PRECAUTIONS:

- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers
- ♦ Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
- ♦ Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.
- If overseeding wait 4 weeks after grass seeds sprout before applying this product.

APPLICATION INSTRUCTIONS:

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

An advisory for fertilizers containing iron:

5. This product will stain sidewalks and driveways. Sweep off any product which may have drifted onto these areas.

(Tabular Format)

Application Schedule	Cultural Practices		
1 - 2 days before application	Mow lawn to normal height 1 to 2 days before application.		
1 day before application	Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering.		
Day of application	Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.		
1 - 2 days after application	Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.		

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application. Cultivars vary in tolerance to this product. Do not apply this product to 'Floratam' St. Augustinegrass.

The maximum application rate to turf is 1.1 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

DISTRIBUTION AND COVERAGE:

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Page 5 of 10 EPA Reg. No. 2217-827



Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.).

NOTE TO SUPPLEMENTAL REGISTRANTS: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

	length	
X	width	
=_	Total Sq.Ft.	· ILLUSTRATION .
	Non-Lawn Areas	
=_	Total Sq. Ft. To Be Treated	
ľ	•	

Bag Size, Application Rate, Pounds Pounds/1000 Sq.Ft.		Treated Area for This Bag Size, Square Feet		
3.2		4,700		
3.4		4,400		
15	3.6	4,200		
	4.0	3,700		
	3.2	5,000		
	3.4	4,700		
16	3.6	4,400		
	4.0	4,000		
3.2		5,300		
		5,000		
17	3.6	4,700		
	4.0	4,200		
3.2		5,600		
	3.4	5,300		
18	3.6	5,000		
4.0		4,500		
	3.2	6,400		
	3.4	5,900		
20	3.6	5,600		
4.0		5,000		

SPREADERS AND CALIBRATION:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product. Choose one of these application or distribution patterns:

- 1. **Single Pass** Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.
- 2. **Double Pass** Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions or when stoppings are
- 3. Cross Hatching Reduce the application rate by one-half of the recommended fale, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the Wheel tracks of the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

CALIBRATION INSTRUCTIONS:

Calibrate (check) your spreader before beginning your application and adjust the settings if necessary.

- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 295 square feet and adjust to apply at the rate of 1 pound of product per 295 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Spreader Settings	
Scott	XX
Sears	XX
Lawn Beauty	XX
Jackson	XX

IMPORTANT: It is important to follow directions as indicated because over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

[Alternate Warranty Statement for use by Supplemental Distributor]

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law. Manufacturer DISCLAIMS ANY LIABILITY FOR CGMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

APPENDIX

I. Advertising claims appropriate for the bag size:

- ◆ (Fifteen, Sixteen, Seventeen, Eighteen, Twenty, Thirty Two, or Forty Eight) Pounds Covers Up To <u>X</u>,000 Square Feet.
- Controls (Kills) (desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge) and over (number) other (more) broadleaf weeds.
- Helps build a stronger, deeper root system.
- Feeds your lawn with time-released nitrogen for extended greening.
- Encourages early Spring green-up.
- Timed-release fertilizer for professional results
- Use when broadleaf weeds are actively growing.
- Feeds for up to (time period as designated by subregistrant). (At 72°)
- Quick greening & extended feeding for up to 8 weeks.
- Nitrogen (N) contains timed-release for growing a thick green lawn.
- Phosphorous (P) improves root growth & plant storage capacity.
- Potassium (K) improves plant hardiness & drought resistance.
- For a thick, green weed-free lawn.1
- Greens fast & lasts.
- ◆ Apply in Fall!
- Apply in Fall When Weeds are Growing!
- Kills [Listed] Fall Weeds [As Listed] and Greens the Lawn!
- Apply Anytime Weeds are Growing!
- Kills [Listed] Weeds & Greens the Lawn!
- (Before Photo) and (After Photo) for the Expert Gardener Weed & Feed
- (Before Photo) and (After Photo) for the Expert Gardener Winterizer Weed & Feed
- Kills 200+ [Listed] Weeds [As Listed] While Providing Winter Lawn Protection and Early Spring Green-Up!
- Kills 200+ [Listed] Weeds [As Listed] including Dandelions, Clover and Other Common Lawn Weeds
- ◆ Formulated with Scotts[®] Poly-S[®] Technology for Longer Lasting Greening and Feeding
- Do not use on Dichondra, Carpetgrass or Floratam, a type of St. Augustinegrass.
- Treats & Feeds 5,000 sq ft (1/8 acre = 5,445 sq ft)
- ◆ Expert Gardener[®] Fertilizer Program (chart)
- For Best Results! Use the Expert Gardener Fertilizer Program!
- ◆ Controls 200+ [Listed] Weeds [As Listed]! Including:
- When to Apply: Apply when broadleaf weeds are actively growing in the early fall.
- When to Apply: Apply when broadleaf weeds are actively growing in spring or early fall
- Spreading Tips: Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts* spreader. For additional spreader settings, please call 1-800-543-TURF (8873). Calibrate your spreader before beginning your application and adjust settings if necessary.
- What to Expect: Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.
- Expert Gardener Guarantee
- When you choose Expert Gardener* products, you can be assured of high quality and performance in your lawn and garden. If for any reason you are not completely satisfied with this product, we will refund your money. Please mail your name, address and proof of purchase (dated receipt and package URC/barcode) to the address below. Also, to ensure quality control, please write us a brief note explaining your experience with our product and reason for seeking a refund.
- Contact Us: Consumer Helpline 1-866-369-2533

- Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.gggdatory-info-sc.com
- Nitrogen (N) For Deep Green Color
- ♦ Phosphorus (P) For Strong Roots
- Potassium (K) For Overall Plant Health
- Application: Anytime Weeds Are Growing
- ◆ Contains Poly-S" Slow-Release Technology for Longer-Lasting Greening
- Kills and [&] Controls over 200 Broadleaf Weeds, Including Dandelions, Clovers, Plantains (full list of broadleaf weeds on back panel).
- Non-Burning Formulation (when used as directed)
- Contains:
 - 0.723% 2,4-D, DMA Salt
 - ♦ 0.148% MCPP-p, DMA Salt
 - 0.088 2,4-DP-p, DMA Salt
- ◆ Optional Warranty Statements:
 - NOTICE: Buyer assumes all risks of use, storage or handling of this product.
 - **GUARANTEE:** If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.
- New. Built-In Spreader.
- Kills [Listed] Weeds
- Builds Strong, Deep Roots
- Water Smart(TM) Formula [graphic of water drop]
- Makes the Most of Every Drop(TM)
- Improves Lawn's Ability to Absorb Water & Nutrients
- 28-4-2 [fertilizer grade]
- 28-0-3 [fertilizer grade]
- Apply to a wet lawn
- Well-fed lawns are more efficient at using available water.
- Feeding with Scotts® Water Smart^(TM) Formula [products] help(s) grow a thick lawn with more roots to better absorb water and nutrients. This helps protect your lawn against heat, drought and other stresses.
- Feed regularly for your best lawn.
 - [graphic of grass density] One Annual Feeding
 - [graphic of grass density] 2-3 Annual Feedings
 - [graphic of grass density] 4 Annual Feedings
- Scotts No-Quibble Guarantee® If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.
- Get expert advice for your lawn NOW! Want information on what to use and when? Sign up today to get free expert advice specifically for your lawn
- [computer mouse icon] Visit our website: www.scotts.com
- [phone icon] Consumer Helpline: 1-800-543-TURF (8873)
- Easy To Apply
 - Step 1: Flip: Remove seal under cap
 - Step 2: Tilt: Tilt bottle as shown
 - Step 3: Feed: To prevent over application, tilt bottle as shown when applying the product. Walk at a steady pace and shake bottle side-to-side using your wrist. Apply product uniformly in rows.
 - Step 4: Clean-Up: To keep lawn products from reaching storm sewers, sweep or blowers stray particles from hard surfaces back onto the lawn.
- Tip: 1.000 sq ft (20x50 ft) is approximately half the size of a tennis court [tennis court mage]
- Questions/Comments. [computer mouse icon] www.scotts.com 1-800-543-TURF (8873) [phone icon] (8873) [phone icon]

130F13

- Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn.
- Kills 200+ [Listed] Weeds [As Listed]
- Promotes thick, green lawns
- · Feeds for a thick, green lawn
- Grows strong roots
- Contains time-released nitrogen for prolonged, even feeding

	Early Spring	Mid/Late Spring	Summer	Fall	Grass Type
Morman	Crabgrass Preventer	Weed & Feed	Lawn Food	Lawn Food or Weed & Feed	Bluegrass, Rye, Fescues, Bermuda, Zoysia
Sowhern	Crabgrass Preventer	Weed & Feed	Lawn Food	Lawn Food or Weed & Feed	Bermuda, Fescues, Bahia
hera	Nothing until grass starts to green up	St. Augustine Weed & Feed or Lawn Food	Lawn Food	Lawn Food or St. Augustine Weed & Feed	St. Augustine, Centipede, Zoysia

NOTIFICATION

JAN 3 0 2009



