

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

James L. Kunstman, Ph.D. PBI/Gordon Corporation P.O. Box 014090 Kansas City, MO 64101

MAY 1 5 2013

Subject:

Notification per PR Notice 2007-4: Update Container Disposal Instructions

EH-1391 Herbicide

EPA Reg. No. 2217-844

Application Dated: April 18, 2013

Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 for the subject product.

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label changes requested fall within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions regarding this letter, contact me at (703) 347-0257 or rudick.maggie@epa.gov.

Sincerely,

Maggie Rudick

Herbicide Branch

Registration Division (7505P)

Please read instructions on reverse be	efore completin (m.		For	n Approved	d. (3 No. 207	'0-0060. Approv	val Expires 2-28-95
	\t, ,				Registration	ODD	ldentifier Number
♦EPA En	United Sta Nvironmental Prot	•	Vaency		i		
	Washington, D		Agency		Amendme	nt	
	9 ,				Other		
	Application	on for P	esticide - S	ection	.I		
1. Company/Product Number			2. EPA Produc	_		3. Proposed	Classification
2217	7-844		Kal	ole Bo D	avis	_	
4. Company/Product (Name)	l laubiatala		PM#		T 05	X None	Restricted
***	Herbicide				-Team 25		
 Name and Address of Applicant PBI/Gordon Corporation 	(Include ZIP Code)		6. Expedited i				ection 3(c)(3) and labeling to:
Post Office Box 014090			EPA Reg. No.			iii compositioi	rand labeling to.
Kansas City, Missouri 64101			_				
Check if this is a ne	ew address		Product Name				
		Sect	ion – II		 		
Amendment - Explain below.			Final pr	inted labels	s in response to	NOTIFICA	Tion
·							-
Resubmission in response to A	gency letter dated		"Me To	o" Applicati	on.	MAY 15	2013
Notification - Explain below.			Other -	Explain be	low.	·	-0,0
Explanation: Use additional page(s	s) if necessary. (For section	n I and Section	on II.)	e-mail to	jkunstman@pbig	ordon.com FA	X: 816-421-2731
I. NOTIFICATION: N							
Request to update the				hin the g	uidelines of F	PR Notice 20	07-4. Please
refer to pages 14 and 1 Notification of label change per PR N				· 1 · r	D.M. (1 2007		CEDA
regulations at 40 CFR §§ 156.10, 150							
Statement of Formula for this produc	t. I understand that it is a v	violation of I	18 U.S.C. Sec. 10	01 to willfu	ılly make any fal	lse statement to	EPA. I further
understand that if the amended label may be in violation of FIFRA and I m							136, this product
may be in violation of 111 1d2 and 1 m	ay so subject to enjoy come.		on – III	12	<i>ana</i> 17 <i>oj</i> 111 10		
Material This Product Will Be Pack	aged In:						
Child-Resistant Packaging Unit Pa	ackaging	Water 9	Soluble Packaging	,	2. Type of Cont	ainer	
Yes*	es	Ye	es		Metal ☑ Plastic		
No No	io	No.)		Glass		
* Certification must If "Yes"	No. per	If "Yes"	No. pe	r	Paper		
	ackaging wgt. container	Packag	e wgt. Contai	ner	Other (S	Specify)	
3. Location of Net Contents Information	n 4. Size(s) Re	etail Containe		5. Lo	cation of Label D		
Label Container	1 ' '	uart, ½ Gallo			On Label		•
					On Labeling acco	ompanying prod	ugi:
6. Manner in Which Label is Affixed to	Product Lithogr	raph	☐ Ot	ther		<u>e c</u>	<u>t. t</u> .
	Paper	glued				eeët	
	Stencil						
			on – IV		· · · · · · · · · · · · · · · · · · ·	<u> </u>	. C. C.
Contact Point (Complete items direct	tly below for identification of	Title	be contacted, if i	necessary,		,	lude Area Code)
Name James L. Kunstma	an, Ph.D.		ctor of Regula	tory Serv			0-6292
	Certifica	ation	-				A ^c pplication
I certify that the statements I have I acknowledge that any knowingly						Recei	ved
both under applicable law.	.a.oo or morodaling stateme	-	aonable by fille	- imprisor	mioni oi		Stamped)
2. Signature		3. Title					
(lan I Z		Directo	or of Regulato	ory Servic	ces		
4. Typed Name		5. Date					
James L. Kunstma	an, Ph.D.	J. Date	April 18, 2	013			
			. , –				

An Employee-Owned Company

1217 W. 12th STREET · P.O. BOX 014090 KANSAS CITY, MISSOURI 64101-0090 816-421-4070 · 1-800-821-7925 FAX: 816-474-0462

April 18, 2013

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Mr. Kable Bo Davis, (PM-25)

Dear Mr. Davis:

NOTIFICATION: Notification of label change per PR Notice 2007-4. EH-1391 Herbicide (EPA Reg. No. 2217-844).

I. Notification of label change per PR Notice 2007-4.

Request to update the Storage and Disposal section of sublabel 2 within the guidelines of PR Notice 2007-4. Please refer to pages 14 and 15 of the enclosed draft labeling.

II. Certification Statement:

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penaties under sections 12 and 14 of FIFRA.

III. Enclosures:

MAY 1 5 2013

- 1. Application for Pesticide Amendment (EPA Form 8570-1)
- 2. One (1) copy of the draft labeling with annotations
- 3. One (1) copy of the draft labeling without notations.
- 4. One CD containing electronic copies of the draft labeling.

If you have any questions, please call me at 816-460-6292 or contact me at jkunstman@pbigordon.com

Sincerely,

James. L. Kunstman, Ph.D. Director of Regulatory Services

EH-1391 HERBICIDE

EPA Reg. No. 2217-844

Three sublabels represent the entire master label of EH-1391 Herbicide. Please refer to the appropriate section of the labeling as shown as follows:

Section	•			
Sublabel 1				
Sublabel 2				
Sublabel 3		 · · · · · · · · · · · · · · · · · · ·		
Appendix				
Document Control Information		 		

NOTIFICATION MAY 1 5 2013

SUBLABEL 1

EH-1391 HERBICIDE

NOTIFICATION
MAY 1 5 2013

EPA Reg. No. 2217-844

Alternate Brand Names: GroundWork Ready-To-Use 2% Weed & Grass Killer, Gordon's Pronto Fast Acting Weed & Grass Killer, Gordon's Pronto Fast Acting Weed & Grass Killer - Foams on Contact

ACTIVE INGREDIENT:

Isopropylamine salt of glyphosate*

OTHER INGREDIENTS:

100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Wash thoroughly with soap and water.

If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes Remove contact lenses, if present, after the first 5 minutes, then continues Call a poison control center or doctor for treatment advice. 	
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. 	ζες 1. τςςς
if swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or Do not give anything by mouth to an unconscious person. 	r doctór.

^{*}N-(phosphonomethyl)glycine

Environmental Hazards

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater and rinsate.

[Environmental Hazards (alternate per PRN 2008-1)

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.]

Physical Or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Keep children and pets out of treated areas until sprays have dried.

1. How It Works

Facts and Tips

- EH-1391 Herbicide is a general purpose herbicide designed as a ready-to-use spray to control a broad spectrum of emerged annual and perennial weed species. EH-1391 Herbicide will control unwanted species of grass, sedge, broadleaf weeds, and certain woody species around trees, shrubs and fences.
- This product kills plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves down to the roots.
- This product is nonselective and will kill or injure all vegetation contacted by the spray. Do not spray plants or grasses you like—they may die too. Not recommended for spot weed control in lawns since this product kills lawn grasses.
- Any product not absorbed by plants breaks down into natural materials without moving in or on the soil to untreated plants.
- This product has no soil residual activity and thus will not affect nearby untreated plants.
- Plants that have not emerged at the time of application will not be controlled.
- Best results will be obtained when the weed species are actively growing.

2. Tough Weed Control (Annuals, Perennials, and Brush) CONTROLS WEEDS AND GRASSES INCLUDING:

Lawn and Turfgrass Spec EH-1391 Herbicide will prov		pression) of these lawn and to	urfgrass species:
Bahiagrass Bentgrass, colonial, creeping and redtop Bermudagrass, common and hybrid Bluegrass, annual, rough,	Bromegrass, downy and smooth Buffalograss* Canarygrass, reed Carpetgrass* Centipedegrass	Fescue, chewings, hard, red, tall and sheep Foxtail, green, yellow, creeping, meadow and giant Johnsongrass	Orchardgrass Paspalum spp Ryegrass, annual (Italian), perennial, and hardy St. Augustinegrass Zoysiagrass or Japanese
and Kentucky		Kikuyugrass	lawngrass
* Not for use in California			

Woody Species			;
EH-1391 Herbicide	will provide full or partial c	ontrol (suppression) of thes	e woody species:
Alder	Elderberry	Mock orange*	Quaking aspen
Blackberry	Honeysuckle	Manzanita	Raspberry
Barberry*	Hickory	Pine	Sumac

Ceanothus spp.	Kudzu	Poison ivy	Trumpetcreeper
Cherry	Lantana	Poison oak	Wild blackberry
Coyotebrush	Maple	Poison hemlock	Western red cedar*
Dewberry	Multiflora rose	Oak	Willow
Elm			
* Not for use in Califo	ornia		

Annual and Perennial We			
EH-1391 Herbicide will pro	vide full or partial control (su	ppression) of these annual and	perennial weed species:
Artichoke, Jerusalem	Fiddleneck	Oleander*	Tansy ragwort
Barnyardgrass	Field bindweed	Oxalis	Tansy mustard
Beggarweed	Field pennycress	Pampasgrass	Texas panicum
Bittercress	Field sandbur	Pennsylvania smartweed	Thistle
Black medic	Filaree	Pennywort	Timothy
Broadleaf plantain	Florida pusley	Pigweed	Toadflax
Buckhorn plantain	Garden spurge	Plantain	Torpedograss
Bull thistle	Goosegrass	Prickly lettuce	Trailing lantana
Burclover	Green foxtail	Primrose	Tumble mustard
Canada thistle	Guineagrass	Prostrate spurge	Vaseygrass
Chickweed	Hemp sesbania	Puncturevine	Velvetleaf
Clover	Henbit	Purple cudweed	Virginia-creeper
Cocklebur	Horsenettle	Purple nutsedge	White clover
Common groundsel	Horseradish	Purslane	Whitetop
Common mullien	Horseweed/marestail	Quackgrass	Wild barley
Common ragweed	Ironweed	Ragweed	Wild carrot
Coral-bead	Knapweed	Red clover	Wild geranium
Crabgrass	Knawel	Red sorrel*	Wild morningglory
Creeping beggarweed	Knotweed	Redroot pigweed	Wild mustard
Creeping Charlie	Lambsquarters	Shepherdspurse	Wild oats
Curly dock	London rocket	Smartweed	Wild sweet potato
Dallisgrass	Lovegrass	Smooth catsear	Witchgrass
Dandelion	Mallow	Smooth pigweed	Yarrow
Dogfennel	Mayweed	Sowthistle	Yellow nutsedge
Evening primrose	Milkweed	Spotted spurge	Yellow oxalis
Fall panicum	Mouseear chickweed	Spurge	Yellow starthistle
False dandelion	Nimblewill	Sunflower	and many other weeds
Fennel			
* Not for use in California	_		

3. Where To Use

For General Weed Control and Spot Spraying

- Along and on sidewalks, gravel/brick walkways, paths, curbs, paved areas (private roads and streets), driveways, parking areas, recreational areas.
- · Along ornamental lawns and turfgrass.
- Around or near ornamental/shade trees, fruit and nut trees, grapevines, shrubs, groundcovers, flower beds, mulched landscape beds.
- Along fences, hedgerows, under fencelines.
- Around rocks, signposts, light posts, or on fringe areas.
- Around or along buildings/structures, foundations, patios, houses, mobile homes, vacant lots, urban areas, domestic outdoor premises, and other areas associated with household or home life.
- In areas for lawn replacement or garden plot preparation.
- Urban areas where total vegetation control is desired.
- Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fencelines, around shelterbelts and other areas where nonselective weed control is desired.

For Landscaping

- Prepare areas for planting of ornamentals, trees, shrubs, desert landscapes, rock gardens, flower beds or similar plantings.
- Delay mowing, tilling, or disturbing the treated vegetation for at least seven (7) days after application of this product for the best results. However, if the treated vegetation is not disturbed, planting or transplanting of trees, flowers, shrubs, bedding plants may be done at one (1) day after application.

For Wildlife Areas

- Prepare areas for planting wildlife food.
- Control undesirable and exotic plants to allow for recovery or planting of native species.
- Make treatments to remove unwanted plants. Take care to keep spray off desirable plants.

For Brush Control Foliar Sprays for Brush

- Cut back brush that is taller than 5 feet and then spray regrowth.
- For maximum activity, weeds should be growing vigorously at the time of the postemergent applications. The most favorable period for treatment occurs after full leaf stage in the spring and continues into early summer. Brush treated in the late fall may not be fully controlled.
- Spray uniformly the foliage of the vegetation to be controlled. Apply for full coverage and thorough brush/weed contact.
- Follow-up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to-control species including blackberry, poison oak, and kudzu. Another choice for effective brush control is Gordon's Cleanout® Brush and Stump Spray. Gordon's Cleanout® Brush and Stump Spray is recommended to control undesirable woody plants including buckbrush, poison ivy, multiflora rose, and sumac. Foliar applications of Gordon's Cleanout® Brush and Stump Spray are appropriate during the period of active growth of the brush. Cut stump, basal bark, and frill treatments of Gordon's Cleanout® Brush and Stump Spray may be applied any time of year.

Cut-Surface Stump Treatment (Not for use in California)

- Cut stump treatments are effective on stumps with diameters larger than 3-4 inches in diameter.
- Cut trees and brush close to the soil surface. Apply this to the freshly cut surface of the stump *immediately* after cutting. Delays in application may reduce the effectiveness.
- Apply for full and thorough coverage of the cut surface. Treat the entire circumference of the stump by drenching the cambium (the outermost wood between the pith and the bark).
- Treatment will control or suppress many types of woody brush and trees, such as alder, bluegum eucalyptus, madrone, oak, giant reed, saltcedar, sweetgum and tan oak.
- The most favorable period for cut stump treatment occurs after full leaf stage in the spring and continues into early summer. Brush and trees treated in the late fall may not be fully controlled.
- Follow-up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to-control species. Another choice for effective control with a cut-surface stump treatment is Gordon's Cleanout® Brush and Stump Spray.

For Lawn Renovation and Lawn Replacement

- Spot treatments or broadcast applications can be used to establish or renovate warm season and cool season turfgrass from seed or planting stock. This product may be applied before planting, seeding, sprigging, plugging, or sodding.
- Use EH-1391 Herbicide to remove the existing lawn and kill weeds and grasses. Uniform and complete spray coverage of all existing vegetation is required.
- Bermudagrass, nutsedge, torpedograss and deep rooted perennials may require a second treatment in 5 to 7 days.
- For best results, apply in spring or fall when daytime temperatures are at least 60° F.
- Best results are obtained when one mowing is skipped before and after the application of this product.

- If the lawn is dry, water lawn before applying EH-1391 Herbicide. Water again 2 to 3 days after application.
- Seeding, sprigging, plugging or sodding may be done at seven (7) days after application of this product. Prepare the soil for planting by raking, rototilling, vertical cutting, or other tillage method.
- Rake up and remove loosened thatch or debris.
- Mix in starter fertilizer and soil amendments if needed. Establish the lawn according to local cultural practices.

For Vine Control

- If vines are growing up poles, fences, or tree trunks with mature bark, cut vines to a height of 3 to 4 feet and spray vines thoroughly.
- If vines are climbing shrubs or tree trunks with green bark, cut vines at base and treat as directed for stumps or spray regrowth. If spraying regrowth, shield shrubs and green bark from spray drift with a piece of cardboard or plastic.

4. How And When To Apply

- EH-1391 Herbicide is formulated for "spot" treatment of individual weeds and grasses. This product is ready-to-use (RTU) and is applied directly from the container fitted with a trigger sprayer. Adjust the sprayer tip to deliver coarse spray droplets to the individual weeds. Direct the spray to the target weeds—the ones you want to kill—to provide uniform and thorough coverage. Apply on a spray-to-wet basis.
- Trigger sprayers that deliver droplets that foam on contact: EH-1391 is formulated for "spot" treatment of individual weeds and grasses. This product is ready-to-use (RTU) and is applied directly from the container fitted with a trigger sprayer. This nozzle delivers spray droplets (stream) that foams on contact with the plant leaves. This trigger sprayer improves the accuracy of spray placement and reduces the dripping on desirable plants. Direct the spray to the target weeds (or to individual weeds) and apply on a spray-to-wet basis. Allows you to see treated weeds. Ideal for tight spots like flower beds and gardens.
- [Supplemental registrants may include or omit instructions for use as an edger as presented in this paragraph.] EH-1391 Herbicide is an easy edger for use around desirable plants and vegetation. To use, remove cap and fasten the built-in sprinkler applicator. For edging, turn container upside down. Walk slowly holding nozzle 2 to 6 inches above [weeds and grasses][area to be treated]. Thoroughly wet the foliage of the unwanted vegetation. A one-gallon container covers a 2 inch band 1,200 feet long. For larger area coverage, adjust the width of the spray pattern by varying the height of the container.]
- Battery-operated trigger sprayer: This product is applied directly from the container using the
 battery-operated trigger sprayer. The sprayer is packaged with the battery installed. To use, remove
 the cap from the container, insert the pickup tube and thread cap attached to pickup tube to the
 container. Move the safety lock located below the trigger to the "ON" position. Pull trigger, waiting
 several seconds for fluid to prime. Adjust the sprayer tip to deliver coarse spray droplets. To store,
 move the safety lock located below the trigger to the "LOCK" position.
- Apply when the weeds are growing actively and air temperatures are above 60°F.
- Spray [Use] when air is calm to prevent drift to desirable plants.
- When treating around desirable plants, shield plants from drift with a sheet of cardboard or plastic. If desirable plant is accidentally sprayed, rinse off immediately with water.
- One (1) application will kill most of the weeds listed.
- Weeds usually begin to wilt within hours with complete kill in 1 to 2 weeks
- Rain or watering 2 hours after application will not wash away effectiveness
- Weed control may be reduced under adverse environmental conditions, e.g., drought, cool
 temperatures or misapplication. Difficult to control weeds, dense infestations/populations, or
 heavy/dense growth (canopy) of the target species may require additional applications. Two (2) or
 more applications at four (4) weeks intervals may be required for season long control.
- Ornamental flowers, trees and shrubs may be planted 1 day after application.
- This 32 fl.oz. (1.0 quart) retail container size treats approximately 250 weeds or up to 875 sq.ft.

- This 64 fl.oz. (1/2 gallon) retail container size treats approximately 500 weeds or up to 1,750 sq.ft.
- This 128 fl.oz. (1.0 gallon) retail container size treats approximately 1,000 weeds or up to 3,500 sq.ft.

[OPTIONAL TEXT:]

GARDENS (vegetables, herbs, small fruits and berries):

This product may be applied before planting or transplanting of vegetables (garden crops), herbs and small fruits and berries. This product may be used as a broad spectrum postemergent herbicide prior to planting to control annual and perennial weeds including field bindweed, johnsongrass, or dallisgrass. Spray the undesirable vegetation on a spray-to-wet basis. Uniform and complete spray coverage of all existing vegetation is required.

Planting intervals:

Blackberry, blueberry, boysenberry, currant, dewberry, elderberry, gooseberry, huckleberry, Loganberry, olallieberry, red raspberry and black raspberry	1 day
Artichoke (Jerusalem), beans (all), beet greens, broccoli, Brussels sprout, cabbage (all), carrot, cauliflower, celery, chard (Swiss), chicory, collards, endive, horseradish, kale, kohlrabi, leek, lentil, lettuce, mustard greens, okra, onion, rape greens, parsley, parsnip, potato, radish, rhubarb, rutabaga, spinach, sweet potato, turnip and yams	1 day
Corn, cucumber, eggplant, garlic, gourds, melons (all), pea (all), pepper (all), pumpkin, squash (summer and winter), tomatillo, tomato (by seed) and watercress	3 days
Tomatoes from transplants and all berries, fruits, vegetables and herbs not on previous lists	30 days

Harvest Intervals:

Allow a minimum of 14 days between the last application of this product and harvest of berries: Blackberry, blueberry, boysenberry, currant, dewberry, elderberry, gooseberry, huckleberry, Loganberry, olallieberry, red raspberry and black raspberry. [END OPTIONAL TEXT]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Pronto® is a registered trademark of PBI/GORDON Corporation

SUBLABEL 2

EH-1391 HERBICIDE

EPA Reg. No. 2217-844

Alternate Brand Names: GroundWork Ready-To-Use 2% Weed & Grass Killer, Gordon's Pronto Fast Acting Weed & Grass Killer, Gordon's Pronto Fast Acting Weed & Grass Killer - Foams on Contact

ACTIVE INGREDIENT:

Isopropylamine salt of glyphosate*	2.0%
OTHER INGREDIENTS:	<u>98.0%</u>
TOTAL	100.00%

^{*}N-(phosphonomethyl)glycine

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Wash thoroughly with soap and water.

First Aid	
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
•	ct container or label with you when calling a poison control center or doctor or going for additional information in case of emergency, call toll free 1-877-800-5556.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater and rinsate.

[Environmental Hazards (alternate per PRN 2008-1)

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.]

Physical Or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Keep people and pets off treated areas until spray solution has dried.

EH-1391 Herbicide is intended to be applied by occupational users, non-commercial users, and landscape personnel for use in institutional, ornamental, and residential/domestic sites. Institutional sites are defined as areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses, and office buildings. Ornamental sites include area around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings. Finally, residential/domestic sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals or convalescent homes.

Facts and Tips

- EH-1391Herbicide is a ready-to use herbicide intended to control a broad spectrum of emerged annual and perennial weed species. EH-1391 Herbicide will control unwanted species of grass, sedge, broadleaf weeds, and certain woody species around trees, shrubs, fences.
- This product kills plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves down to the roots.
- This product is nonselective and will kill or injure all vegetation contacted by the spray. Do not spray plants or grasses you like—they may die too. Not recommended for spot weed control in lawns since this product kills lawn grasses.
- This product has no soil residual activity and thus will not affect nearby untreated plants.
- Plants that have not emerged at the time of application will not be controlled.
- Best results will be obtained when the weed species are actively growing. Apply when the weeds are growing actively and air temperatures are above 60°F.

1. Tough Weed Control (Annuals, Perennials, and Brush) CONTROLS WEEDS AND GRASSES INCLUDING:

Lawn and Turfgrass Spec EH-1391 Herbicide will prov		pression) of these lawn and t	urfgrass species:
Bahiagrass Bentgrass, colonial, creeping and redtop Bermudagrass, common and hybrid Bluegrass, annual, rough, and Kentucky	Bromegrass, downy and smooth Buffalograss* Canarygrass, reed Carpetgrass* Centipedegrass	Fescue, chewings, hard, red, tall and sheep Foxtail, green, yellow, creeping, meadow and giant Kikuyugrass	Orchardgrass Paspalum spp Ryegrass, annual (Italian), perennial, and hardy St. Augustinegrass Zoysiagrass or Japanese lawngrass
* Not for use in California			

Alder	Elderberry	Mock orange*-	Quaking aspen
Blackberry	Honeysuckie	Manzanita	Raspberry
Barberry*	Hickory	Pine	Sumac
Ceanothus spp.	Kudzu	Poison ivy	Trumpetcreeper
Cherry	Lantana	Poison oak	Wild blackberry
Coyotebrush	Maple	Poison hemlock	Western red cedar*
Dewberry	Multiflora rose	Oak	Willow
Elm			[

Annual and Perennial W EH-1391 Herbicide will pro		uppression) of these annual and	l perennial weed species:
Artichoke, Jerusalem	Fiddleneck	Oleander*	Tansy ragwort
Barnyardgrass	Field bindweed	Oxalis	Tansy mustard
Beggarweed	Field pennycress	Pampasgrass	Texas panicum
Bittercress	Field sandbur	Pennsylvania smartweed	Thistle
Black medic	Filaree	Pennywort	Timothy
Broadleaf plantain	Florida pusley	Pigweed	Toadflax
Buckhorn plantain	Garden spurge	Plantain	Torpedograss
Bull thistle	Goosegrass	Prickly lettuce	Trailing lantana
Burclover	Green foxtail	Primrose	Tumble mustard
Canada thistle	Guineagrass	Prostrate spurge	Vaseygrass
Chickweed	Hemp sesbania	Puncturevine	Velvetleaf
Clover	Henbit	Purple cudweed	Virginia-creeper
Cocklebur	Horsenettle	Purple nutsedge	White clover
Common groundsel	Horseradish	Purslane	Whitetop
Common mullien	Horseweed/marestail	Quackgrass	Wild barley
Common ragweed	Ironweed	Ragweed	Wild carrot
Coral-bead	Johnsongrass	Red clover	Wild geranium
Crabgrass	Knapweed	Red sorrel*	Wild morningglory
Creeping beggarweed	Knawel	Redroot pigweed	Wild mustard
Creeping Charlie	Knotweed	Shepherdspurse	Wild oats
Curly dock	Lambsquarters	Smartweed	Wild sweet potato
Dallisgrass	London rocket	Smooth catsear	Witchgrass
Dandelion	Lovegrass	Smooth pigweed	Yarrow
Dogfennel	Mallow	Sowthistle	Yellow nutsedge
Evening primrose	Mayweed	Spotted spurge	Yellow oxalis
Fall panicum	Milkweed	Spurge	Yellow starthistle
False dandelion	Mouseear chickweed	Sunflower	and many other weeds
Fennel	Nimblewill		
*Not for use in California			

2. Where To Use

- For General Weed Control and Spot Spraying
- Along and on sidewalks, gravel/brick walkways, paths, curbs, paved areas (private roads and streets), driveways, parking areas, recreational areas.
- Along ornamental lawns and turfgrass.
- Around or near ornamental/shade trees, fruit and nut trees, shrubs, groundcovers, flower beds, mulched landscape beds.
- · Along fences, hedgerows, under fencelines.
- Around rocks, signposts, light posts, or on fringe areas.
- Around or along buildings/structures, foundations, patios, houses, mobile homes, vacant lots, urban areas, domestic outdoor premises, and other areas associated with household or home life.
- In areas for lawn replacement or lawn renovation.

- Urban areas where total vegetation control is desired.
- Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fencelines, around shelterbelts and other areas where nonselective weed control is desired.
- Weed control may be reduced under adverse environmental conditions, e.g., drought, cool temperatures, or misapplication. Difficult to control weeds, dense infestations/populations, or heavy/dense growth (canopy) of the target species may require additional applications. Two (2) or more applications at four (4) weeks intervals may by required for season long control.

3. How To Apply

Trigger sprayers, compressed air sprayers, and small power sprayers provide excellent coverage for spot treatments of annual and perennial weeds.

A. Trigger sprayers:

- EH 1391 Herbicide is formulated for "spot" treatment of individual weeds and grasses. This product is ready-to-use (RTU) and is applied directly from the container fitted with a trigger sprayer. Adjust the sprayer tip to deliver coarse spray droplets to the individual weeds. Direct the spray to the target weeds—the ones you want to kill—to provide uniform and thorough coverage. Apply on a spray-to-wet basis. Additional applications may be necessary to control perennials.
- **B. Compressed air sprayers and small power sprayers** including but not limited to hand operated, compressed air sprayers with tank of 1 to 3 gallons, backpack sprayers or knapsack sprayers, and small power (engine driven) sprayers fitted with low-pressure wands.
 - EH-1391 Herbicide is formulated as a ready-to-use (RTU) herbicide. No dilution with water is necessary. Pressurize the sprayer and adjust the nozzle or wand to deliver coarse droplets in a narrow or wide spray pattern. Direct the spray to the leaves and stems of target weeds (or to individual weeds) and apply on a spray-to-wet basis. Additional applications may be necessary to control perennials.
 - This product is nonselective and will injure or kill all plants contacted with spray droplets. Avoid contact with leaves and stems of desirable plants.

For Landscaping

- Prepare areas for planting of ornamentals, trees, shrubs, desert landscapes, rock gardens, flower beds or similar plantings.
- Delay mowing, tilling, or disturbing the treated vegetation for at least seven (7) days after application of this product for the best results. However, if the treated vegetation is not disturbed, planting or transplanting of trees, flowers, shrubs, bedding plants may be done at one (1) day after application.

Table 1: Use rates of site preparation of landscape plantings for control of emerged weeds.		
	Amount of Ready-To-Use Product, Fluid Ounces / 1,000 sq.ft.	Amount of Ready-To-Use Product, Gallons per Acre (A)
Annual Weeds	18 to 36 fluid ounces / 1,000 sq.ft.	6 to 12 gallons of product /A
Perennial Weeds	36 to 86 fluid ounces / 1,000 sq.ft.	12 to 30 gallons of product /A

For Brush Control Foliar Sprays for Brush:

- Cut back brush that is taller than 5 feet and then spray regrowth.
- For maximum activity, weeds should be growing vigorously at the time of the postemergent applications. The most favorable period for treatment occurs after full leaf stage in the spring and continues into early summer. Brush treated in the late fall may not be fully controlled.
- Apply this product with hand operated equipment. Spray uniformly the foliage of the vegetation to be controlled. Apply for full coverage and thorough brush/weed contact.
- Follow-up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to-control species including blackberry, poison oak, and

kudzu. Another choice for effective brush control is Gordon's Cleanout® Brush and Stump Spray. Gordon's Cleanout® Brush and Stump Spray is recommended to control undesirable woody plants including buckbrush, poison ivy, multiflora rose, and sumac. Foliar applications of Gordon's Cleanout® Brush and Stump Spray are appropriate during the period of active growth of the brush. Cut stump, basal bark, and frill treatments of Gordon's Cleanout® Brush and Stump Spray may be applied any time of year.

Cut-Surface Stump Treatment (Not for use in California)

- Cut stump treatments are effective on stumps with diameters larger than 3-4 inches in diameter.
- Cut trees and brush close to the soil surface. Apply this to the freshly cut surface of the stump *immediately* after cutting. Delays in application may reduce the effectiveness.
- Apply this product undiluted using suitable equipment to provide complete coverage of the cambium.
 Apply for full and thorough coverage of the cut surface. Treat the entire circumference of the stump by drenching the cambium (the outermost wood between the pith and the bark).
- Treatment will control or suppress many types of woody brush and trees, such as alder, bluegum eucalyptus, madrone, oak, giant reed, saltcedar, sweetgum and tan oak.
- The most favorable period for cut stump treatment occurs after full leaf stage in the spring and continues into early summer. Brush and trees treated in the late fall may not be fully controlled.
- Follow-up applications of this product or with a different herbicide may be needed under less favorable growing conditions and for hard-to-control species. Another choice for effective control with a cut-surface stump treatment is Gordon's Cleanout® Brush and Stump Spray.

For Lawn Renovation and Lawn Replacement

- Spot treatments or broadcast applications can be used to establish or renovate warm season and cool season turfgrass from seed or planting stock. This product may be applied before planting, seeding, sprigging, plugging, or sodding.
- Use EH-1391 Herbicide to remove the existing lawn and kill weeds and grasses. Uniform and complete spray coverage of all existing vegetation is required.
- Bermudagrass, nutsedge, torpedograss and deep rooted perennials may require a second treatment in 5 to 7days.


For best results, apply in spring or fall when daytime temperatures are at least 60° F.

	r site preparation of lawn and turfgrass seed ontrol emerged weeds.	dbeds (prior to establishment) and for
	Amount of Ready-To-Use Product, Fluid Ounces / 1,000 sq.ft.	Amount of Ready-To-Use Product, Gallons per Acre (A)
Annual Weeds	18 to 36 fluid ounces / 1,000 sq.ft.	6 to 12 gallons of product /A
Perennial Weeds	36 to 86 fluid ounces / 1,000 sq.ft.	12 to 30 gallons of product /A

- Seeding, sprigging, plugging or sodding may be done at seven (7) days after application of this product. Prepare the soil for planting by raking, rototilling, vertical cutting, or other tillage method.
- Rake up and remove loosened thatch or debris.
- Mix in starter fertilizer and soil amendments if needed. Establish the lawn according to local cultural practices.

For Vine Control

- If vines are growing up poles, fences, or tree trunks with mature bark, cut vines to a height of 3 to 4 feet and spray vines thoroughly.
- If vines are climbing shrubs or tree trunks with green bark, cut vines at base and treat as directed for stumps or spray regrowth. If spraying regrowth, shield shrubs and green bark from spray drift with a piece of cardboard or plastic. Apply this product undiluted with hand operated equipment. Thorough coverage of the foliage is necessary for the best results.



STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

PESTICIDE STORAGE: STORE ABOVE 10° F (12°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room (68°F or 20°C) for several days to allow crystals to redissolve, then shake well before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:] CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. [Triple rinse container (or equivalent) promptly after emptying.]

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. [Triple rinse container (or equivalent) promptly after emptying.]

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill

the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[OR .

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

[For Refillable Containers:]

CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Pronto® is a registered trademark of PBI/GORDON Corporation

SUBLABEL 3

EH-1391 HERBICIDE

EPA Reg. No. 2217-844

Alternate Brand Names: GroundWork Ready-To-Use 2% Weed & Grass Killer, Gordon's Pronto Fast Acting Weed & Grass Killer, Gordon's Pronto Fast Acting Weed & Grass Killer - Foams on Contact

ACTIVE INGREDIENT:

· Isopropylamine salt of glyphosate*	2.0%
OTHER INGREDIENTS:	<u>98.0%</u>
TOTAL	100.00%

^{*}N-(phosphonomethyl)glycine

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Harmful if swallowed. Wash thoroughly with soap and water.

First Aid	·
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
	ct container or label with you when calling a poison control center or doctor or going for additional information in case of emergency, call toll free 1-877-800-5556.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater and rinsate.

[Environmental Hazards (alternate per PRN 2008-1)

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.]

Physical Or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Keep children and pets out of treated areas until sprays have dried.

1. How It Works

Facts and Tips

- EH-1391 Herbicide is a general purpose herbicide designed as a ready-to-use spray to control a
 broad spectrum of emerged annual and perennial weed species. EH-1391 Herbicide will control
 unwanted species of grass, sedge, broadleaf weeds, and certain woody species around trees,
 shrubs, fences.
- This product kills plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves down to the roots.
- This product is nonselective and will kill or injure all vegetation contacted by the spray. Do not spray plants or grasses you like—they may die too. *Not recommended for spot weed control in lawns since this product kills lawn grasses.*
- Any product not absorbed by plants breaks down into natural materials without moving in or on the soil to untreated plants.
- This product has no soil residual activity and thus will not affect nearby untreated plants.
- Plants that have not emerged at the time of application will not be controlled.
- Best results will be obtained when the weed species are actively growing.

2. Tough Weed Control

EH-1391 Herbicide will provide full or partial control (suppression) for these annual and perennial weed species:

Lawn and Turfgrass such as Bermudagrass, Bluegrass, Fescue, Zoysiagrass and other grasses.

Woody Species such as Poison ivy, Poison oak, Sumac, Trumpetcreeper and others.

Annual and Perennial Weed Species such as Goosegrass, Knotweed, Spurge, Torpedograss, Yellow nutsedge and many other weeds.

Restrictions for weed species in California (Not for use in California): Buffalograss, Carpetgrass, Barberry, Mock orange, Western red cedar, Oleander and Red sorrel.

3. Where To Use

For General Weed Control and Spot Spraying

- Along and on sidewalks, gravel/brick walkways, paths, curbs, paved areas (private roads and streets), driveways, parking areas, recreational areas.
- Along ornamental lawns and turfgrass.
- Around or near ornamental/shade trees, fruit and nut trees, grapevines, shrubs, groundcovers, flower beds, mulched landscape beds.
- Along fences, hedgerows, under fencelines.
- Around rocks, signposts, light posts, or on fringe areas.
- Around or along buildings/structures, foundations, patios, houses, mobile homes, vacant lots, urban areas, domestic outdoor premises, and other areas associated with household or home life.
- Urban areas where total vegetation control is desired.
- Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fencelines, around shelterbelts and other areas where nonselective weed control is desired.

4. How And When To Apply

- EH-1391 Herbicide is formulated for "spot" treatment of individual weeds and grasses. This product is ready-to-use (RTU) and is applied directly from the container fitted with a trigger sprayer. Adjust the sprayer tip to deliver coarse spray droplets to the individual weeds. Direct the spray to the target weeds—the ones you want to kill—to provide uniform and thorough coverage. Apply on a spray-to-wet basis.
- Apply when the weeds are growing actively and air temperatures are above 60°F.
- Spray when air is calm to prevent drift to desirable plants.
- When treating around desirable plants, shield plants from drift with a sheet of cardboard or plastic. If desirable plant is accidentally sprayed, rinse off immediately with water.
- One (1) application will kill most of the weeds listed.
- Weeds usually begin to wilt within hours with complete kill in 1 to 2 weeks
- Rain or watering 2 hours after application will not wash away effectiveness
- Weed control may be reduced under adverse environmental conditions, e.g., drought, cool
 temperatures or misapplication. Difficult to control weeds, dense infestations/populations, or
 heavy/dense growth (canopy) of the target species may require additional applications. Two (2) or
 more applications at four (4) weeks intervals may be required for season-long control.
- This 32 fl.oz. (1.0 quart) retail container size treats approximately 250 weeds or up to 875 sq.ft.
- This 64 fl. oz. (1/2 gallon) retail container size treats approximately 500 weeds or up to 1,750 sg. ft.
- This 128 fl. oz. (1/2 gallon) retail container size treats approximately 500 weeds or up to 3,500 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Pronto® is a registered trademark of PBI/GORDON Corporation

APPENDIX FOR ALL SUBLABELS

1. Statements which may appear on different label components depending on packaging
configuration.
See next panel for additional Precautionary Statements and First Aid
Net Contents:
• EPA Est. No
2. Advertising claims that may be presented on container labeling, advertisements, brochures,
and other marketing/sales promotional materials:
Ready-To-Use no mixing
For nonselective postemergent control of unwanted vegetation
For nonselective weed control on noncrop/residential areas
Contains glyphosate
Quick and easy to use
No leaks or mess
Easy to store
No mixing required
 Covers approximately # sq. ft. [dependent upon container size]
 Systemic weed and grass killer for spot treatment of unwanted weeds and grasses
Kills weeds and grasses listed on this label
Kills to the root
No soil residual activity
 Kills roots and all in 2 to 4 weeks
Kills even the toughest weeds
Eliminates undesirable vegetation in 2 to 4 weeks
• For use around flower beds, trees and shrubs, fences, walks, mulch beds, gravel and driveways
For use around flower beds, trees, shrubs, fences and walks
Ideal for vegetable garden weed control
• [Rain-Proof in hours][Rain-Proof in less than hours].
• Results in as little as 24 hours

- [For product information and a complete list of weeds controlled] [For more information] [For questions or comments], [call toll-free 1 8xx-yyy-zzzz or visit (website address)]
- · Fast acting
- · Rainproof in 2 hours
- [Even] Kills the root[s]
- Use in gardens, patios, driveways and walks
- The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer, and Buyer assumes the risk of any such use.
- Results in 24 Hours
- Glyphosate 2%
- With Battery-Operated Sprayer
- Rociador de Potencia para Hierba y Mala Hierba [Power Sprayer for Grass and Weeds]
- Treated lawns [areas] can be re-seeded 7 days after application
- Satisfaction Guaranteed: If for any reason you are not completely satisfied with the performance or results of this GreenThumb® product, it will be replaced free of charge. Simply provide proof of purchase and return the product and/or unused portion to the place of purchase.
- · Rain fast in hours-
- · Rain-fast in two hours
- For outdoor use
- · For spot treatment of individual weeds
- · Allows grass reseeding in one week
- Controls Weeds
- Kills Roots
- Covers [up to] XXX sq.ft.

Advertising claims for Trigger Sprayers that deliver droplets that foam on contact:

- Foams on contact see where you sprayed
- This sprayer can improve the accuracy of spray placement
- Combine the effectiveness of glyphosate for target weeds with the application safety for desirable (nontarget) plants.
- · Kills weeds without dripping onto desirable plants.
- Foams on application.
- · Ideal for target weeds in tight spots.

DOCUMENT CONTROL INFORMATION

- 1. Unique Label Identifier: 002217-00844.20130418.notif-proposed-highlighted.doc
- 2. Reason for Issue: Sublabel 2 storage and disposal