

2217-853

4-9-2002

47

GORDON'S[®] PERMETHRIN 10 LIVESTOCK AND PREMISE SPRAY

For Use On Livestock: Cattle (Beef and Dairy), Goats, Sheep, Swine, Horses and Poultry.

- ◆ *For Agricultural/Commercial Use—Not for Household Use*
- ◆ *A Broad Spectrum Multi-Use Insecticide*

ACTIVE INGREDIENT:

Permethrin..... 10.0%

INERT INGREDIENTS:..... 90.0%

TOTAL 100.0%

Cis/trans ratio: Max. 65% (+/-) trans and min. 35% (+/-) cis.
Contains Petroleum Distillates

ACCEPTED
APR 9 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2217-853

KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panel for First Aid and Additional Precautionary Statements.

NET CONTENTS: 1 PINT, 1 QUART, or 1 GALLON

APXXXXXX
EPA REG. NO. 2217-ILG
EPA EST. NO. 2217-KS-1
MANUFACTURED BY:



2/7



READ THE ENTIRE LABEL BEFORE EACH USE. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

FOR PET USE - USE ONLY ON DOGS

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: Harmful if inhaled. Causes moderate eye injury and skin irritation. Do not get on skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing vapor or spray mist and contact with eyes. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Wear dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any R, P or HE filter for applying indoors as a space spray or fog.

FIRST AID	
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.</p>	
Note to Physician and Veterinarians:	<ul style="list-style-type: none"> • Contains petroleum distillate - vomiting may cause aspiration pneumonia. • Do not administer or instill milk, cream or other substances containing vegetable or animal fats which enhance absorption of lipophilic substances.

3/7

ENVIRONMENTAL HAZARDS:

This product is highly toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the high water mark. Do not apply where runoff is likely to occur. Do not apply when wind speeds exceed 10 mph. Do not contaminate water when disposing of equipment washwaters. Shrimp and crab may be killed if application rates recommended on this label are exceeded. Use with caution where these are important resources.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply this product or allow drift when bees are actively visiting treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS:

Combustible. Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shake or stir product thoroughly before using or reusing diluted emulsion. To prepare dilutions, the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water and blend thoroughly. Do not hold dilutions for more than 24 hours.

Do not allow spray treatment to contaminate pasture land, crop land, poultry ranges or water supplies. Do not use on crops used for food, forage or pasture.

APPLICATION SITES	PESTS CONTROLLED	USE DIRECTIONS	DILUTION RATE	APPLICATION RATE
ANIMAL PREMISE USE: SURFACE SPRAY AND SPACE SPRAY FOR CATTLE BARNs, SWINE HOUSES, POULTRY HOUSES, HORSE BARNs, DAIRIES, MILKING PARLORS				
AS SURFACE SPRAY: FARM PREMISES (Barns, Dairies, Loafing Sheds, Milking Parlors, Feed Lots, Stables, Paddocks, Poultry and Livestock Housing.	Horse Flies, Stable Flies, Lesser House Flies and Other Manure Breeding Flies; Fleas; Lice; Mites; Ticks, Including Deer Tick. Aids In Control of Cockroaches, Mosquitoes, Spiders.	Observe (follow) application intervals greater than 2 weeks. Do not contaminate animal's feed or water by spray. Do not use in milk rooms. For maximum effectiveness, a combination of localized application and space treatment is recommended.	For initial clean up of severe insect infestation, dilute at a rate of 1 part concentrate in 19 parts water: <u>Amt. Product per Water Volume</u> 6.7 fl.ozs. 1.0 gallon 8 fl.ozs. 1.2 gallons 16 fl.ozs. (1pt.) 2.4 gallons 32 fl.ozs. (1qt.) 4.0 gallons For normal infestations, dilute 1 part concentrate in 100 parts water. <u>Amt. Product per Water Volume</u> 1.33 fl.ozs. 1.0 gallon 8 fl.ozs. 6.25 gallons 16 fl.ozs. (1pt.) 12.5 gallons 32 fl.ozs. (1qt.) 25.0 gallons	Spray walls and surfaces thoroughly but do not allow runoff to occur; about 1 gallon per 750 - 1,000 sq.ft.
AS A SPACE SPRAY: Cattle Barns, Horse Barns, Swine Houses, Kennels, Milking Parlors, Milk Rooms, Dairies, Poultry Houses, Feed Lots, Stables, and Paddocks.	Flies, Mosquitoes, Gnats.	FOG OR FINE MIST: Direct the spray toward the ceiling and upper corners until the area is filled with mist. Use about 2 fl.oz. per 1,000 sq.ft. of space For best results, close doors and windows before spraying and keep closed for 10-15 minutes. Vacate the treated area and ventilate before reoccupying. Observe (follow) application intervals greater than 2 weeks. For maximum effectiveness, a combination of localized application and space treatment is recommended.	<u>Amt. Product per Water Volume</u> 1.33 fl.ozs. 1.0 gallon 8 fl.ozs. 6.25 gallons 16 fl.ozs. (1pt.) 12.5 gallons 32 fl.ozs. (1qt.) 25.0 gallons	Direct the spray toward the ceiling and upper corners until the area is filled with mist. Use about 2 fl.oz. per 1,000 sq.ft. of space.

APPLICATION SITES	PESTS CONTROLLED	USE DIRECTIONS	DILUTION RATE	APPLICATION RATE
LIVESTOCK AND POULTRY USE:				
DIRECT APPLICATIONS ON CATTLE (BEEF AND DAIRY), GOATS, SHEEP, SWINE, HORSES, AND POULTRY				
LIVESTOCK SPRAY: Dairy or Beef Cattle, Horses, Sheep, and Goats.	Face Flies, Horn Flies, Stable Flies, Mosquitoes, Lice, Mites, Ticks, Including Deer Tick.	Spray to thoroughly cover entire animal. For lice or mites, a second treatment is recommended 14-21 days later. Spray lactating dairy cows only after milking is completed.	<u>Amt. Product per Water Volume</u> 8 fl.ozs. 50 gallons 16 fl.ozs. (1pt.) 100 gallons 32 fl.ozs. (1qt.) 200 gallons	
	Horn Flies, Face Flies, Stable Flies, Ear Ticks.	SPOT TREATMENT - LOW PRESSURE SPRAY: Apply ½ fl.oz. per ear or 2-4 fl.oz. on face or 12-16 fl.oz. along backline.	<u>Amt. Product per Water Volume</u> 2 fl.ozs. 1.0 gallons 16 fl.ozs. (1pt.) 2.5 gallons 32 fl.ozs. (1qt.) 5.0 gallons	
	Face Flies, Horn Flies, Horse Flies, Ear Ticks.	BACK RUBBER: Recharge backrubber as needed. Spray lactating dairy cows only after milking is completed.	<u>Amt. Product per Water Volume</u> 8 fl.ozs. 5 gallons 16 fl.ozs. (1pt.) 10 gallons 32 fl.ozs. (1qt.) 20 gallons	
Swine (Do not ship swine for slaughter within 5 days of last treatment.)	Mange Mites	Spray, dip or sponge animals. Retreat after 14 days, spraying walls and floor and space bedding to kill late hatching, developing stages.	<u>Amt. Product per Water Volume</u> 8 fl.ozs. 25 gallons 16 fl.ozs. (1pt.) 50 gallons 32 fl.ozs. (1qt.) 100 gallons	
	Blow Flies, Flies, Mosquitoes, Hog Lice, Fleas, Ticks, Including Deer Tick.	Spray, dip or sponge to apply 1 pint of spray solution per pig, especially around ears.	<u>Amt. Product per Water Volume</u> 8 fl.ozs. 25 gallons 16 fl.ozs. (1pt.) 50 gallons 32 fl.ozs. (1qt.) 100 gallons	
Poultry	Poultry Mites, Northern Fowl Mites, Lice	Spray at the rate of ½ fl.oz. of spray solution per bird, or 1 gallon of spray solution per 100 birds, with a fine mist. Spray roosts, walls and nests or cages.	<u>Amt. Product per Water Volume</u> 8 fl.ozs. 1.25 gallons 16 fl.ozs. (1pt.) 25 gallons 32 fl.ozs. (1qt.) 50 gallons	

5/7

APPLICATION SITES	PESTS CONTROLLED	USE DIRECTIONS	DILUTION RATE	APPLICATION RATE								
DOG PREMISES AND DIRECT APPLICATIONS TO DOGS.												
DOG PREMISES: Kennels, Dog Houses, Runs and Yards	Fleas, Flies, Ticks, Lice Mange (adult and larvae).	<p>SURFACE SPRAY: Thoroughly wet pest breeding or resting areas but do not allow runoff to occur. Most efficient pest control is accomplished when heavy pressure and good contact are achieved.</p> <p>Timing and frequency of applications should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray feed, food or water. Repeat as needed but not more often than once every 2 weeks.</p>	<table border="0"> <tr> <td colspan="2"><u>Amt. Product per Water Volume</u></td> </tr> <tr> <td>8 fl.ozs.</td> <td>6.25 gallons</td> </tr> <tr> <td>16 fl.ozs. (1pt.)</td> <td>12.5 gallons</td> </tr> <tr> <td>32 fl.ozs. (1qt.)</td> <td>25.0 gallons</td> </tr> </table>	<u>Amt. Product per Water Volume</u>		8 fl.ozs.	6.25 gallons	16 fl.ozs. (1pt.)	12.5 gallons	32 fl.ozs. (1qt.)	25.0 gallons	
<u>Amt. Product per Water Volume</u>												
8 fl.ozs.	6.25 gallons											
16 fl.ozs. (1pt.)	12.5 gallons											
32 fl.ozs. (1qt.)	25.0 gallons											
PESTS ON DOGS (Do not treat puppies less than 12 weeks old)	Fleas; Ticks, Including Deer Ticks; Lice; Mange	Thoroughly wet the animal by dipping, sponging or spraying. Allow animal to dry in a warm place without rinsing or toweling. Gives long-lasting protection against reinfestation and applications at intervals of 2-3 weeks. Do not repeat treatment for 7 days. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.	<table border="0"> <tr> <td colspan="2"><u>Amt. Product per Water Volume</u></td> </tr> <tr> <td>8 fl.ozs.</td> <td>12.5 gallons</td> </tr> <tr> <td>16 fl.ozs. (1pt.)</td> <td>25 gallons</td> </tr> <tr> <td>32 fl.ozs. (1qt.)</td> <td>50 gallons</td> </tr> </table>	<u>Amt. Product per Water Volume</u>		8 fl.ozs.	12.5 gallons	16 fl.ozs. (1pt.)	25 gallons	32 fl.ozs. (1qt.)	50 gallons	
<u>Amt. Product per Water Volume</u>												
8 fl.ozs.	12.5 gallons											
16 fl.ozs. (1pt.)	25 gallons											
32 fl.ozs. (1qt.)	50 gallons											

6/7

7/7

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: For Metal Containers - Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. For Plastic Containers - Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.