2217-855	7-21-2003/10	071141011/10	91 1	-		1/4		
Please read instructions on	reverse before completing form. United S Environmental Pro Washington,	otection Agency		Registrati	on	Approval Expires 05-31-98 OPP Identifier Number		
	Applica	tion for Pesticid	e - Section	I				
1. Company/Product Nun	nber 2217-855	2. EPA F	Product Manage Joanne Mi		3. Proj	posed Classification		
4. Company/Product (Nat 1382 EC	ne) C Residential_Herbicide	PM#	(PM 23))		None Restricted		
5. Name and Address of PBI/Gordon Corpora PO Box 014090 Kansas City, Missour	(b)(i), my EPA Re	 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name 						
		Section - II		<u></u>				
Amendment - Explain Resubmission in resp Notification - Explain	onse to Agency letter dated		Final printed label Agency letter date Me Too" Applicat Dther - Explain be	ion.	NOT	FICATION 2 1 2003		
Explanation: Use additional page(s) if necessary. (For section I and Section II.) <u>Notification of Additional Brand Name:</u> We ask to add an additional brand name of "Speed Zone St. Augustine Residential" with the "Gordon's ProForm Professional Formulations" logo. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C.Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.								
		Section - III						
1. Material This Product Will Child-Resistant Packaging Yes* No * Certification must be submitted			ckaging No. per container	2. Type of Con Metal Plastic Glass Paper	tainer	· · · ·		
3. Location of Net Contents In		Retail Container 3, 16, 32 & 64 fluid ounces 1, 30, 55 gallon	s 🗹 d	cation of Label I On Label On Labeling acco	Directions	product		
6. Manner in Which Label is Affixed to Product Paper glued Stenciled								
		Section - IV				<u></u>		
Name	ems directly below for identification g Martens	Title Manager of R	m*m*	Te	lephone N	ny o. <i>(incluce Area Code)</i> 16/460-6287		
I certify that the stateme I acknowledge that any I both under applicable la 2. Signature	Certifi nts I have made on this form and a knowingly false or misleading state w.	cation	true, accurate a by fine or impriso	nd complete. nment or	6.	Date Application Received (Stamped)		
4. Typed Name Craig	Martens	5. Date July 1	, 2003					

EPA Form 8570-5 (Rev. 2-92) Previous editions are obsolete.



Speed+Zone AI GI STINE RESIDENTIAI

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester														10.49
Mecoprop-p, acid											•	• •		2.66
Dicamba, acid														
Carfentrazone-ethyl														0.54
INERT INGREDIENTS:							•							85.64
									Ţ	Ö	T	A	Ľ	100.00

THIS PRODUCT CONTAINS:

0.52 lbs. 2.4-dichlorophenoxyacetic acid equivalent per gallon or 6.96%. 0.20 lb. (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 2.66%.

- 0.05 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.67%.
- 0.04 lb. Ethyl a,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1 2,4-triazoi-1-yi]-4-tluorobenzenepropanote or 0.54%

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN CAUTION

See next column for additional Precautionary Statements

First	Aid
FIISU	Alu

If swallowed:	Call a poison control center or doctor immediately for treatment advice.
	Do not induce vomiting unless told to by a poison control center or doctor.
1	 Do not give any liquid to the person.
	Do not give anything by mouth to an unconscious
ł	(Continued on next column)

First	Aid	(continued)
-------	-----	-------------

If on skin:	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia.



PRECAUTIONARY STATEMENTS

Hazarda to Humana and Domestic Animala

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin. eves, or clothing. Harmful if swallowed. When using this product, wear longsleeved shirt, long pants, socks, shoes, and chemical resistant gloves.

Personal Hygiene Statements:

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promotly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons; Persons engaged in open pouring of this product must also wear coveralis or a chemical resistant apron.

Containers of 5 gallons or more: Do not open-pour from this container. A mechanical system (such as probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

ENVIRONMENTAL HAZARDS:

Carlentrazone-ethyl is very toxic to algae and moderately toxic to fish. This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aguatic invertebrates and nontarget plants. Do not apply directly to water.

Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing, loading and disposal sites. Caution should be exercised when handling 2.4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

PRODUCT DESCRIPTION:

This Trimec[®] product is a combination of four proven weed killers: 2,4-D, mecoprop-p, dicamba and carlentrazone-ethyl, Postemergent applications of this product control a wide range of broadleaf lawn weeds in residential lawns including dandelion, chickweed, spurge, pennywort (dollarweed), knotweed, plantain, and clover.

- This product offers these advantages:
- · Rapid and effective weed control for common and troublesome weed species in turforass
- Superior cool weather performance.

 Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed within hours of the application and plant death can occur within 7-14 days.

WHERE TO USE:

This product is intended for use on most ornamental lawns including:

Cool Season Turfgrass	Warm Seeson Turfgrase			
Kentucky bluegrass	Common bermudagrass			
Perennial ryegrass	Hybrid bermudagrass			
Tall fescue	Bahiagrass			
Red or fine leaf lescues	Zoysiagrass			
Colonial bentgrass	Buffalograss			
Mixtures of cool season species	St. Augustinegrass			
	Centipedegrass			


Turforasa tolerance:

The turforass tolerance to Speed-Zone® St. Augustine Residential may vary and temporary turiorass vellowing may occur on St. Augustinegrass and certain varieties (F1) hybrids of hybrid bermudagrass.

Speed-Zone St. Augustine Residential may injure certain turforass species. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90°F.

Prohibitions:

- . Do not use this product on carpetorass, dichondra or lawns containing desirable clovers.
- . Do not apply this product to St. Augustinegrass during spring green-up which is the transition period between dormancy and active growth.
- · Do not spray roots of trees and ornamentals.
- . Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result.
- Do not exceed specified dosages for any area. Se particularly careful within dripline of trees and other ornamental species.



WHEN TO APPLY:"

Spray when broadleat weeds are young and actively growing for best results. Speed-Zone St. Augustine Residential may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2-6 week intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are susceptible.

CULTURAL TIPS FOR IMPROVED CONTROL:

Irrigation:

 Do not apply this product immediately before rainfall or irrigation. Rain or watering within 4 hours after treatment may reduce effectiveness.

Mowing:

Delay mowing 1 to 2 days before and after the application of this product.

Reseeding Interval

Treated areas may be reserved 2 weeks after application.

THREE EASY STEPS TO MIX AND APPLY:

- Determine (Measure) the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is presented as fl.oz. (tablespoons) per 1,000 sq. ft. of lawn.
- 2. Use hand-operated sprayers (backpack sprayers, compression sprayers, knapsack sprayers or tank type) or hose end sprayers. Carefully measure and mix the proper amount of product with water. See below for instructions for the various types of sprayers.
- 3. Spray evenly over the lawn.

HOW MUCH TO USE:

I. BROADCAST TREATMENTS FOR COOL SEASON TURFGRASS: Use rates of Speed-Zone St. Augustine Residential on cool season turfgrass as broadcast treatments are presented in Table 1.

TABLE 1. RATE RECOMMENDATIONS FOR COOL SEASON TUREGRASS.

[Amount of		Traded Area, sq. R.	
Species	Product 81. oz./1,000 są. It.	8.8 8. ez. sprayer jar er sulali containen	1.0 pf. (16 H. oz.) sprayer jar er rotail centainer	1.0 qt. (3211.oz.) Sprayer jar or rotail containor
Kentucky bluegrass	15-2.0	4,000-5,390 sq. ft	8,000-10,600 sq. ft	16 000-21,300 sq. ft
Perennial cyegrass	1.5-2.0	4,000-5,300 sq. ft	8,000-10,600 sq. M	16,000-21,300 sq. ft
Tall lescae	1.5-2.0	4,000-5,300 sq. R	8,000-10,600 sq. It	16,090-21,380 sq. ft
Red or fine leaf lescues	1.5-2.0	4,000-5,300 sq R	8,000-10,600 sq. R	16,000-21,300 sq. R
Colonial betsigrass	1.5-2.0	4,008-5,309 sq. ft	8,000-10,600 sq. R	16,000-21,300 sq. it
Mixtures of cool season species	1.5-2.0	4,000-5,300 sq. ft	8,000-10,600 sq. 8	16,000-21,300 sq. h

HOW TO APPLY:

- FOR ORTHO HOSE-END SPRAYERS FOR COOL SEASON TURFGRASS • For Ortho® Dial'N Spray Hose-End Sprayer: Measure 2.0 (I. oz. (4 Tbs.)
- into sprayer jar. Do not add water. Set dial on 1 oz. and spray uniformly on 1,000 sq. ft. of lawn.
- For Ortho® Plant & Weed Sprayette® Hose-End Sprayer: Measure 2.0 fl. oz. (4 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 sq. ft.
- For Gilmout[®] 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl. oz. and Dilution Rate of 80:1:
- For 1.5 ft oz. of product per 1,000 sd. ft.: Measure 10 ft. oz. of product into sprayer jar. Add water or fill with water to the 32 ft. oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq. ft. of lawn.
- For 2.0 II. oz. of product per 1.000 so. ft.: Measure 13.3 fl. oz. of product into sprayer jar. Add water or fill with water to the 32 fl. oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq. ft. of lawn.
- For Gilmour[®] Metering Dial Insecticide & Fertilizer Sprayer: This sprayer can be set to meter tablespoons per gallon of water or can be set to teaspoons per gallon of water by using the converting orifice in the top of the mixing head.
- For 1.5 fl. oz. of product per 1.000 sq. ft.; Turn the metering dial until the desired setting number is next to the setting arrows on the side of the mixing head. Set the dial at 1 tablespoon per gallon or 3 teaspoons per gallon. Add herbicide to the sprayer bottle. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. By squeezing the handle, you will be automatically spraying the proper spray concentration.
- For 2.0 fl. oz. of product per 1.000 sci. ft.. Turn the metering dial until the desired setting number is next to the setting arrows on the side of the mixing head. Set the dial at 1 tablespoon per gallon or 4 teaspoons per gallon. Add herbicide to the sprayer bottle. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. By squeezing the handle, you will be automatically spraying the proper spray concentration.

The maximum application rate to turf is 0.5 pound 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

II. SPOT TREATMENTS FOR COOL SEASON TURFGRASS.

HAND OPERATED SPRAYERS:

For cool season turtgrass, mix 1.5-2.0 fl. oz. of Speed-Zone St. Augustine Residential per 1.0 gallon of water for treatment of approximately 1,000 sq. ft. of turtgrass. Apply any time the emerged broadleal weeds are susceptible. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants.

Spray volume, gais. of water	Fluid ownces (II, oz.) of product	Tablespoone (ths.) of product	Area to be treated, sq. if.
1 gallon	1.5-2.0 fl. oz.	3.0- 4.0 tbs.	1,000 sq. ft.
2 gailons	3.0-4.0 fl. oz	6.0- 8.0 tbs.	2,000 sq. R.
3 gallons	4.5-6.0 fl. oz	9.0-12.0 tbs.	3,000 sq. ft.

Use rates of Speed-Zone St. Augustine Residential on cool season turigrass as broadcast treatments are presented in Table 2.

TABLE 2. RATE RECOMMENDATIONS FOR WARM SEASON TURFGRASS.

	Januari of	Treated Area, 29. N.						
Species	Preduct R. uz./1,000 sq. R.	0.0 II. ez. sprayer jar er retail container	f.0 pl. (16 0. oz.) sprayor jar or rotali custaioor	1.0 et. (32 fl. ez.) sprayer jar er selait container				
Common bernudagrass	1.0-2.0	4,000-8,000 sq. R	8 000-16 000 sq. ft	16,000-32,000 sq. R				
Hybrid bermudageass	075-15	5,300-10,600 sq. ft	10,600-21,309 sq. ft	21,300-43,000 sq. R				
Bahingrass	0.75-1.5	5,300-10,600 sq. ft	10,600-21,300 sq. ft	21,300-43,000 sq. #				
Zoysiagrass	0.75-1.5	5,300-10,600 sq.ft	10,600-21,300 sq. ft	21,300-43,000 sq. R				
Bultalograss	0.75-1.5	5,300-10,600 sq. it	10,600-21,300 sq. ft	21,300-43,000 sq. R				
SL Augustinegrass	0.75-1.5	5,300-10,600 sq. ft.	10,600-21,300 sq. R.	21,300-43,000 sq. ft.				
Centipydegrass	0.75-1.5	5,300-10,600 sq. ft.	10,600-21,300 sq. A.	21,300-43,000 sq. ft.				

HOW TO APPLY:

FOR ORTHO HOSE-END SPRAYERS FOR WARM SEASON TURFGRASS

- For Ortho[®] Dial N Spray Hose-End Sprayer: Measure 1½ II. oz. (3 Tbs.) into sprayer jar. Do not add water. Set dial to 1 Tbs. and spray uniformly on 1,000 sq. ft. of lawn.
- For Ortho[®] Plant & Weed Sprayette[®] Hose-End Sprayer: Measure 1½ fl. oz. (3 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray uniformly on 1,000 sq. ft. of lawn.
- For Gilmour[®] 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl. oz. and Dilution Rate of 80:1:
- For 0.75 II. oz. of product per 1.000 sq. It.; Measure 5.0 II. oz. of product into sprayer jar. Add water or fill with water to the 32 II. oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq. ft, of lawn.
- For 1.0 fl.oz. of product per 1.000 sq. ft.; Measure 6.7 fl. oz. of product into sprayer jar. Add water or fill with water to the 32 fl. oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq. ft. of lawn.
- For 1.5 fl. oz. of product per 1,000 sg. ft.: Measure 10.0 fl. oz. of product into sprayer jar. Add water or fill with water to the 32 fl. oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sg. ft. of lawn.
- For 2.0 fl. oz. of product per 1,000 sg. fl.: Measure 13.3 fl. oz. of product into sprayer jar. Add water or fill with water to the 32 fl. oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sg. ft. of lawn.
- For Gilmour[#] Metering Dial Insecticide & Fertilizer Sprayer: This sprayer can be set to meter tablespoons per gallon of water or can be set to teaspoons per gallon of water by using the converting orifice in the top of the mixing head.
- For 1.0 fl. oz. of product per 1.000 sq. fl.; Turn the metering dial until the desired setting number is next to the setting arrows on the side of the mixing head. Set the dial at 2 teaspoons per gallon. Add herbicide to the sprayer bottle. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. By squeezing the handle, you will be automatically spraying the proper spray concentration.

6571006

Page 2 of 3



An Employee-Opined Company 1217 West 12th Street Kansas City, Missouri 64101

www.pbigordon.com



NET CONTENTS: ____ U.S. GAL.

657/7-2003 AP011503 EPA REG. NO. 2217-855 EPA EST. NO. 2217-KS-1

- For 1.5 fl. oz. of product per 1.000 sq. ft.: Turn the metering dial until the desired setting number is next to the setting arrows on the side of the mixing head. Set the dial at 1 tablespoon per gallon or 3 teaspoons per gallon. Add herbicide to the sprayer bottle. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. By squeezing the handle, you will be automatically spraying the proper spray concentration.
- For 2.0 fl. oz. of product per 1,000 so. ft.: Turn the metering dial until the desired setting number is next to the setting arrows on the side of the mixing head. Set the dial at 1 tablespoon per gallon or 4 teaspoons per gallon. Add herbicide to the sprayer bottle. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. By squeezing the handle, you will be automatically spraying the proper spray concentration.

SPOT TREATMENTS FOR WARM SEASON TURFGRASS:

HAND OPERATED SPRAYERS.

For warm season turigrass, mix 0.75-1.5 fl. oz. of Speed-Zone St. Augustine Residential per 1.0 gallon of water for treatment of approximately 1,000 sq. ft. of turigrass. Apply any time the emerged broadleal weeds are susceptible. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants.

Spray volume, gals. of water	Field cences (it. oz.) of preduct	Tablespoons (ths.) of product	Area to be irealed, ay, It.
1 gallon	0.75-1.5 fl. oz.	1.5-3.0 tbs.	1,000 sq. ft.
2 gailons	1.5-3.0 fl. oz	3.0-6.0 tbs,	2.000 sq. ft.
3 galions	2.25-4.5 fl. oz.	4.5-9.0 tbs.	3,000 sq. ft.

Equal measures: 1.0 fl. oz. = 2 tablespoons.

The maximum application rate to turl is 0.5 pound 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

BROADLEAF WEEDS CONTROLLED:

Speed-Zone St. Augustine Residential will control or suppress the following broadleaf weeds and will control or suppress other broadleaf weeds that are susceptible to 2,4-D. A partial list is presented below.

BROADLEAF WEEDS

Aster, white heath & white prairie	Cinquetoil Clover	Field oxeye-daisy (*creeping oxeye)
Bedstraw	Compassplant	Filaree, whitestern
Beggarweed,	Curly dock	& redstem
creeping	Dandelion	Florida betony
Bindweed	Dayflower	Florida pusley
Black medic	Deadnettle	Ground ivy
Broadleaf plantain	Dock	Groundsei
Buckhorn plantain	Doglennel	Hawkweed
Bull thistle	English daisy	Healail
Burclover	False dandelion	Henbit
Burdock, common	(*spotted catsear &	Innocence
Buttercup, creeping	common catsear)	(Blue-eyed Mary)
Carpetweed	Field bindweed	Lambsquarters
Chickweed, common	(*morningglory &	Lawn burweed
Chicory	creeping jenny)	Lespedeza, common

Mallow, common Matchweed Mouseear chickweed Old world diamond flower Oxalis ("vellow woodsorrel & creeping woodsorrel) Parsley-piert Pennsylvania smartweed Pennywort ("dollarweed) Pepperweed Pigweed Pineappleweed Plantain

*Synonyms

Poison ivv Poison oak Prostrate knotweed (*knolweed) Puncturevine Purple cudweed Purslane Ragweed Redweed Red sorrel ("sheep sorrel) Shepherdspurse Spotted source Spurge Thistle Veronica (*corn speedwell)

Virginia buttonweed White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort) Wild carrot Wild garitc Wild garanium Wild entuce Wild mustard Wild onion Wild strawberry Wild onion Wild strawberry Wild violet Yarrow Yellow rocket

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toil free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, brinadoes, hurricanes, etc. These factors are beyond the control of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

SPEED-ZONE® is a registered trademark of PBI/Gordon Corporation.



An Employee-Owned Company 1217 West 12th Street Kanshs City, Missouri 64101



NET CONTENTS: ____ U.S. GAL.

657/7-2003 AP011503 EPA REG. NO. 2217-855 EPA EST. NO. 2217-KS-1

6571006

and the second second second second

Page 3 of 3