

2217-855

09/04/2008

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

SEP - 4 2008

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

James L. Kunstman  
PBI/Gordon Corporation  
1217 West 12<sup>th</sup> St.  
P.O. Box 014090  
Kansas City, Missouri 64101-0090

Dear Dr. Kunstman:

SUBJECT: RED Label Amendment  
EH-1382 EC Residential Herbicide  
EPA Registration No. 2217-855

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- 1) A footnote must be added to the ingredient statement, per the product chemistry review, which reads "Contains Petroleum Distillate."
- 2) As specified in the acute toxicity review, a phone number for emergency medical treatment information should be added to the label.
- 3) The first paragraph of the Environmental Hazards text currently on the label must be revised to read as follows:

"This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate."

- 4) The following text must be added to the label:

"Do not apply as a fine mist because of potential of injury to desirable plants."


"Do no apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do no apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not

over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.”

5) - Under the **Broadcast Treatments for Cool Season Turfgrass** section, add the following statements:

“Limited to 2 broadcast applications per year. Minimum of 30 days between applications is required.”

Sincerely yours,



Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

3/12

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
SEP - 4 2008

# 1382 EC RESIDENTIAL

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

## EPA Reg. No. 2217-855

2217-855

### ACTIVE INGREDIENT:

2,4-D, 2-ethylhexyl ester .....	10.49%
Mecoprop-p, acid .....	2.66%
Dicamba, acid .....	0.67%
Carfentrazone-ethyl .....	0.54%

### INERT INGREDIENTS:

.....	<u>85.64%</u>
TOTAL.....	100.00%

### THIS PRODUCT CONTAINS:

- 0.52 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.96%.
- 0.20 lbs. (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 2.66%.
- 0.05 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.67%.
- 0.04 lbs. Ethyl  $\alpha$ ,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate or 0.54%

Isomer Specific by AOAC Methods.  
TRIMEC® is a registered trademark of PBI/Gordon Corporation

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

## PRECAUTIONARY STATEMENTS

### Hazards to Human and Domestic Animals

**CAUTION:** Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves.

### Personal Hygiene Statements:

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.



<b>First Aid</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillates - vomiting may cause aspiration pneumonia.	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **General Precautions and Restrictions:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### **PRODUCT DESCRIPTION:**

This Trimec® product is a combination of four proven weed killers: 2,4-D, mecoprop-p, dicamba and carfentrazone-ethyl. Postemergent applications of this product control a wide range of broadleaf lawn weeds in residential lawns including dandelion, chickweed, spurge, pennywort (dollarweed), knotweed, plantain, clover and ground ivy (creeping Charlie).

This product offers these advantages:

- Rapid and effective weed control for common and troublesome weed species in turfgrass.
- Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed within hours of the application and plant death can occur within 7 - 14 days.

**WHERE TO USE:**

This product is intended for use on most ornamental lawns including:

Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass Perennial ryegrass Tall fescue Red or fine leaf fescues Colonial bentgrass Mixtures of cool season species	Common bermudagrass Hybrid bermudagrass Bahiagrass Zoysiagrass Buffalograss St. Augustinegrass Centipedegrass

**Turfgrass tolerance:**

The turfgrass tolerance to 1382 EC Residential may vary and temporary turfgrass yellowing may occur on St. Augustinegrass and certain varieties of hybrid bermudagrass.

1382 EC Residential may injure certain turfgrass species. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90°F.

**Prohibitions:**

- Do not use this product on carpetgrass, dichondra or lawns containing desirable clovers.
- Do not apply this product to 'Floritam' or Bitterblue' varieties of St. Augustinegrass in Florida.
- Do not apply this product to St. Augustinegrass during spring green-up which is the transition period between dormancy and active growth.
- Do not spray roots of trees and ornamentals.
- Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result.
- Do not exceed specified dosages for any area. Be particularly careful within dripline of trees and other ornamental species.

**WHEN TO APPLY:**

Spray when broadleaf weeds are young and actively growing for best results.

1382 EC Residential may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2 - 6 week intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are susceptible.

**CULTURAL TIPS FOR IMPROVED CONTROL:**

**Irrigation:**

- Do not apply this product immediately before rainfall or irrigation. Rain or watering within 4 hours after treatment may reduce effectiveness.

**Mowing:**

- Delay mowing 1 to 2 days before and after the application of this product.

**Reseeding Interval:**

- Treated areas may be reseeded 2 weeks after application.

**THREE EASY STEPS TO MIX AND APPLY:**

1. Determine (Measure) the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is presented as fl.oz. (tablespoons) per 1,000 sq.ft. of lawn.

- 2. Use hand-operated sprayers (backpack sprayers, compression sprayers, knapsack sprayers or tank type) or hose end sprayers. Carefully measure and mix the proper amount of product with water. See below for instructions for the various types of sprayers.
- 3. Spray evenly over the lawn.

**HOW MUCH TO USE:**

**1. BROADCAST TREATMENTS FOR COOL SEASON TURFGRASS:**

Use rates of 1382 EC Residential on cool season turfgrass as broadcast treatments are presented in Table 1.

TABLE 1. RATE RECOMMENDATIONS FOR COOL SEASON TURFGRASS.				
Species	Amount of Product fl.oz./1,000 sq.ft.	Treated Area, sq.ft.		
		8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container
Kentucky bluegrass	1.5 - 2.0	4,000 - 5,300 sq.ft	8,000 - 10,600 sq.ft	16,000 - 21,300 sq.ft
Perennial ryegrass	1.5 - 2.0	4,000 - 5,300 sq.ft	8,000 - 10,600 sq.ft	16,000 - 21,300 sq.ft
Tall fescue	1.5 - 2.0	4,000 - 5,300 sq.ft	8,000 - 10,600 sq.ft	16,000 - 21,300 sq.ft
Red or fine leaf fescues	1.5 - 2.0	4,000 - 5,300 sq.ft	8,000 - 10,600 sq.ft	16,000 - 21,300 sq.ft
Colonial bentgrass	1.5 - 2.0	4,000 - 5,300 sq.ft	8,000 - 10,600 sq.ft	16,000 - 21,300 sq.ft
Mixtures of cool season species	1.5 - 2.0	4,000 - 5,300 sq.ft	8,000 - 10,600 sq. ft	16,000 - 21,300 sq.ft

**HOW TO APPLY:**

**FOR ORTHO HOSE-END SPRAYERS FOR COOL SEASON TURFGRASS**

- **For Ortho® Dial'N Spray Hose-End Sprayer:** Measure 2.0 fl.oz. (4 Tbs.) into sprayer jar. Do not add water. Set dial on 1 oz. and spray uniformly on 1,000 sq.ft. of lawn.
- **For Ortho® Plant & Weed Sprayette® Hose-End Sprayer:** Measure 2.0 fl.oz. (4 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 sq.ft.
- **For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Rate of 80:1:**
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Measure 10 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Measure 13.3 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
- **For Gilmour® Metering Dial Insecticide & Fertilizer Sprayer:**
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Set the dial to 1 tablespoon (3 teaspoons) per gallon. Add 1.5 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Set the dial to 4 teaspoons per gallon. Add 2 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.

The maximum application rate is 3.0 fl. oz. of product/1,000 sq.ft. per application (0.5 lbs 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. The maximum seasonal rate is 6.0 fl. oz. of product/1,000 sq.ft. (1.0 lbs 2,4-D acid equivalent per acre), excluding spot treatments.

**2. SPOT TREATMENTS FOR COOL SEASON TURFGRASS.**

**HAND OPERATED OR PUMP-UP SPRAYERS:**

For cool season turfgrass, mix 1.5 - 2.0 fl.oz. of 1382 EC Residential per 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. Apply any time the emerged broadleaf weeds are susceptible. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants.

Spray volume, gals. Of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	1.5 - 2.0 fl.oz.	3.0 - 4.0 tbs.	1,000 sq.ft.
2 gallons	3.0 - 4.0 fl.oz.	6.0 - 8.0 tbs.	2,000 sq.ft.
3 gallons	4.5 - 6.0 fl.oz.	9.0 - 12.0 tbs.	3,000 sq.ft.

Equal measures: 1.0 fl.oz. = 2 tablespoons. 1 tablespoon = 3 teaspoons

Use rates of 1382 EC Residential on cool season turfgrass as broadcast treatments are presented in Table 2.

**TABLE 2. RATE RECOMMENDATIONS FOR WARM SEASON TURFGRASS.**

Species	Amount of Product Fl.oz./1,000 sq.ft.	Treated Area, sq.ft.		
		8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container
Common bermudagrass	1.0 - 2.0	4,000 - 8,000 sq.ft	8,000 - 16,000 sq.ft	16,000 - 32,000 sq.ft
Hybrid bermudagrass	0.75 - 1.5	5,300 - 10,600 sq.ft	10,600 - 21,300 sq. ft	21,300 - 43,000 sq.ft
Bahiagrass	0.75 - 1.5	5,300 - 10,600 sq.ft	10,600 - 21,300 sq. ft	21,300 - 43,000 sq.ft
Zoysiagrass	0.75 - 1.5	5,300 - 10,600 sq.ft	10,600 - 21,300 sq.ft	21,300 - 43,000 sq.ft
Buffalograss	0.75 - 1.5	5,300 - 10,600 sq.ft	10,600 - 21,300 sq.ft	21,300 - 43,000 sq.ft
St. Augustinegrass	0.75 - 1.5	5,300 - 10,600 sq.ft	10,600 - 21,300 sq.ft	21,300 - 43,000 sq.ft
Centipedegrass	0.75 - 1.5	5,300 - 10,600 sq.ft	10,600 - 21,300 sq.ft	21,300 - 43,000 sq.ft

**HOW TO APPLY:**

**FOR ORTHO HOSE-END SPRAYERS FOR WARM SEASON TURFGRASS**

- **For Ortho® Dial'N Spray Hose-End Sprayer:** Measure 1.5 fl.oz. (3 Tbs.) into sprayer jar. Do not add water. Set dial to 1 Tbs. and spray uniformly on 1,000 sq.ft. of lawn.
- **For Ortho® Plant & Weed Sprayette® Hose-End Sprayer:** Measure 1.5 fl.oz. (3 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray uniformly on 1,000 sq.ft. of lawn.
- **For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Rate of 80:1:**
  - For 0.75 fl.oz. of product per 1,000 sq.ft.: Measure 5.0 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 1.0 fl.oz. of product per 1,000 sq.ft.: Measure 6.7 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Measure 10.0 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Measure 13.3 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
- **For Gilmour® Metering Dial Insecticide & Fertilizer Sprayer:**
  - For 1.0 fl.oz. of product per 1,000 sq.ft.: Set the dial to 2 teaspoons per gallon. Add 1.0 ounce of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire

- o contents uniformly on area to be treated.
- o **For 1.5 fl.oz. of product per 1,000 sq.ft.:** Set the dial to 1 tablespoons (3 teaspoons) per gallon. Add 1.5 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.
- o **For 2.0 fl.oz. of product per 1,000 sq.ft.:** Set the dial to 4 teaspoons per gallon. Add 2 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.

**SPOT TREATMENTS FOR WARM SEASON TURFGRASS:  
HAND OPERATED OR PUMP-UP SPRAYERS.**

For warm season turfgrass, mix 0.75 - 1.5 fl.oz. of 1382 EC Residential per 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. Apply any time the emerged broadleaf weeds are susceptible. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants.

Spray volume, gals. Of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	0.75 - 1.5 fl.oz.	1.5 - 3.0 tbs.	1,000 sq.ft.
2 gallons	1.5 - 3.0 fl.oz.	3.0 - 6.0 tbs.	2,000 sq.ft.
3 gallons	2.25 - 4.5 fl.oz.	4.5 - 9.0 tbs.	3,000 sq.ft.
Equal measures: 1.0 fl.oz. = 2 tablespoons. 1 tablespoon = 3 teaspoons			

The maximum application rate is 3.0 fl. oz. of product/1,000 sq.ft. per application (0.5 lbs 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. The maximum seasonal rate is 6.0 fl. oz. of product/1,000 sq.ft. (1.0 lbs 2,4-D acid equivalent per acre), excluding spot treatments.

**Ready-to-use hose-end sprayers supplied by Green Garden Products Company® and other suppliers.** Dilution factor of 200:1 with a sprayer jar capacity of 8 and 16 fl. oz.

This retail sprayer bottle will cover the approximate areas described below when this product is applied to established turfgrass. Over-application or rates above those recommended on this label can cause turf injury.

**Four Steps To Apply:**

1. Measure the *total* (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
  2. Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the OFF position.  
To BEGIN spraying: Turn on water at faucet. Point nozzle toward lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Spray evenly over the measured area.
  3. SPRAY: Spray evenly over the measured area. Walk at a steady pace. Refer to the view strip [graduated scale] on the side of the container to guide coverage. The xx fl. oz. container will treat xxxx sq.ft. of turfgrass.
  4. To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose
- 
1. Measure the *total* (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
  2. Connect a garden hose to the hose-end sprayer. Make sure the control valve on the



nozzle is in the OFF position.

To BEGIN spraying: Turn on water at faucet. Point nozzle toward lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Spray evenly over the measured area.

3. **SPRAY:** Spray evenly over the measured area. Walk at a steady pace. Refer to the view strip [graduated scale] on the side of the container to guide coverage. The xx fl. oz. container will treat xxxx sq.ft. of turfgrass. Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the OFF position.
4. **To STOP spraying:** Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose

Species	8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container
Kentucky bluegrass Perennial ryegrass Tall fescue Red or fine leaf fescues Colonial bentgrass Mixtures of cool season species	5,000 sq.ft.	10,000 sq.ft.
Common bermudagrass Hybrid bermudagrass Bahia grass Zoysiagrass Buffalograss St. Augustinegrass <sup>1</sup> Centipedegrass	10,000 sq.ft.	20,000 sq.ft.
Do not apply this product to 'Floratum' or Bitterblue' varieties of St. Augustinegrass in Florida.		
The maximum application rate is 3.0 fl. oz. of product/1,000 sq.ft. per application (0.5 lbs 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. The maximum seasonal rate is 6.0 fl. oz. of product/1,000 sq.ft. (1.0 lb 2,4-D acid equivalent per acre), excluding spot treatments.		

**BROADLEAF WEEDS CONTROLLED:**

1382 EC Residential will control or suppress the following broadleaf weeds and will control or suppress other broadleaf weeds that are susceptible to 2,4-D. A partial list is presented below.

BROADLEAF WEEDS			
Aster, white heath & white prairie	Dogfennel	Mallow, common	Shepherdspurse
Bedstraw	English daisy	Matchweed	Spotted spurge
Beggarweed, creeping	False dandelion (*spotted catsear & common catsear)	Mouseear chickweed	Spurge
Bindweed	Field bindweed	Old world diamond flower	Thistle
Black medic	(*morningglory & creeping jenny)	<i>Oxalis</i> (*yellow woodsorrel & creeping woodsorrel)	<i>Veronica</i> (*corn speedwell)
Broadleaf plantain	Field oxeye-daisy (*creeping oxeye)	Parsley-piert	Virginia buttonweed
Buckhorn plantain	Filaree, whitestem & redstem	Pennsylvania smartweed	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Bull thistle	Florida betony	Pennywort (*dollarweed)	Wild carrot
Burclover	Florida pusley	Pepperweed	Wild garlic
Burdock, common	Ground ivy (*creeping Charlie)	Pigweed	Wild geranium
Buttercup, creeping	Groundsel	Pineappleweed	Wild lettuce
Carpetweed	Hawkweed	Plantain	Wild mustard
Chickweed, common	Healall	Poison ivy	Wild onion
Chicory	Henbit	Poison oak	Wild strawberry
Cinquefoil	Innocence (Blue-eyed Mary)	Prostrate knotweed (*knotweed)	Wild violet
Clover	Lambsquarters	Puncturevine	Yarrow
Compassplant	Lawn burweed	Purple cudweed	Yellow rocket
Curly dock	Lespedeza, common	Purslane	
Dandelion		Ragweed	
Dayflower		Redweed	
Deadnettle		Red sorrel (*sheep sorrel)	
Dock			
*Synonyms			

## **STORAGE & DISPOSAL**

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**CONTAINER DISPOSAL:** **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (800-XXX-XXXX) for disposal instructions. Never place unused product down any indoor or outdoor drain.

## **LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

## **APPENDIX**

### **1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

### **2. Advertising claims appropriate for the bag size:**

- Rainfast in as little as 3 hours
- Rainfast [Rainproof] in 3 hours
- Visible results in just 24 hours
- Clover control in as little as one week
- Proven cool-weather performance
- Consistently fast control on tough weeds like clover, plantain, ground ivy and spurge [various other listed weeds].
- [This product] Smokes weeds, fast
- From the makers of Trimec® herbicides.
- For information call XXX-XXX-XXXX [contact [www.xxx-xxxx.com](http://www.xxx-xxxx.com)]

## **DOCUMENT CONTROL INFORMATION**

1. Unique Label Identifier: 002217-00855.20071105

**2. Previous Version:** AP052907

**3. Reason for Issue:** Various changes pertaining to re-registration. Re-format label.