



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

March 25, 2025

Eric D. Smith  
Director, Regulatory Affairs  
PBI/Gordon Corporation  
P.O. Box 860350  
Shawnee, KS 66286

Subject: Label Amendment - Registration Review Mitigation for Carfentrazone-ethyl & Mecoprop-p  
Product Name: 1382 EC RESIDENTIAL HERBICIDE  
EPA Registration Number: 2217-855  
Application Dates: May 1, 2018; July 5, 2022; & January 30, 2024  
Decision Numbers: 596732 & 596733

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Carfentrazone-ethyl & Mecoprop-p Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at (202) 566-0636, or via email at [carr.caleb@epa.gov](mailto:carr.caleb@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read 'Linda Arrington', with a stylized flourish at the end.

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

# 1382 EC RESIDENTIAL

## EPA Reg. No. 2217-855

### ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester .....	10.49%
Mecoprop-p, acid .....	2.66%
Dicamba, acid .....	0.67%
Carfentrazone-ethyl .....	0.54%
OTHER INGREDIENTS: .....	<u>85.64%</u>
TOTAL .....	100.00%

### THIS PRODUCT CONTAINS:

- 0.52 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.96%.
- 0.20 lb (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 2.66%.
- 0.05 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.67%.
- 0.04 lb Ethyl  $\alpha$ ,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate or 0.54%


Isomer Specific by AOAC Methods.

Contains Petroleum Distillate.

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**



See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid.

**Net Contents:** \_\_\_\_\_

**EPA Est. No.** \_\_\_\_\_

**Company Name and Address:**

PBI/Gordon Corporation

P.O. Box 860350

Shawnee, Kansas 66286



## **PRECAUTIONARY STATEMENTS**

### **Hazards to Human and Domestic Animals**

**CAUTION:** Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves.

### **Personal Hygiene Statements**

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

<b>First Aid</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give any liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillate - vomiting may cause aspiration pneumonia.	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**Fish Advisory Statement:** This product may be hazardous to aquatic organisms, particularly in clear, shallow water bodies that are adjacent to treated areas. Transport to water by runoff or spray drift of this product in areas where surface water is present, or intertidal areas below the mean high water mark, should be avoided. Do not contaminate water when disposing of equipment wash water or rinsate.

**Non-target Organism Advisory Statement:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **1. Product Description**

This Trimec® product is a combination of four proven weed killers: 2,4-D, mecoprop-p, dicamba and carfentrazone-ethyl. Postemergent applications of this product control a wide range of listed broadleaf lawn weeds in residential lawns including dandelion, chickweed, spurge, pennywort (dollarweed), knotweed, plantain, clover and ground ivy (creeping Charlie).

This product offers these advantages:

- Rapid and effective weed control for common and troublesome weed species as listed in turfgrass
- Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed within hours of the application and plant death can occur within 7 to 14 days.

## 2. Use Restrictions

- Only use for sites, pests, and application methods specified on this labeling.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- The maximum broadcast application rate is 3.0 fl.oz. of product/1,000 sq.ft. per application [(0.5 lb 2,4-D ae, 0.2 lb MCPP-p ae, 0.05 lb dicamba ae, and 0.04 lb carfentrazone ai per acre per application)].
- The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications.
- The maximum annual broadcast application rate is 6.0 fl.oz. of product/1,000 sq.ft. [(1.0 lb 2,4-D ae, 0.4 lb MCPP-p ae, 0.1 lb dicamba ae, and 0.08 lb carfentrazone ai per acre per year)], excluding spot treatments.
- The maximum spot treatment application rate is 3.0 fl.oz. of product/1,000 sq.ft. per application [(0.5 lb 2,4-D ae, 0.2 lb MCPP-p ae, 0.05 lb dicamba ae, and 0.04 lb carfentrazone ai per acre per application)].
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on carpetgrass, dichondra or lawns containing desirable clovers.
- Do not apply this product to 'Floritam' or Bitterblue' varieties of St. Augustinegrass in Florida.
- Do not apply this product to St. Augustinegrass during spring green-up which is the transition period between dormancy and active growth.
- Do not spray roots of trees and ornamentals.
- Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result.
- Do not exceed specified dosages for any area. Be particularly careful within dripline of trees and other ornamental species.
- Do not apply as a fine mist because of potential injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product runoff, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.

## 3. Where To Use

This product is intended for use on most ornamental lawns including:

- **Cool Season Turfgrass:** Kentucky bluegrass, Perennial ryegrass, tall fescue, Red or fine leaf fescues, colonial bentgrass, and mixtures of cool-season species
- **Warm Season Turfgrass:** common Bermudagrass, hybrid Bermudagrass, bahiagrass, zoysiagrass, buffalograss, St. Augustinegrass, and centipedegrass

Turfgrass tolerance: The turfgrass tolerance to 1382 EC Residential may vary and temporary turfgrass

yellowing may occur on St. Augustinegrass and certain varieties of hybrid bermudagrass. 1382 EC Residential may injure certain turfgrass species. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90°F.

#### 4. When To Apply

Spray when broadleaf weeds are young and actively growing for best results.

1382 EC Residential may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2 to 6 week intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are susceptible.

#### CULTURAL TIPS FOR IMPROVED CONTROL:

##### Irrigation:

- Do not apply this product immediately before rainfall or irrigation. Rain or watering within 4 hours after treatment may reduce effectiveness.

##### Mowing:

- Delay mowing 1 to 2 days before and after the application of this product.

##### Reseeding Interval:

- Treated areas may be reseeded 2 weeks after application.

#### THREE EASY STEPS TO MIX AND APPLY:

1. Determine (Measure) the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is presented as fl.oz. (tablespoons) per 1,000 sq.ft. of lawn.
2. Use hand-operated sprayers (backpack sprayers, compression sprayers, knapsack sprayers or tank type) or hose end sprayers. Carefully measure and mix the proper amount of product with water. See below for instructions for the various types of sprayers.
3. Spray evenly over the lawn.

#### 5. How Much To Use

##### BROADCAST TREATMENTS FOR COOL SEASON TURFGRASS:

Use rates of 1382 EC Residential on cool season turfgrass as broadcast treatments are presented in Table 1.

TABLE 1. RATE RECOMMENDATIONS FOR COOL SEASON TURFGRASS.				
Species	Amount of Product	Treated Area, sq.ft.		
	fl.oz./1,000 sq.ft.	8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container
Kentucky bluegrass, Perennial ryegrass, Tall fescue, Red or fine leaf fescues, Colonial bentgrass, Mixtures of cool season species	1.5 to 2.0	4,000 to 5,300 sq.ft.	8,000 to 10,600 sq.ft.	16,000 to 21,300 sq.ft.

## 6. How To Apply

### FOR ORTHO HOSE-END SPRAYERS FOR COOL SEASON TURFGRASS

- **For Ortho® Dial'N Spray Hose-End Sprayer:** Measure 2.0 fl.oz. (4 Tbs.) into sprayer jar. Do not add water. Set dial on 1 oz. and spray uniformly on 1,000 sq.ft. of lawn.
- **For Ortho® Plant & Weed Sprayette® Hose-End Sprayer:** Measure 2.0 fl.oz. (4 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 sq.ft.
- **For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Rate of 80:1:**
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Measure 10 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Measure 13.3 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
- **For Gilmour® Metering Dial Insecticide & Fertilizer Sprayer:**
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Set the dial to 1 tablespoon (3 teaspoons) per gallon. Add 1.5 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Set the dial to 4 teaspoons per gallon. Add 2 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.

### SPOT TREATMENTS FOR COOL SEASON TURFGRASS.

#### HAND OPERATED OR PUMP-UP SPRAYERS:

For cool season turfgrass, mix 1.5 to 2.0 fl.oz. of 1382 EC Residential per 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. Apply any time the emerged broadleaf weeds are susceptible. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants.

Spray volume, gals. Of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	1.5 to 2.0 fl.oz.	3.0 to 4.0 tbs.	1,000 sq.ft.
2 gallons	3.0 to 4.0 fl.oz.	6.0 to 8.0 tbs.	2,000 sq.ft.
3 gallons	4.5 to 6.0 fl.oz.	9.0 to 12.0 tbs.	3,000 sq.ft.
Equal measures: 1.0 fl.oz. = 2 tablespoons. 1 tablespoon = 3 teaspoons			

Use rates of 1382 EC Residential on warm season turfgrass as broadcast treatments are presented in Table 2.

**TABLE 2. RATE RECOMMENDATIONS FOR WARM SEASON TURFGRASS.**

Species	Amount of Product	Treated Area, sq.ft.		
	Fl.oz./1,000 sq.ft.	8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container
Common bermudagrass	1.0 to 2.0	4,000 to 8,000 sq.ft	8,000 to 16,000 sq.ft.	16,000 to 32,000 sq.ft.



Hybrid bermudagrass, Bahia grass, zoysiagrass, Buffalograss, St. Augustine grass, Centipedegrass	0.75 to 1.5	5,300 to 10,600 sq.ft	10,600 to 21,300 sq. ft.	21,300 to 43,000 sq.ft.
---	-------------	-----------------------	--------------------------	-------------------------

#### FOR ORTHO HOSE-END SPRAYERS FOR WARM SEASON TURFGRASS

- **For Ortho® Dial'N Spray Hose-End Sprayer:** Measure 1.5 fl.oz. (3 Tbs.) into sprayer jar. Do not add water. Set dial to 1 Tbs. and spray uniformly on 1,000 sq.ft. of lawn.
- **For Ortho® Plant & Weed Sprayette® Hose-End Sprayer:** Measure 1.5 fl.oz. (3 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray uniformly on 1,000 sq.ft. of lawn.
- **For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Rate of 80:1:**
  - For 0.75 fl.oz. of product per 1,000 sq.ft.: Measure 5.0 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 1.0 fl.oz. of product per 1,000 sq.ft.: Measure 6.7 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Measure 10.0 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Measure 13.3 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.
- **For Gilmour® Metering Dial Insecticide & Fertilizer Sprayer:**
  - For 1.0 fl.oz. of product per 1,000 sq.ft.: Set the dial to 2 teaspoons per gallon. Add 1.0 ounce of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.
  - For 1.5 fl.oz. of product per 1,000 sq.ft.: Set the dial to 1 tablespoons (3 teaspoons) per gallon. Add 1.5 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.
  - For 2.0 fl.oz. of product per 1,000 sq.ft.: Set the dial to 4 teaspoons per gallon. Add 2 ounces of herbicide to the sprayer bottle for each 1,000 sq. ft. of lawn to be treated. Insert siphon tube into the base of the sprayer head. Attach water hose to sprayer. Squeeze the handle and spray entire contents uniformly on area to be treated.

#### SPOT TREATMENTS FOR WARM SEASON TURFGRASS: HAND OPERATED OR PUMP-UP SPRAYERS.

For warm season turfgrass, mix 0.75 to 1.5 fl.oz. of 1382 EC Residential per 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. Apply any time the emerged broadleaf weeds are susceptible. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants.

Spray volume, gals. Of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	0.75 to 1.5 fl.oz.	1.5 to 3.0 tbs.	1,000 sq.ft.
2 gallons	1.5 to 3.0 fl.oz.	3.0 to 6.0 tbs.	2,000 sq.ft.
3 gallons	2.25 to 4.5 fl.oz.	4.5 to 9.0 tbs.	3,000 sq.ft.
Equal measures: 1.0 fl.oz. = 2 tablespoons. 1 tablespoon = 3 teaspoons			

**Ready-to-use hose-end sprayers supplied by Green Garden Products Company® and other suppliers.** Dilution factor of 200:1 with a sprayer jar capacity of 8 and 16 fl.oz.



This retail sprayer bottle will cover the approximate areas described below when this product is applied to established turfgrass. Over-application or rates above those recommended on this label can cause turf injury.

#### Four Steps To Apply:

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
2. Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the OFF position.  
To BEGIN spraying: Turn on water at faucet. Point nozzle toward lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Spray evenly over the measured area.
3. SPRAY: Spray evenly over the measured area. Walk at a steady pace. Refer to the view strip [graduated scale] on the side of the container to guide coverage. The xx fl.oz. container will treat xxxx sq.ft. of turfgrass.
4. To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose

Species	8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container
Kentucky bluegrass Perennial ryegrass Tall fescue Red or fine leaf fescues Colonial bentgrass Mixtures of cool season species	5,000 sq.ft.	10,000 sq.ft.
Common bermudagrass Hybrid bermudagrass Bahia grass Zoysiagrass Buffalograss St. Augustinegrass <sup>1</sup> Centipedegrass	10,000 sq.ft.	20,000 sq.ft.
<sup>1</sup> Do not apply this product to 'Floratum' or Bitterblue' varieties of St. Augustinegrass in Florida.		

## 5. Broadleaf Weeds Controlled [79]

1382 EC Residential will control or suppress the following broadleaf weeds.

Aster, white heath & white prairie	Curly dock	Hawkweed
Bedstraw	Dandelion	Healall
Beggarweed, creeping	Dayflower	Henbit
Bindweed	Deadnettle	Innocence (Blue-eyed Mary)
Black medic	Dock	Lambsquarters
Broadleaf plantain	Dogfennel	Lawn burweed
Buckhorn plantain	English daisy	Lespedeza, common
Bull thistle	False dandelion (*spotted catsear & common catsear)	Mallow, common
Burclover	Field bindweed (*morningglory & creeping jenny)	Matchweed
Burdock, common	Field oxeye-daisy (*creeping oxeye)	Mouseear chickweed
Buttercup, creeping	Filaree, whitestem & redstem	Old world diamond flower
Carpetweed	Florida betony	<i>Oxalis</i> (*yellow woodsorrel & creeping woodsorrel)
Chickweed, common	Florida pusley	Parsley-piert
Chicory	Ground ivy (*creeping Charlie)	Pennsylvania smartweed
Cinquefoil	Groundsel	Pennywort (*dollarweed)
Clover		Pepperweed
Compassplant		

Pigweed  
Pineappleweed  
Plantain  
Poison ivy  
Poison oak  
Prostrate knotweed (\*knotweed)  
Puncturevine  
Purple cudweed  
Purslane  
Ragweed  
Redweed

Red sorrel (\*sheep sorrel)  
Shepherdspurse  
Spotted spurge  
Spurge  
Thistle  
*Veronica* (\*corn speedwell)  
Virginia buttonweed  
White clover (\*Dutch clover,  
honeysuckle clover, white trefoil,  
& purplewort)  
Wild carrot

Wild garlic  
Wild geranium  
Wild lettuce  
Wild mustard  
Wild onion  
Wild strawberry  
Wild violet  
Yarrow  
Yellow rocket

\*Synonyms

## STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

## APPENDIX

### 1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

#### Herbicide Benefit Claims:

- Clover control in as little as one week
- Consistently fast control on tough weeds like clover, plantain, ground ivy and spurge [various other listed weeds].
- [This product] Smokes weeds, fast
- Consistently fast – controls tough weeds like clover, plantain, ground ivy, spurge & other listed weeds

General Marketing Claims:

- Rainfast in as little as 3 hours
- Rainfast [Rainproof] in 3 hours
- Visible results in just 24 hours
- Proven cool-weather performance
- From the makers of Trimec® herbicides.
- For information call [contact pbigordon.com]
- In the event of a medical or chemical emergency, contact ChemTel Inc North America 1-800-255-3924

## **DOCUMENT CONTROL INFORMATION**

**1. Unique Label Identifier:** 002217-00855.20240130.amend-proposed-clean.doc

**2. Reason for Issue:** Comments on MCPP-p and Carfentrazone-ethyl Interim Registration Review Decision

<b>Internal Usage Notes</b>
Associated Container Label: SpeedZone St. Augustine Residential; Supplemental Distributors