

2217-859

03/10/2005

1/22

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 2217-859	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) EH-1401 Herbicide	PM# Product Manager—Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
MAR 10 2005

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION per Pesticide Registration Notice (PRN) 98-10:

- We ask to add an alternate set of spreader settings which may be presented on the labeling of our supplemental distributors. This is presented in its entirety on page 9 (sublabel 1) and page 17 (sublabel 2) of the enclosed labeling.
- We ask to add one advertising claim which may be presented on the labeling of our supplemental distributors. This additional claim appears highlighted on page 21 of the enclosed labeling.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.1

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name James L. Kunstman, Ph.D.	Title Director of Regulatory Services	Telephone No. (Include Area Code) 816-430-6292
Certification <i>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</i>		6. Date Application Received (Stamped)
2. Signature 	3. Title Director of Regulatory Services	
4. Typed Name James L. Kunstman, Ph.D.	5. Date February 21, 2005	

2/22

SUBLABEL 1: Intended for commercial users.

EH-1401 HERBICIDE

In the State of New York: This product may be applied only by commercial applicators at no more than 218 pounds (0.35 lb. of active ingredient) per acre (5.0 lb. per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

ACTIVE INGREDIENTS:

2,4-D.....	0.64%
Mecoprop-p.....	0.14%
Dicamba.....	0.06%
Dithiopyr.....	0.16%

OTHER INGREDIENTS:	<u>99.00%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 12.80 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
- 2.80 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
- 1.20 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
- 3.20 lbs dithiopyr per ton

CAS Registry Numbers:

- 2,4-D CAS No. 94-75-7
- Mecoprop-p CAS No. 16484-77-8
- Dicamba CAS No. 1918-00-9
- Dithiopyr CAS No. 97886-45-8

Isomer Specific by AOAC Methods
Contains the single isomer form of Mecoprop-p.

EPA Est. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.

DIMENSION® is a registered trademark of Dow AgroSciences, L.L.C.
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N).....	x.x%
x.x% Ammoniacal Nitrogen	
x.x% Water Insoluble Nitrogen	
x.x% Urea Nitrogen	
x.x% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅).....	x.x%
Soluble Potash (K ₂ O).....	x.x%
Iron (Fe).....	x.x%
Nutrient Sources: (Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer).	

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements.

NOTICE: Read the entire label. Use only according to label directions.

NET WEIGHT: (16, 18, 20, 30,40, or 50) POUNDS

AP022105
 EPA Reg. No. 2217-859
MANUFACTURED BY:

All Employee-Owned Company
 1217 West 12th Street
 Kansas City, Missouri 64101
 Telephone: 1-800-821-7925

NOTIFICATION
MAR 16 2005



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

3/22

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or if swallowed. Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

Personal Hygiene Statements: After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-800-xxx-xxxx.</p>	

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water.

Drift or runoff from treated turf may adversely affect nontarget plants and aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Do not contaminate water when disposing of equipment washwaters.

When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOG. OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

Re-entry Restrictions—Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Read all precautions and directions for use carefully before applying this product.

4/22

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Paper and plastic bags: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

EH-1401 Herbicide provides postemergent control of over 200 broadleaf weeds and preemergent or early postemergent control of many annual weed grasses in established cool and warm season turf.

ORNAMENTAL LAWNS AND TURFGRASS:

WHERE TO USE:

EH-1401 Herbicide is intended to be applied by lawn care operators, maintenance applicators, and service technicians for use in ornamental lawns and turf established in institutional, ornamental, and residential/domestic sites. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (excluding greens), and office buildings. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings. Finally, residential/domestic sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

EH-1401 Herbicide is recommended for the following established turf species:

Established Cool Season Turfgrasses—Kentucky bluegrass, perennial ryegrass, fine and tall fescues.

Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

Established Warm Season Turfgrasses—Bahia grass, bermudagrass, buffalograss, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam').

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Do not use EH-1401 Herbicide on creeping bentgrass, colonial bentgrass, dichondra, carpetgrass, or 'Floritam' St. Augustinegrass.

Note: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area at recommended use rates, prior to initiating large scale use.

USE PRECAUTIONS AND RESTRICTIONS

Precautions for Reseeding, Overseeding, or Sprigging:

- Reseeding, overseeding or sprigging of areas treated with this product within 10 weeks after a single application of this product, or within 16 weeks after a split application program totaling 5.0 lbs./1,000 sq. ft. (0.38 lb. active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.

Use of EH-1401 Herbicide on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

Do not use this product where desirable clovers are present or apply (spread) the granules onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosage for any area; be particularly careful within dripline of trees and other ornamental species.

Do not use clippings from treated areas for mulching, vegetables, or fruit trees.

Do not contaminate domestic or irrigation waters.

- WEEDS CONTROLLED -			
When applied as directed, EH-1401 Herbicide will control the following weeds:			
Annual Bluegrass (<i>Poa annua</i>)	Barnyardgrass	Crowfootgrass	Kikuyugrass (seedling)
Annual Ryegrass	Crabgrass (large, smooth, and Southern)	Foxtail (green and yellow) Goosegrass	Smutgrass (seedling)

- WEEDS CONTROLLED (Continued) -

Broadleaf Weeds:

Annual yellow sweetclover	Clover, white	Musk thistle	Spiny amaranth
Aster	Cockle	Mustard	Spiny cocklebur
Austrian fieldcress	Cocklebur	Narrowleaf plantain	Spiny sowthistle
Bedstraw	Common mullein	Narrowleaf vetch	Spotted catsear
Beggarticks	Creeping jenny	Nettle	Spotted spurge
Betony, Florida	Cudweed	Orange hawkweed	Spurweed
Bindweed, field	Curly dock	Oriental cocklebur	Stinging nettle
Bird vetch	Daisy, English	Oxalis	Strawberry, India mock
Bitter wintercress	Daisy fleabane	Parsley-piert	Tall nettle
Bittercress, hairy	Daisy, oxeye	Parsnip	Tall vervain
Bitterweed	Dandelion	Pearlwort	Tansy ragwort
Black-eyed Susan	Dichondra	Pennycress	Tansy mustard
Black medic	Dogfennel	Pepperweed	Tanweed
Black mustard	Dollarweed	Pigweed	Thistle
Blackseed plantain	False dandelion	Pineappleweed	Trailing crownvetch
Blessed thistle	Falseflax	Pineywoods bedstraw	Tumble mustard
Bloodflower milkweed	False sunflower	Plains tickseed	Tumble pigweed
Blue lettuce	Fiddleneck	Plantain	Velvetleaf
Blue vervain	Florida pusley	Poison ivy	Venice mallow
Bracted plantain	Frenchweed	Poison oak	Virginia buttonweed
Brassbuttons	Galinsoga	Pokeweed	Virginia creeper
Bristly oxtongue	Goathead	Poorjoe	Virginia pepperweed
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly lettuce	Western clematis
Broomweed	Gumweed	Prickly sida	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild aster
Bull thistle	Healall	Prostrate spurge	Wild buckwheat
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild carrot
Burclover	Heath aster	Puncturevine	Wild four-o'clock
Burdock	Hedge bindweed	Purslane, common	Wild garlic
Burning nettle	Hedge mustard	Ragweed	Wild geranium
Burweed	Henbit	Red sorrel	Wild lettuce
Buttercup	Hoary cress	Redroot pigweed	Wild marigold
Buttonweed	Hoary plantain	Redstem filaree	Wild mustard
Canada thistle	Hoary vervain	Rough cinquefoil	Wild onion
Carolina geranium	Horsenettle	Rough fleabane	Wild parsnip
Carpetweed	Jimsonweed	Russian pigweed	Wild radish
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild rape
Catnip	Knotweed	Scotch thistle	Wild sweet potato
Catsear	Kochia	Sheep sorrel	Wild vetch
Chickweed, common	Lambsquarters	Sheperdspurse	Woodsorrel
Chickweed, mouseear	Lespedeza	Slender plantain	Woolly croton
Chicory	Mallow	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Matchweed	Smooth dock	Woolly plantain
Clover, crimson	Mexicanweed	Smooth pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Morningglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, sweet	Mugwort	Speedwell	and other broadleaf weeds

WHEN TO APPLY

Established Lawns: Apply when broadleaf weeds are actively growing and prior to germination or tillering of annual grasses. EH-1401 Herbicide may be applied as a single application or as a split application in spring, summer or fall.

Newly Seeded Areas: When establishing a new lawn of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied 2 weeks after the first sign of grass germination. For other turfgrass varieties, wait until the grass is well established (3 mowings) before applying this product.

HOW TO APPLY

For optimum weed control apply to recently mowed turf when foliage is moist. When possible, delay mowing or irrigation for several days after treatment. Uniform coverage is essential for best results.

Application Directions for Turf

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The dithiopyr component in this product is not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Use of split (half-rate applications) spaced 5-10 weeks apart may provide improved weed control.

Control of Crabgrass

Preemergence and Early Postemergence Control

- This product provides preemergent control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass.
- This product also provides early postemergent control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergent crabgrass control is a 2-4 week longer period of time (window of application) during which crabgrass control can be achieved.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer or fall as described in the following use programs.

Spring Applications

For applications made in the spring or early summer, use an appropriate application rate corresponding to one of the two control programs listed in Table 1. Specific rate recommendations are based on the user's location, the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the users' location within a region.

Use Program 1 for preemergent control at sites where turfgrass is cut relatively high (e.g., homeowner lawns). This program provides 3-5 months of preemergent crabgrass control. This program will also provide early postemergent control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

Use Program 2 for preemergent control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways). And (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergent crabgrass control. This program may also be used for early postemergent control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

Maximum use rate restrictions:

- In the State of New York: This product may be applied only by commercial applicators at no more than 218 pounds (0.35 lb. of active ingredient) per acre (5.0 lb. per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.
- The maximum application rate to turf is 1.4 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Table 1. Use Rates

Region	Application Rates	Program 1	Program 2
All states except NY, and parts of states not included in transition, south, coastal south or west.	Lbs./acre	130	152
	Lbs./1,000 sq. ft.	3.0	3.5
	Lbs. ai/acre	0.21	0.25
New York State only	Lbs./acre	152	218
	Lbs. /1,000 sq. ft. ft.	3.5	5.0
	Lbs. ai/acre	0.25	0.35
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT & RI.	Lbs./acre	152	218
	Lbs. /1,000 sq. ft.	3.5	5.0
	Lbs. ai/acre	0.25	0.35
South: AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX.	Lbs./acre	218	152 + 152
	Lbs. /1,000 sq. ft.	5.0	3.5 + 3.5†
	Lbs. ai/acre	0.35	0.25 + 0.25
Coastal South: HI, FL & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.	Lbs./acre	152 + 152	196 + 196
	Lbs. /1,000 sq. ft.	3.5 + 3.5†	4.5 + 4.5†
	Lbs. ai/acre	0.25 + 0.25	0.31 + 0.31
West: AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	Lbs./acre	130—152	152—196
	Lbs. /1,000 sq. ft.	3.0—3.5†	3.5—4.5†
	Lbs. ai/acre	0.21—0.25	0.25—0.31

† Split applications may be made 5—10 weeks apart.

Use of Split or Sequential Applications: Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with the rates in Table 1 being split between two applications made 5—10 weeks apart. When longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application, provided the maximum application rate per year is not exceeded. Refer to the "Maximum Use Rates" section above and separate rate restrictions for the State of New York.

Fall Applications:

This product can also be applied in the late summer through early fall (late August through November) at use rates listed in Table 1 for "Program 2" to provide control of crabgrass through the early part of the next following spring for season long control of crabgrass.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS:

- This product may be applied with rotary-type spreaders. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.
- Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.
- Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips or overlaps during application
- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 285 square feet and adjust to apply at the rate of 1 pound of product per 285 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
- For 4.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 225 square feet and adjust to apply at the rate of 1 pound of product per 225 square feet.
- For 5.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 200 square feet and adjust to apply at the rate of 1 pound of product per 200 square feet.
- Check equipment frequently to verify calibration and proper functioning.

Spreader	Delivers ___ lbs./1,000 sq. ft.				
	3.0 lbs.	3.5 lbs	4.0 lbs	4.5 lbs	5.0 lbs.
LESCO® Rotary calibration gauge setting #	X	X	X	X	X
Ortho® Drop (5000)	X	X	X	X	X
Ortho® Rotary (6000)	X	X	X	X	X
Precision® Red Devil® Rotary	X	X	X	X	X
Quaker® Rotary	X	X	X	X	X
Republic® EZ Drop	X	X	X	X	X
Republic® EZ Grow	X	X	X	X	X
Republic® EZ Rotary	X	X	X	X	X
Scotts® Drop (PF3)	X	X	X	X	X
Scotts® Easy Green	X	X	X	X	X
Scotts® Speedy Green	X	X	X	X	X
True Temper Rotary	X	X	X	X	X

These spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.


If your spreader model does not appear below, please call toll-free 800-xxx-xxxx, 8 A.M. – 5 P.M. EDT, Monday—Friday for additional model settings.

10/22

Alternate Spreader Settings:


Suggested Spreader Settings

These spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Apply in a single pass at a rate of 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.

Broadcast Spreaders	Setting
Agr-Fab® Rotary	4
Cyclone® Rotary	3½
Earthway® Rotary	14
Greenview® Jet-Spreader® SS1	14
Lesco® Rotary (dial A-S)(number dial)	H/14
Red Devil® Rotary (1-20)	5
Republic® Rotary	4
Scotts® EasyGreen®	23
Scotts® EdgeGuard®	4½
Scotts® SpeedyGreen® 2000, 3000	4½
Sears® Craftsman Rotary	4
Sta-Green® Rotary (1-10)	2½
True Temper® Rotary	3½
Vigoro® Rotary (1-10)	2½



Apply in a single pass at a rate of 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders	Setting
Agr-Fab® Drop	4
Greenview® Drop-Spreader® XP1	18
Red Devil® Drop (1-20)	9
Republic® Drop	5
Scotts® Drop AccuGreen (2-18)	7
Scotts® Drop PF1 (2-18)	5
Sta-Green® Drop (1-5)	3½



If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

[Note: Additional spreader models and settings may be added or substituted by the supplemental registrant provided they deliver the labeled rate of product.]

[Note: Spreader settings may be varied by the supplemental registrant as different grades of fertilizer are used, provided they deliver the labeled rate of pesticide.]

WARRANTY

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to manufacturer, and buyer assumes the risk of any such use.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on this label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of seller. All such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at sellers election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall seller be liable for consequential or incidental damages or losses.

SUBLABEL 2: Intended for homeowners.

EH-1401 HERBICIDE

ACTIVE INGREDIENTS:

2,4-D.....	0.64%
Mecoprop-p.....	0.14%
Dicamba.....	0.06%
Dithiopyr.....	0.16%

OTHER INGREDIENTS:	<u>99.00%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 12.80 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
- 2.80 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
- 1.20 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
- 3.20 lbs dithiopyr per ton
- Isomer Specific by AOAC Methods
- Contains the single isomer form of Mecoprop-p.

CAS Registry Numbers:

- 2,4-D CAS No. 94-75-7
- Mecoprop-p CAS No. 16484-77-8
- Dicamba CAS No. 1918-00-9
- Dithiopyr CAS No. 97886-45-8

EPA File Symbol 2217-ILO

DIMENSION® is a registered trademark of Dow AgroSciences, L.L.C.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N).....	x.x%
x.x% Ammoniacal Nitrogen	
x.x% Water Insoluble Nitrogen	
x.x% Urea Nitrogen	
x.x% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅).....	x.x%
Soluble Potash (K ₂ O).....	x.x%
Iron (Fe).....	x.x%
Nutrient Sources: (Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer).	

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements.

NOTICE: Read the entire label. Use only according to label directions.

In New York State: Do not exceed one application at the recommended rate of 3.6 lbs. of product per 1,000 sq.ft. (0.25 lb. active ingredient of dithiopyr per acre).

NET WEIGHT: (16, 18, 20, 30,40, or 50) POUNDS

EPA Reg. No. 2217-859

MANUFACTURED BY:



1217 West 12th Street
Kansas City, Missouri 64101

Telephone: 1-800-821-7925



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

13/22

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or if swallowed. Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

Personal Hygiene Statements: After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-800-xxx-xxxx.</p>	

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water.

Drift or runoff from treated turf may adversely affect nontarget plants and aquatic organisms in adjacent aquatic sites. Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water when disposing of equipment washwaters.

When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD, OR FOR COMMERCIAL SEED OR FOR RESEARCH PURPOSES.

Re-entry Restrictions: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Read all precautions and directions for use carefully before applying this product.

STORAGE AND DISPOSAL

STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.

DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available
If partly filled: Call your local solid waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

In case of emergency endangering the environment involving this product, call Chemtrec day or night at (800) 424-9300.

GENERAL INFORMATION

EH-1401 Herbicide will control all your lawn's weeds in one application. EH-1401 Herbicide will kill over 200 lawn weeds commonly found in home lawns including dandelion, chickweed, thistle and clover, and will prevent crabgrass and other undesirable annual grasses. EH-1401 Herbicide also contains a long-lasting, timed release fertilizer that has all the nutrients your lawn needs for 2-3 months, depending on grass variety.

EH-1401 Herbicide is recommended for the following established turf species:

Established Cool Season Turfgrasses—Kentucky bluegrass, perennial ryegrass, fine and tall fescues.

Established Warm Season Turfgrasses—Bahia grass, bermudagrass, buffalograss, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam').

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Do not use EH-1401 Herbicide on colonial bentgrass, dichondra, carpetgrass, or 'Floritam' St. Augustinegrass.

If the lawn has been seeded recently, wait until after the third mowing before using this product. Do not seed, overseed or sprig the treated area for 10 weeks after application.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

Use of EH-1401 Herbicide on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

Do not use this product where desirable clovers are present or spread (apply) the granules onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosage for any area; be particularly careful within dripline of trees and other ornamental species.

Do not use clippings from treated areas for mulching, vegetables, or fruit trees.

Do not contaminate domestic or irrigation waters.

- WEEDS CONTROLLED -

When applied as directed, EH-1401 Herbicide will control the following weeds:

Annual Grass Weeds:

Annual Bluegrass (<i>Poa annua</i>)	Crowfootgrass	Kikuyugrass (seedling)
Annual Ryegrass	Foxtail (green and yellow)	Perennial Ryegrass
Barnyardgrass	Goosegrass	Smutgrass (seedling)
Crabgrass		


Established Broadleaf Weeds:			
Annual yellow sweetclover	Clover, white	Musk thistle	Spiny amaranth
Aster	Cockle	Mustard	Spiny cocklebur
Austrian fieldcress	Cocklebur	Narrowleaf plantain	Spiny sowthistle
Bedstraw	Common mullein	Narrowleaf vetch	Spotted catsear
Beggarticks	Creeping jenny	Nettle	Spotted spurge
Betony, Florida	Cudweed	Orange hawkweed	Spurweed
Bindweed, field	Curly dock	Oriental cocklebur	Stinging nettle
Bird vetch	Daisy, English	Oxalis	Strawberry, India mock
Bitter wintercress	Daisy fleabane	Parsley-piert	Tall nettle
Bittercress, hairy	Daisy, oxeye	Parsnip	Tall vervain
Bitterweed	Dandelion	Pearlwort	Tansy ragwort
Black-eyed Susan	Dichondra	Pennycress	Tansy mustard
Black medic	Dogfennel	Pepperweed	Tanweed
Black mustard	Dollarweed	Pigweed	Thistle
Blackseed plantain	False dandelion	Pineappleweed	Trailing crownvetch
Blessed thistle	Falseflax	Pineywoods bedstraw	Tumble mustard
Bloodflower milkweed	False sunflower	Plains tickseed	Tumble pigweed
Blue lettuce	Fiddleneck	Plantain	Velvetleaf
Blue vervain	Florida pusley	Poison ivy	Venice mallow
Bracted plantain	Frenchweed	Poison oak	Virginia buttonweed
Brassbuttons	Galinsoga	Pokeweed	Virginia creeper
Bristly oxtongue	Goathead	Poorjoe	Virginia pepperweed
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly lettuce	Western clematis
Broomweed	Gumweed	Prickly sida	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild aster
Bull thistle	Healall	Prostrate spurge	Wild buckwheat
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild carrot
Burclover	Heath aster	Puncturevine	Wild four-o'clock
Burdock	Hedge bindweed	Purslane, common	Wild garlic
Burning nettle	Hedge mustard	Ragweed	Wild geranium
Burweed	Henbit	Red sorrel	Wild lettuce
Buttercup	Hoary cress	Redroot pigweed	Wild marigold
Buttonweed	Hoary plantain	Redstem filaree	Wild mustard
Canada thistle	Hoary vervain	Rough cinquefoil	Wild onion
Carolina geranium	Horsenettle	Rough fleabane	Wild parsnip
Carpetweed	Jimsonweed	Russian pigweed	Wild radish
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild rape
Catnip	Knotweed	Scotch thistle	Wild sweet potato
Catsear	Kochia	Sheep sorrel	Wild vetch
Chickweed, common	Lambsquarters	Sheperdspurse	Woodsorrel
Chickweed, mouseear	Lespedeza	Slender plantain	Woolly croton
Chicory	Mallow	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Matchweed	Smooth dock	Woolly plantain
Clover, crimson	Mexicanweed	Smooth pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Morningglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, sweet	Mugwort	Speedwell	and other broadleaf weeds

WHEN TO APPLY

EH-1401 Herbicide can be used either in the spring or fall. EH-1401 Herbicide should be used when the grass and weeds are actively growing and the soil is moist. Do not apply under drought conditions.

In the spring, the ideal time to apply EH-1401 Herbicide is after the forsythia has bloomed or when the first dandelion flowers appear. In southern areas (see map), make a second application of EH-1401 Herbicide in 6-8 weeks for best weed and crabgrass control and for full season fertilization.

In the fall, apply EH-1401 Herbicide when the weather cools and both the grass and weeds are actively growing, generally around Labor Day.



HOW TO APPLY

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Do not apply this product onto desirable plants or shrubs as damage may result. If the product gets onto other plants immediately brush or wash it off the foliage.

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

HOW MUCH TO USE

Apply 3.6 pounds of product per 1,000 sq. ft. of turfgrass.

The maximum application rate is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

CALIBRATION INSTRUCTIONS

- For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- Check equipment frequently to verify calibration

Drop Spreader Settings:

Apply in a single pass at a rate of approximately 3.6 lbs per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working toward the center. Overlap the coverage slightly, and shut off the spreader before making hard turns.

Drop Spreader	Setting
Greenview® Drop Spred® XP1	16
Greenview® Drop Spred® III	5
Agri-Fab® Drop	4
Red Devil Drop	9
Republic Drop	5
Scotts Drop	5

Broadcast Spreader Settings:

Apply in a single pass at a rate of approximately 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. A little overlap coverage is desirable.

Broadcast Spreader	Setting
Greenview® Jet-Spred® SS1	14
Greenview® Jet-Spred® III	4½
Agri-Fab® Rotary	4
Cyclone Rotary	3½
Earthway Rotary	14
Lesco Rotary	H
Red Devil Rotary (1-20)	8
Republic Rotary	4
Sears Craftsman Rotary	4½
Scotts Speedy Green®	5

To calibrate your spreader: 1 lb of material should cover approximately 275 sq. ft.

If your spreader model is not listed, go to our website at www.lebsea.com, or call toll-free 800-233-0628, 8 A.M.—5P.M. Monday—Friday, for additional model settings.

SUGGESTED SPREADER SETTINGS:


These spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

If your spreader model does not appear below, please call toll-free 800-xxx-xxxx, 8 A.M. – 5 P.M. EDT, Monday—Friday for additional model settings.

Alternate Spreader Settings:


Suggested Spreader Settings

These spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Apply in a single pass at a rate of 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.

Broadcast Spreaders	Setting
Agri-Fab® Rotary	4
Cyclone® Rotary	3½
Earthway® Rotary	14
Greenview® Jet-Spreader® SS1	14
Lescoc® Rotary (dial A-SI/number dial)	H/14
Red Devil® Rotary (1-20)	5
Republic® Rotary	4
Scotts® EasyGreen®	23
Scotts® EdgeGuard®	4½
Scotts® SpeedyGreen® 2000, 3000	4½
Sears® Craftsman Rotary	4
Sta-Green® Rotary (1-10)	2½
True Temper® Rotary	3½
Vigoro® Rotary (1-10)	2½



Apply in a single pass at a rate of 3.6 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders	Setting
Agri-Fab® Drop	4
Greenview® Drop-Spreader® XP1	18
Red Devil® Drop (1-20)	9
Republic® Drop	5
Scotts® Drop AccuGreen (2-18)	7
Scotts® Drop PF1 (2-18)	5
Sta-Green® Drop (1-5)	3½



If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

[Note: Additional spreader models and settings may be added or substituted by the supplemental registrant provided they deliver the labeled rate of product.]

[Note: Spreader settings may be varied by the supplemental registrant as different grades of fertilizer are used, provided they deliver the labeled rate of pesticide.]

WARRANTY

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to manufacturer, and buyer assumes the risk of any such use.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on this label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of seller. All such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at sellers election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless seller is promptly notified of such loss or damage in writing. In no case shall seller be liable for consequential or incidental damages or losses.

APPENDIX

I. Advertising claims that may be presented on the container:

- ◆ **Kills Dandelion**
- ◆ **Prevents Crabgrass and Fertilizes**
- ◆ **Weed 'n Feed Plus Crabgrass Preventer**
- ◆ **Fertilize Once for Whole Season**—Feeds and greens with slow release fertilizer for up to three months.
- ◆ **Only one application needed with new nitrogen technologies.**
- ◆ **Eliminates lawn weeds**—crabgrass, dandelions, viney weeds, even clover.
- ◆ **Extended time-release feeding of nitrogen lasts into the summer.**
- ◆ Greenview guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem and simply send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.
- ◆ **This xxx bag covers up to xx sq. ft. of turfgrass (lawn):**

Bag Size, Pounds	Application Rate, Pounds/1,000 Sq.Ft.	Treated Area for This Bag Size, Square Feet
16	3.0	5,300
	3.5	4,600
	4.0	4,000
	4.5	3,600
	5.0	3,200
18	3.0	6,000
	3.5	5,100
	4.0	4,500
	4.5	4,000
	5.0	3,600
20	3.0	6,700
	3.5	5,700
	4.0	5,000
	4.5	4,400
	5.0	4,000
30	3.0	10,000
	3.5	8,600
	4.0	7,500
	4.5	6,700
	5.0	6,000
40	3.0	13,300
	3.5	11,400
	4.0	10,000
	4.5	8,900
	5.0	8,000
50	3.0	16,700
	3.5	14,300
	4.0	12,500
	4.5	11,000
	5.0	10,000

II. Additional Labeling Requirements for Fertilizers and State Registrations:

- ◆ For paper box board containers, the following statement may appear: **This container is constructed from 85% recycled box board.**
- ◆ If the fertilizer guarantee contains iron, this following statement must appear: *This product contains secondary nutrient iron. Iron may stain concrete and should not be applied on dry or water dampened concrete and should be removed from these area promptly after application by sweeping or blowing. Do not wash off with water.*
- ◆ If the recommended use rate applies less than 1 pound nitrogen per 1,000 sq. ft. per application, the following statement must appear: *Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq. ft.*
- ◆ For products distributed in the State of Washington, the following statement must appear: *Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.xxx.com> .*
- ◆ Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. nitrogen per 1,000 sq. ft.

III. Addition of fertilizer grade, guaranteed analysis and advertising claims:

GUARANTEED FERTILIZER ANALYSIS—24-2-8	
Total Nitrogen (N).....	24%
2.0% Ammoniacal Nitrogen	
8.5% Water Insoluble Nitrogen*	
4.3% Urea Nitrogen	
9.2% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ O ₅).....	2%
Soluble Potash (K ₂ O).....	8%
Total Sulfur (S).....	2.6%
2.6% Combined Sulfur (S)	
Nutrient Sources: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Muriate of Potash.	
Chlorine (Cl) not more than.....	7.0%
*17.7% Slowly Available Nitrogen from Methylene Ureas.	F699
Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-lebsea.com .	

WHAT IS SLOW RELEASE NITROGEN?	
Greenview Fairway Formula Fertilizers utilize a unique, slow release nitrogen source developed for professionals.	
METH-EX[®]	Provides slow, continuous feeding for up to 12 weeks. Initially some nitrogen is released by water to provide early spring green-up. As soil temperatures warm, nitrogen is gradually released through microbial action that minimizes surge growth.
17.7% Slowly Available Nitrogen/24% Total Nitrogen = 7/10 of the Nitrogen is Slow Release	
0.9 lb. Nitrogen/1,000 sq. ft.	

- One Product Kills Broadleaf Weeds and Prevents Crabgrass
- 4 Weed Control Products in 1
- Replaces Steps 1 and 2 of a 4 Step Program
- Fertilizes all season—longer than most other leading brands
- In spring apply once when weeds are growing (see back panel on when to apply)



- Good Housekeeping Seal (see attached front bag copy)

22/22

- Stops More Weeds In One Application!
- Apply once in Spring when weeds are growing