

2217-864

12/01/2008

1/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

1 DEC 2008

James L. Kunstman  
PBI/Gordon Corp.  
1217 West 12<sup>th</sup> Street  
P.O. Box 014090  
Kansas City, MO 64101

Dear Dr. Kunstman :

Subject: Label Amendment for RED: 2,4-D, Dicamba and MCPP-p  
1381 EC Residential  
EPA Registration Number 2217-864


The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Per the acute toxicity review, the "IF ON SKIN" heading in the first aid statements must be revised to read "IF ON SKIN OR CLOTHING."
2. Per the acute toxicity review, the text in **bold type** below must be added Hazards to Humans and Domestic Animals:  
  
"CAUTION  
Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. **Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber, or viton.** Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."
3. Per the 2,4-D, Dicamba and MCPP-p combined label table, the first 2 sentences of the Environmental Hazard section must be revised to read:  
  
"This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark."
4. Per the product chemistry review, the heading "Storage" in the Storage and Disposal section should be changed to "Pesticide Storage."

5. Per the combined label table, the following text must be added to the label:
- “Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated areas to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.”
- “Do not apply as a fine mist because of potential for injury to desirable plants.”
6. Per the combined label table, the text “Wait at least 30 days between applications” must be added to the label.
7. Change sentence under Limited Warranty and Disclaimer to read “**To the extent consistent with applicable law**, the manufacturer neither makes nor intends any other express or implied warranties including any warranty of merchantability or fitness for a particular purpose, which are expressly disclaimed.”
8. Please note: Final product reregistration cannot be considered until after all active ingredients in this product are eligible for reregistration.
9. Add the following statements to the labeling: Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA Web Site: <http://www.epa.gov/espp>.

One copy of the label stamped “Accepted with Comments” is enclosed for your records. Please submit one copy of the final printed label before the product is released for shipment. If you have any questions, please contact Tracy White by phone at (703) 308-0042 or via email at [white.tracy@epa.gov](mailto:white.tracy@epa.gov).

Sincerely,



Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure

3/10

# 1381 EC RESIDENTIAL

## EPA Reg. No. 2217-864

### ACTIVE INGREDIENT:

2,4-D, 2-ethylhexyl ester .....	28.57%
Mecoprop-p acid .....	5.88%
Dicamba acid .....	1.71%
Carfentrazone-ethyl. ....	0.62%

INERT INGREDIENTS:..... 63.22%

TOTAL..... 100.00%

2,4-dichlorophenoxyacetic acid equivalent 18.95%.

Isomer Specific By AOAC Methods.

Contains petroleum distillates.

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**ACCEPTED  
with COMMENTS**

**In EPA Letter Dated:**

**1 DEC 2008**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

2217-864

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### **Personal Hygiene Statements**

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

<b>First Aid</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give <b>any</b> liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

<b>If on skin:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact 1-877-800-5556 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillates - vomiting may cause aspiration pneumonia.	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **General Precautions and Restrictions:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### **PRODUCT DESCRIPTION:**

This Trimec® product is a combination of four proven weed killers: 2,4-D, mecoprop-p, dicamba and carfentrazone-ethyl. Postemergent applications of this product control a wide range of broadleaf lawn weeds in residential lawns including dandelion, chickweed, spurge, pennywort (dollarweed), knotweed, plantain, and clover.

This product offers these advantages:

- Rapid and effective weed control for common and troublesome weed species in turfgrass
- Superior cool weather performance.
- Injury symptoms on broadleaf weeds can be noticed within hours of the application and plant death can occur within 7 to 14 days.

### **WHERE TO USE:**

This product is intended for use on most ornamental lawns including:

Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass Perennial ryegrass Tall fescue Red or fine leaf fescues Colonial bentgrass Mixtures of cool season species	Common bermudagrass Hybrid bermudagrass Zoysiagrass

#### **Turfgrass tolerance:**

The turfgrass tolerance to 1381 EC Residential may vary and temporary turfgrass yellowing may occur on certain varieties (hybrids) of hybrid bermudagrass.

1381 EC Residential may injure certain turfgrass species. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 85°F.

#### **Prohibitions:**

- Do not use this product on bahiagrass, buffalograss, carpetgrass, centipedegrass, dichondra, St. Augustinegrass, or lawns containing desirable clovers.
- Do not spray roots of trees and ornamentals.
- Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result.
- Do not exceed specified dosages for any area. Be particularly careful within dripline of trees and other ornamental species.

#### **Limitations on 2,4-D applications to residential turfgrass**

The maximum application rate is 1.5 fl. oz. of product per 1,000 sq.ft. per application (0.8 lbs 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. The maximum seasonal rate is 3.0 fl. oz. of product per 1,000 sq.ft. (1.6 lbs 2,4-D acid equivalent per acre), excluding spot treatments.

#### **WHEN TO APPLY:**

Spray when broadleaf weeds are young and actively growing for best results.

1381 EC Residential may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments.

Broadcast (entire areas) applications or follow-up applications at 2 to 6 week intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are susceptible.

#### **CULTURAL TIPS FOR IMPROVED CONTROL:**

##### **Irrigation:**

- Do not apply this product immediately before rainfall or irrigation. Rain or watering within 4 hours after treatment may reduce effectiveness.

##### **Mowing:**

- Delay mowing 1 to 2 days before and after the application of this product.

##### **Reseeding Interval:**

- Treated areas may be reseeded 2 weeks after application.

### THREE EASY STEPS TO MIX AND APPLY:

1. Determine (Measure) the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is presented as fl.oz. (tablespoons) per 1,000 sq.ft. of lawn.
2. Use hand-operated sprayers (backpack sprayers, compression (pump-up) sprayers, knapsack sprayers or tank type) or hose end sprayers. Carefully measure and mix the proper amount of product with water. See below for instructions for the various types of sprayers.
3. Spray evenly over the lawn.

### HOW MUCH TO USE:

#### 1. BROADCAST TREATMENTS FOR COOL SEASON TURFGRASS:

Use rates of 1381 EC Residential on cool season turfgrass as broadcast treatments are presented in Table 1.

TABLE 1. RATE RECOMMENDATIONS FOR COOL SEASON TURFGRASS.					
Species	Amount of Product Fl.oz./1,000 sq.ft.	Approximate Treated Area, sq. ft.			
		8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container	20 fl. oz. sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container
Kentucky bluegrass	1.4	5,700 sq. ft.	11,400 sq. ft.	14,300 sq. ft.	22,800 sq. ft.
Perennial ryegrass	1.4	5,700 sq. ft.	11,400 sq. ft.	14,300 sq. ft.	22,800 sq. ft.
Tall fescue	1.4	5,700 sq. ft.	11,400 sq. ft.	14,300 sq. ft.	22,800 sq. ft.
Red or fine leaf fescues	1.4	5,700 sq. ft.	11,600 sq. ft.	14,300 sq. ft.	23,200 sq. ft.
Colonial bentgrass	1.4	5,700 sq. ft.	11,400 sq. ft.	14,300 sq. ft.	22,800 sq. ft.
Mixtures of cool season species	1.4	5,700 sq. ft.	11,400 sq. ft.	14,300 sq. ft.	22,800 sq. ft.

### HOW TO APPLY:

#### FOR ORTHO HOSE-END SPRAYERS FOR COOL SEASON TURFGRASS

- For Ortho® Dial'N Spray Hose-End Sprayer or Gilmour® Dial-a-Mix® Lawn and Garden Sprayer:

**Set the dial**—Set dial on 3 tsp. (1 Tbs.) per gallon.

**Fill**—Determine [Measure] the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is 1.4 fl. oz. per 1,000 sq. ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area.

**Spray**—Spray evenly over the lawn.

**Example:** Measure 4.0 fl.oz. (8 Tbs.) into sprayer jar. Do not add water. Set dial on 3 teaspoons (tsp.) and spray uniformly on 3,000 sq.ft. of lawn.

- For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Ratio of 80:1:

For 1.4 fl.oz. of product per 1,000 sq.ft.: Measure 9 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.

### SPOT TREATMENTS FOR COOL SEASON TURFGRASS.

#### HAND OPERATED SPRAYERS:

For cool season turfgrass, mix 1.0 fl.oz. of 1381 EC Residential per 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

Spray preparation for spot treatments on cool season turfgrass			
Spray volume, gals. Of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	1.0 fl.oz.	2.0 tbs.	1,000 sq.ft.
2 gallons	2.0 fl.oz.	4.0 tbs.	2,000 sq.ft.
3 gallons	3.0 fl.oz.	6.0 tbs.	3,000 sq.ft.
Equal measures: 1.0 fl.oz. = 2 tablespoons.			

## 2. BROADCAST TREATMENTS FOR WARM SEASON TURFGRASS

Use rates of 1381 EC Residential on warm season turfgrass as broadcast treatments are presented in Table 2.

TABLE 2. RATE RECOMMENDATIONS FOR WARM SEASON TURFGRASS.					
Species	Amount of Product Fl.oz./1,000 sq.ft.	Approximate Treated Area, sq. ft.			
		8.0 fl.oz. sprayer jar or retail container	1.0 pt. (16 fl.oz.) sprayer jar or retail container	20 fl. oz. sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container
Common bermudagrass	1.5	5,400 sq. ft.	10,900 sq. ft.	13,300 sq. ft.	21,800 sq. ft.
Hybrid bermudagrass	1.1	7,300 sq. ft.	14,500 sq. ft.	18,200 sq. ft.	29,100 sq. ft.
Zoysiagrass	1.1	7,300 sq. ft.	14,500 sq. ft.	18,200 sq. ft.	29,100 sq. ft.

### HOW TO APPLY:

#### FOR ORTHO HOSE-END SPRAYERS FOR WARM SEASON TURFGRASS

For common bermudagrass:

- For Ortho® Dial'N Spray Hose-End Sprayer or Gilmour® Dial-a-Mix® Lawn and Garden Sprayer:

**Set the dial**—Set dial on 3 tsp. (1 Tbs.) per gallon.

**Fill**—Determine [Measure] the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is 1.5 fl. oz. per 1,000 sq. ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area.

**Spray**—Spray evenly over the lawn.

**Example:** Measure 4.4 fl.oz. (9 Tbs.) into sprayer jar. Do not add water. Set dial on 3 teaspoons (tsp.) and spray uniformly on 3,000 sq.ft. of lawn.

- For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Ratio of 80:1:

For 1.5 fl.oz. of product per 1,000 sq.ft.: Measure 10.0 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.

For hybrid bermudagrass and zoysiagrass:

- For Ortho® Dial'N Spray Hose-End Sprayer or Gilmour® Dial-a-Mix® Lawn and Garden Sprayer:

**Set the dial**—Set dial on 2 tsp. per gallon.

**Fill**—Determine [Measure] the lawn area to be sprayed. Calculate square feet by multiplying length by width. The application rate for this product is 1.1 fl. oz. per 1,000 sq. ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area.

**Spray**—Spray evenly over the lawn.

**Example:** Measure 3.3 fl.oz. (6.6 Tbs.) into sprayer jar. Do not add water. Set dial on 2 teaspoons (tsp.) and spray uniformly on 3,000 sq.ft. of lawn.

- For Gilmour® 20-Gallon Lawn Sprayer, Model 390B, Sprayer Jar with 32 fl.oz. and Dilution Ratio of 80:1:

For 1.1 fl.oz. of product per 1,000 sq.ft.: Measure 7.3 fl.oz. of product into sprayer jar. Add water or fill with water to the 32 fl.oz. line of the sprayer jar. Spray entire contents uniformly on 6,700 sq.ft. of lawn.

## SPOT TREATMENTS FOR WARM SEASON TURFGRASS:

### HAND OPERATED SPRAYERS.

For warm season turfgrass, mix 0.75 to 1.0 fl. oz. of 1381 EC Residential per 1.0 gallon of water for treatment of approximately 1,000 sq. ft. of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

Spray preparation for spot treatments on warm season turfgrass			
Spray volume, gals. of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	0.75 to 1.0 fl.oz.	1.5 to 2.0 tbs.	1,000 sq.ft.
2 gallons	1.5 to 2.0 fl.oz.	3.0 to 4.0 tbs.	2,000 sq.ft.
3 gallons	2.25 to 3.0 fl.oz.	4.5 to 6.0 tbs.	3,000 sq.ft.
Equal measures: 1.0 fl.oz. = 2 tablespoons.			

### BROADLEAF WEEDS CONTROLLED:

1381 EC Residential will control or suppress the following broadleaf weeds and will control or suppress other broadleaf weeds that are susceptible to 2,4-D. A partial list is presented below.

BROADLEAF WEEDS			
Aster, white heath & white prairie	English daisy	Mallow, common	Shepherdspurse
Bedstraw	False dandelion (*spotted catsear & common catsear)	Matchweed	Spotted spurge
Beggarweed, creeping	Field bindweed	Mouseeear chickweed	Spurge
Bindweed	(*morningglory & creeping jenny)	Old world diamond flower	Thistle
Black medic	Field oxeye-daisy	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Veronica (*corn speedwell)
Broadleaf plantain	(*creeping oxeye)	Parsley-piert	Virginia buttonweed
Buckhorn plantain	Filaree, whitestem & redstem	Pennsylvania smartweed	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Bull thistle	Florida betony	Pennywort (*dollarweed)	Wild carrot
Burclover	Florida pusley	Pepperweed	Wild garlic
Burdock, common	Ground ivy	Pigweed	Wild geranium
Buttercup, creeping	Groundsel	Pineappleweed	Wild lettuce
Carpetweed	Hawkweed	Plantain	Wild mustard
Chickweed, common	Healall	Poison ivy	Wild onion
Chicory	Henbit	Poison oak	Wild strawberry
Cinquefoil	Innocence (Blue-eyed Mary)	Prostrate knotweed (*knotweed)	Wild violet
Clover	Lambsquarters	Puncturevine	Yarrow
Compassplant	Lawn burweed	Purple cudweed	Yellow rocket
Curly dock	Lespedeza, common	Purslane	and many more broadleaf weeds
Dandelion		Ragweed	
Dayflower		Redweed	
Deadnettle		Red sorrel (*sheep sorrel)	
Dock			
Dogfennel			
*Synonyms			



## **STORAGE AND DISPOSAL**

**STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

## **LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

## **Appendix**

### **1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

### **2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- Rainfast in as little as 3 hours
- Rainfast [Rainproof] in 3 hours
- Visible results in just 24 hours
- Clover control in as little as one week
- Proven cool-weather performance
- Consistently fast control on tough weeds like clover, plantain, ground ivy and spurge [various other listed weeds].
- [This product] Smokes weeds, fast
- Dead weeds fast!
- Fast results

10  
10

- Contains the newest chemistry (formula) for weed control
- Contains Carfentrazone
- Contains new active ingredient
- From the makers of Trimec® herbicides.
- For information call XXX-XXX-XXXX [contact [www.xxx.com](http://www.xxx.com)]
- DEAD WEEDS FAST!

<b>GIVE A DANDELION AN INCH AS THEY'LL TAKE A YARD.</b>	
WHAT IT'S FOR:	<ul style="list-style-type: none"> <li>• Use in home lawns</li> <li>• Rapid and effective control of common and troublesome broadleaf weeds</li> </ul>
WHERE TO USE:	<ul style="list-style-type: none"> <li>• Kentucky bluegrass</li> <li>• Perennial ryegrass</li> <li>• Fescues</li> <li>• Bermudagrass</li> <li>• Zoysiagrass</li> </ul>
AMOUNT TO USE:	<ul style="list-style-type: none"> <li>• 1.1 to 1.5 fl. oz. per 1,000 sq. ft. This container will treat up to xx,xxx sq. ft. of lawn.</li> <li>• Refer to complete instructions in Directions for Use section of this booklet.</li> </ul>
QUESTIONS:	<ul style="list-style-type: none"> <li>• For frequently asked questions, visit our website at <a href="http://www.pbigordon.com">www.pbigordon.com</a></li> </ul>

- SPEED-ZONE® is a registered trademark of PBI/Gordon Corporation
- TRIMEC® is a registered trademark of PBI/Gordon Corporation
- ORTHO® Dial'N Spray Hose-End Sprayer is a registered trademark of Monsanto Company
- GILMOUR® Dial-A-Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.

**3. Alternative to the previously approved warranty if the small size of the container labels makes it impractical to present the entire warranty.**

- **NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

## **DOCUMENT CONTROL INFORMATION**

- 1. Unique Label Identifier:** 002217-00864.20071105
- 2. Previous Version:** AP110104
- 3. Reason for Issue:** Various changes pertaining to re-registration. Re-format label.