

2217-865

12-26-2007

1/14



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

2217-865

Date of Issuance:

12-26-07

Term of Issuance:

Name of Pesticide Product:

EH 1404 Herbicide

Name and Address of Registrant (include ZIP Code):

PBI Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.


On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

1. The concentration of the active ingredient Carfentrazone-ethyl must be changed to "0.0023%".
2. To the Hazards to Humans and Domestic Animals add "and before eating, drinking, chewing gum, using tobacco, or using the toilet" after "Wash thoroughly with soap and water after handling".
3. To the label add "Do not apply more than 1.5 lb ai (MCPA) / acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

2/14

4. To the label add under the ingredient statement "This product contains _____ lbs. ai/gallon".	
Signature of Approving Official:  Erik Kraft Acting Team Leader 23 Herbicide Branch, Registration Division (7505P)	Date: 12-24-07

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records.

3/14

Sublabel 1	Gordon's SpeedZone Ready-to-Use Lawn Weed Killer
	Front panel, Precautionary Statements and Directions for Use consistent with an Agency letter dated September 16, 2005.

EH 1404 HERBICIDE

For Home Lawns Only

ACTIVE INGREDIENTS:

MCPA, 2-ethylhexyl ester	0.337%
Mecoprop-p acid	0.066%
Dicamba acid.....	0.018%
Carfentrazone-ethyl	0.002%
INERT INGREDIENTS:	99.577%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

2-methyl-4-chlorophenoxyacetic acid equivalent, 0.216%
 TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

See side panels for additional Precautionary Statements

NET CONTENTS: 24, 32, 64 fl. oz. and 1 Gallon

AP-----
 EPA Reg. No. 2217-865
 EPA Est. No. 2217-KS-1
 MANUFACTURED BY:



Telephone: 1-800-821-7925

ACCEPTED
 with COMMENTS
 in EPA Letter Dated

12-26-07

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.

2217-865



4/1/14

**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS:

This pesticide may be toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.

PRODUCT FACTS:

EH 1404 Herbicide contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

EH 1404 Herbicide offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in turfgrass, e.g. spurge, pennywort (dollarweed), dandelion, and white clover.
- Injury to weeds can be noticed within hours of the application and plant death can occur within 7 - 14 days.

5/14

WHERE TO USE:

On Lawns: Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as:

- Kentucky bluegrass
- Perennial ryegrass
- Tall fescue
- Red or fine leaf fescues
- Mixtures of cool season species
- Common bermudagrass
- Hybrid bermudagrass
- St. Augustinegrass (excluding 'Floritam')
- Zoysiagrass

Prohibitions:

- Do not apply this product to bentgrass, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not apply to 'Floritam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.

APPLICATION SCHEDULES (WHEN TO APPLY):

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 2 to 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant bermudagrass, zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

HOW TO USE:

For Containers with Trigger Sprayers: EH 1404 Herbicide is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

For Refill Containers: EH 1404 Herbicide is ready-to-use and is applied directly from the container. Remove the cap and attach the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

6/14

Trigger Sprayer Instruction for Operation and Care:

1. Remove sprayer from bag.
2. Shake the container well.
3. Unscrew cap from the bottle, and remove the foil seal.
4. Making sure that the tube is properly inserted through hole in cap, insert tube into bottle and screw on cap.
5. Open spray nozzle approximately one full turn.
6. TO PRIME—With sprayer directed at the target area, operate trigger until spray begins (approximately 10 to 15 times).
7. Adjust the spray nozzle for desired spray pattern
8. To store, turn nozzle to "OFF" position and store with trigger head above level of liquid in container to prevent leakage.

BROADLEAF WEEDS CONTROLLED:

EH 1404 Herbicide will control or suppress the following broadleaf weeds:

- BROADLEAF WEEDS -		
Aster, white heath & white prairie	Field oxeye-daisy (*creeping oxeye)	Prostrate knotweed (*knotweed)
Bedstraw	Filaree, whitestem & redstem	Puncturevine
Beggarweed, creeping	Florida pusley	Purple cudweed
Bindweed	Ground ivy	Purslane
Black medic	Groundsel	Ragweed
Broadleaf plantain	Hawkweed	Redweed
Buckhorn plantain	Healall	Red sorrel (*sheep sorrel)
Bull thistle	Henbit	Shepherdspurse
Burclover	Innocence (Blue-eyed Mary)	Spotted spurge
Burdock, common	Lambsquarters	Spurge, prostrate
Buttercup, creeping	Lawn burweed	Thistle
Carpetweed	Lespedeza, common	Veronica (*corn speedwell)
Chickweed, common	Mallow, common	Virginia buttonweed
Chicory	Matchweed	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Cinquefoil	Mouseear chickweed	Wild carrot
Clover	Old world diamond flower	Wild garlic
Compassplant	<i>Oxalis</i> (*yellow woodsorrel & creeping woodsorrel)	Wild geranium
Curly dock	Parsley-piert	Wild lettuce
Dandelion	Pennsylvania smartweed	Wild mustard
Dayflower	Pennywort (*dollarweed)	Wild onion
Deadnettle	Pepperweed	Wild strawberry
Dock	Pigweed	Wild violet
Dogfennel	Pineappleweed	Yarrow
English daisy	Plantain	Yellow rocket
False dandelion (*spotted catsear & common catsear)	Poison ivy	
Field bindweed (*mornningglory & creeping jenny)	Poison oak	

*Synonyms

7/14

STORAGE & DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in an area inaccessible to children and pets.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency (or 1-800-XXX-XXXX) for disposal instruction. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

---END of Sublabel 1---

8/14

Sublabel 2	Text intended for supplemental registrants: Front panel, Precautionary Statements and Directions for Use consistent with our notification dated May 25, 2005.
-------------------	---

EH 1404 HERBICIDE

For Home Lawns Only

ACTIVE INGREDIENTS:

MCPA, 2-ethylhexyl ester	0.337%
Mecoprop-p acid	0.066%
Dicamba acid.....	0.018%
Carfentrazone-ethyl	0.002%

INERT INGREDIENTS:	<u>99.577%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 2-methyl-4-chlorophenoxyacetic acid equivalent, 0.216%
- (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent. 0.066%
- 3,6-dichloro-o-anisic acid equivalent, 0.018%
- Ethyl a,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate, 0.002%

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

See side panels for additional Precautionary Statements

NET CONTENTS: 24, 32, 64 fl. oz. and 1 Gallon

AP.....
EPA Reg. No. 2217-865
EPA Est. No. 2217-KS-1
MANUFACTURED BY:

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
Telephone: 1-800-821-7925



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

9/14

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS:

This pesticide may be toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.

PRODUCT FACTS:

EH 1404 Herbicide contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

EH 1404 Herbicide offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in turfgrass, e.g. spurge, pennywort (dollarweed), dandelion, and white clover.
- Injury to weeds can be noticed within hours of the application and plant death can occur within 7 - 14 days.

WHERE TO USE:

On Lawns: Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as:

Kentucky bluegrass
Perennial ryegrass
Tall fescue
Red or fine leaf fescues
Mixtures of cool season species
Common bermudagrass
Hybrid bermudagrass
St. Augustinegrass (excluding 'Floritam')
Zoysiagrass

10/14

Prohibitions:

- Do not apply this product to bentgrass, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not apply to 'Floritam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.

APPLICATION SCHEDULES (WHEN TO APPLY):

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 2 to 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant bermudagrass, zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

HOW TO USE:

For Containers with Trigger Sprayers: EH 1404 Herbicide is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

For Refill Containers: EH 1404 Herbicide is ready-to-use and is applied directly from the container. Remove the cap and attach the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

Trigger Sprayer Instruction for Operation and Care:

1. Remove sprayer from bag.
2. Shake the container well.
3. Unscrew cap from the bottle, and remove the foil seal.
4. Making sure that the tube is properly inserted through hole in cap, insert tube into bottle and screw on cap.
5. Open spray nozzle approximately one full turn.
6. TO PRIME—With sprayer directed at the target area, operate trigger until spray begins (approximately 10 to 15 times).
7. Adjust the spray nozzle for desired spray pattern
8. To store, turn nozzle to "OFF" position and store with trigger head above level of liquid in container to prevent leakage.

11/14

BROADLEAF WEEDS CONTROLLED:

EH 1404 Herbicide will control or suppress the following broadleaf weeds:

- BROADLEAF WEEDS -		
Aster, white heath & white prairie	Field oxeye-daisy (*creeping oxeye)	Prostrate knotweed (*knotweed)
Bedstraw	Filaree, whitestem & redstem	Puncturevine
Beggarweed, creeping	Florida pusley	Purple cudweed
Bindweed	Ground ivy	Purslane
Black medic	Groundsel	Ragweed
Broadleaf plantain	Hawkweed	Redweed
Buckhorn plantain	Healall	Red sorrel (*sheep sorrel)
Bull thistle	Henbit	Shepherdspurse
Burclover	Innocence (Blue-eyed Mary)	Spotted spurge
Burdock, common	Lambsquarters	Spurge, prostrate
Buttercup, creeping	Lawn burweed	Thistle
Carpetweed	Lespedeza, common	Veronica (*corn speedwell)
Chickweed, common	Mallow, common	Virginia buttonweed
Chicory	Matchweed	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Cinquefoil	Mouseear chickweed	Wild carrot
Clover	Old world diamond flower	Wild garlic
Compassplant	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Wild geranium
Curly dock	Parsley-piert	Wild lettuce
Dandelion	Pennsylvania smartweed	Wild mustard
Dayflower	Pennywort (*dollarweed)	Wild onion
Deadnettle	Pepperweed	Wild strawberry
Dock	Pigweed	Wild violet
Dogfennel	Pineappleweed	Yarrow
English daisy	Plantain	Yellow rocket
False dandelion (*spotted catsear & common catsear)	Poison ivy	
Field bindweed (*morninggloory & creeping jenny)	Poison oak	

*Synonyms

STORAGE & DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in an area inaccessible to children and pets.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency (or 1-800-XXX-XXXX) for disposal instruction. Never place unused product down any indoor or outdoor drain.

12/14

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

18/14

APPENDIX to Sublabel 1 and Sublabel 2

I. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- ◆ Rainfast in as little as 3 hours
- ◆ Rainfast [Rainproof] in 3 hours
- ◆ Visible results in just 24 hours
- ◆ Clover control in as little as one week
- ◆ Proven cool-weather performance
- ◆ Consistently fast control on tough weeds like clover, plantain, ground ivy and spurge [various other listed weeds].
- ◆ For information call XXX-XXX-XXXX [contact www.xxx-xxxx.com]

WHAT IT'S FOR:	<ul style="list-style-type: none"> • Use on home lawns • Rapid and effective control of common and troublesome broadleaf weeds
WHERE TO USE:	<ul style="list-style-type: none"> • Kentucky bluegrass • Perennial ryegrass • Fescues • Bermudagrass • Zoysiagrass
AMOUNT TO USE:	<ul style="list-style-type: none"> • Ready-to-use and is applied directly from the container. • Refer to complete instructions in Directions for Use section of this booklet
QUESTIONS:	<ul style="list-style-type: none"> • For frequently asked questions, visit our website at consumer.pbigordon.com

- **DEAD WEEDS...FAST**
- FOR HOME LAWNS ONLY
- consumer.pbigordon.com
- Performs in both warm and cool weather
- See results in hours
- Kills 200+ Broadleaf Weeds
- Won't harm lawn grass when used as directed
- For Northern and Southern grasses
- EH 1404 Herbicide will control or suppress the following broadleaf weeds and will control or suppress other broadleaf weeds that are susceptible to MCPA.
- Also, adding expanded weed list as presented on following page:

14/14

- BROADLEAF WEEDS -

Annual yellow sweetclover	Cockle	Musk thistle	Spiny amaranth
Aster	Cocklebur	Mustard	Spiny cocklebur
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny sowthistle
Bedstraw	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Beggarticks	Cudweed	Nettle	Spotted spurge
Betony, Florida	Curly dock	Orange hawkweed	Spurweed
Bindweed, field	Daisy, English	Oriental cocklebur	Stinging nettle
Bird vetch	Daisy fleabane	Oxalis	Strawberry, India mock
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall nettle
Bittercress, hairy	Dandelion	Parsnip	Tall vervain
Bitterweed	Dichondra	Pearlwort	Tansy ragwort
Black-eyed Susan	Dogbane	Pennycress	Tansy mustard
Black medic	Dogfennel	Pennywort	Tanweed
Black mustard	Dollarweed	Peppergrass	Thistle
Blackseed plantain	Elderberry	Pepperweed	Trailing crownvetch
Blessed thistle	False dandelion	Pigweed	Tumble mustard
Bloodflower milkweed	Falseflax	Pineywoods bedstraw	Tumble pigweed
Blue lettuce	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue vervain	Fiddleneck	Plantain	Venice mallow
Bracted plantain	Florida pusley	Poison ivy	Virginia buttonweed
Brassbuttons	Frenchweed	Poison oak	Virginia creeper
Bristly oxtongue	Galinsoga	Pokeweed	Virginia pepperweed
Broadleaf dock	Goathead	Poorjoe	Water pennywort
Broadleaf plantain	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broomweed	Ground ivy	Prickly lettuce	Western clematis
Buckhorn	Gumweed	Prickly sida	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild aster
Bull thistle	Healall	Prostrate spurge	Wild buckwheat
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild carrot
Burclover	Heathaster	Puncturevine	Wild four-o'clock
Burdock	Hedge bindweed	Purslane, common	Wild garlic
Burning nettle	Hedge mustard	Ragweed	Wild geranium
Burweed	Hemp	Red sorrel	Wild lettuce
Buttercup	Henbit	Redroot pigweed	Wild marigold
Buttonweed	Hoary cress	Redstem filaree	Wild mustard
Canada thistle	Hoary plantain	Rough cinquefoil	Wild onion
Carolina geranium	Hoary vervain	Rough fleabane	Wild parsnip
Carpetweed	Horsenettle	Russian pigweed	Wild radish
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild rape
Catnip	Knawel	Scarlet pimpernel	Wild strawberry
Catsear	Knotweed	Scotch thistle	Wild sweet potato
Chickweed, common	Kochia	Sheep sorrel	Wild vetch
Chickweed, mouseear	Lambsquarters	Shepherdspurse	Woodsorrel
Chicory	Lespedeza	Slender plantain	Woolly croton
Cinquefoil	Mallow	Smallflower galinsoga	Woolly morningglory
Clover, crimson	Matchweed	Smooth dock	Woolly plantain
Clover, hop	Mexicanweed	Smooth pigweed	Wormseed
Clover, red	Milk vetch	Sorrel	Yarrow
Clover, strawberry	Morningglory	Sowthistle	Yellow rocket
Clover, sweet	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, white	Mugwort	Speedwell	and other broadleaf weeds