



1217 WEST 12TH STREET • P.O. BOX 014090
KANSAS CITY, MISSOURI 64101-0090
816-421-4070 • 1-800-821-7925
FAX: 816-474-0462

August 8, 2006

Ms. Joanne I. Miller (PM-23)
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
Crystal Mall #2, Room 266A
1801 South Bell Street
Arlington, VA 22202

NOTIFICATION

AUG 31 2006

Dear Ms. Miller:

**Subject: Labeling notification for EH-1406 Herbicide (EPA Reg. No. 2217-867)
per Pesticide Registration Notice (PRN) 98-10.**

I. LABELING NOTIFICATION:

A notification was requested on April 11, 2006 regarding the use restrictions in New York. Specifically, we asked to add the following state restrictions to the draft labeling:

- Not for sale, distribution or use in Nassau or Suffolk Counties in New York State.
- New York: Only one application per year of this product is allowed. This product is not allowed to be sold or used in Nassau or Suffolk Counties.

Two (2) copies of the pdf file intended for the printing of 2.5 gallon container labels are enclosed. Please pin-punch one copy label and return in the self-addressed envelope to satisfy state registration requirements.

II. Certification:

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

2/17


III. Enclosures:

The administrative portion of this response includes the following:

1. Application for Pesticide Amendment (EPA Form 8570-1).
2. Two (2) copies of the pdf used for printing container labels.

Please contact me at 816-460-6292 or via e-mail at jkunstman@pbigordon.com if you have questions concerning this submission.

Sincerely,



James L. Kunstman, Ph.D.
Director of Regulatory Services

3/17

NOTIFICATION

AUG 31 2006



SUIRGE®

BROADLEAF HERBICIDE FOR TURF

ACTIVE INGREDIENTS:

Sulfentrazone	0.67%
2,4-D, dimethylamine salt	18.79%
Mecoprop-p, dimethylamine salt	6.80%
Dicamba, dimethylamine salt	3.02%



INERT INGREDIENTS:

70.72%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 0.06 lb. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.67%.
- 1.40 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 15.66%.
- 0.50 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 5.62%.
- 0.22 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.52%.

Isomer Specific by AOAC Method.

**KEEP OUT OF REACH OF CHILDREN
DANGER – PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

See attached booklet for additional Precautionary Statements and First Aid.



0 70251 65526 7

For use by individuals/firms licensed or registered by the State to apply pesticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your State prior to use of this product. *Not for sale, distribution, or use in Nassau or Suffolk Counties in NY State.*


655/8-2006
EPA REG. NO. 2217-867
EPA EST. NO. 2217-KS-1

Manufactured By
G pbi/gordon
corporation
An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
www.pbigordon.com


**From the makers of
TRIMEC Herbicide**

NET CONTENTS: 2.5 U.S. GALLONS

4/17




SPURGE




KNOTWEED




OXALIS*



CLOVER




DANDELION



YELLOW NUTSEDGE**

*Yellow woodsorrel

**Suppression only



Warm-weather weed control - FAST!

For use on: Bluegrass, Fescues, Perennial Ryegrass, Bentgrass, Bermudagrass, Zoysiagrass, Bahiagrass and Buffalograss

- *Fast visual response*
- *Economical - lower cost water-based formulation*
- *Easy cleanup*
- *Warm-weather weed control option*
- *Suitable for applications up to 90°F*
- *Foliar absorption*
- *Reduces call backs*
- *Low odor*
- *Designed for residential and commercial turf applications and many other sites*

Not for sale, distribution, or use in Nassau or Suffolk Counties in NY State.

Manufactured By
G pbi / GORDON CORPORATION

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordcn.com

655/8-2006 AP041106
EPA REG. NO. 2217-867
EPA EST. NO. 2217-KS-1



SURGE

BROADLEAF HERBICIDE FOR TURF



For use by individuals/firms licensed or registered by the State to apply pesticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your State prior to use of this product. Not for sale, distribution, or use in Nassau or Suffolk Counties in NY State.

ACTIVE INGREDIENTS:

Sulfentrazone.	0.67%
2,4-D, dimethylamine salt	18.79%
Mecoprop-p, dimethylamine salt	6.80%
Dicamba, dimethylamine salt.	3.02%

INERT INGREDIENTS: 70.72%
TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 0.06 lb. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.67%.
 - 1.40 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 15.66%.
 - 0.50 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 5.62%.
 - 0.22 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.52%.
- Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN DANGER – PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

See below for additional Precautionary Statements and First Aid.



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if absorbed through skin or if swallowed. Avoid contact with skin, eyes or clothing.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

6/17

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Containers of 5 or more gallons: Do not open-pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

<p>User Safety Recommendations:</p> <p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

(cont. on next column)

AGRICULTURAL USE REQUIREMENTS (cont.)

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

PRODUCT DESCRIPTION

SURGE® Broadleaf Herbicide For Turf contains four active ingredients including sulfentrazone that broaden the spectrum of weed control. These combined herbicides provide limited residual activity at recommended use rates. Sulfentrazone is in the aryl triazolinone family and inhibits protoporphyrinogen oxidase (Protox), a pivotal enzyme in chlorophyll production. Without this key enzyme, a build-up of peroxide-like compounds occur, thus causing the plant cell membranes of weeds to rupture.

SURGE® Broadleaf Herbicide For Turf offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- This product exhibits improved cool-weather performance compared to standard "3-way amines".
- High selectivity (turfgrass safety) in established cool-season turfgrass and warm-season turfgrass.
- Sulfentrazone combinations provide rapid and effective weed control for common and troublesome (tough) weed species in turfgrass, including: dandelion, spurge, white clover and dollarweed (pennywort).
- The speed of action (rate of weed phytotoxicity) and the early weed symptoms are features of sulfentrazone combinations compared to standard "3-way amines". Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 10-14 days.
- This product is generally rainfast in as little as 6 hours.

SPRAY PREPARATION AND TANK MIXTURES

In certain applications, liquid fertilizer may replace part of the water as a diluent.

Water as diluent:

Add one-half the required amount of water to the spray tank, then add SURGE® Broadleaf Herbicide For Turf slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying.

When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Do not use tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer the spray solution to alter the pH range as appropriate.

Liquid fertilizers as diluents:

ALWAYS PREMIX SURGE® Broadleaf Herbicide For Turf WITH WATER BEFORE ADDING TO FLUID FERTILIZERS. For liquid nitrogen solutions such as U.A.N. or urea solutions, use a premix of 1 part of this product with 4 parts of water or use a premix with a 1:4 ratio of product to water. For other fluid fertilizers such as suspensions, use a premix of 1 part of this product with 50 to 60 parts of water.

Use suitable sources and rates of fertilizer based upon local recommendations. Refer to the mixing directions on the labels of the liquid fertilizers. Always perform a jar test for compatibility before large scale mixing.

The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer should not be prepared as a tank mixture.

9/17

Adjuvants and spray additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then adjuvant(s) combined with SURGE® Broadleaf Herbicide For Turf would not be recommended. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury.

GROUND EQUIPMENT

Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate flat fan nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 20-220 gallons per acre with spray pressures adjusted to between 20-40 psi. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essential when using this product.
- Over-application or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm

motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.

Hand-operated sprayers including backpack sprayers, compression sprayers, and knapsack sprayers are appropriate for small turfgrass areas when power equipment is unavailable, uneconomical, or impractical.

This product may cause injury to susceptible/non-target plants at the use site by contacting the foliage, stems, or roots. To prevent injury to susceptible crops and other desirable broadleaf plants including but not limited to cotton, legumes, tobacco, tomatoes, garden/vegetable crops, and ornamentals (flowers, trees, and shrubs) avoid contact with the spray solution, spray droplets, and spray mist (fine droplets). Applications are recommended only when there is no potential hazard from spray drift during dormant and active growth periods. Do not apply when conditions are conducive to spray drift from the use site to untreated areas.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

WHERE TO USE

This product provides selective broadleaf control in warm-season and cool-season turfgrass in five (5) use sites.

- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways, aprons, and roughs), and office buildings.
- **Ornamental sites** include turfgrass established around residences, parks, streets, retail

10/17

outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Agricultural site:** Commercial sod production.
- **Non-cropland sites:** Highway rights-of-way (principal, interstate, county, private, and unpaved roads). Roadsides, roadside ditches, road shoulders, road embankments, dividers, medians, parking areas, and paved areas. Municipal, state, and federal lands. Airports, airfields and military installations. Fuel storage areas. Fencerows. Industrial sites (including but not limited to lumber yards, tank farms, storage areas, vacant lots and production facilities). Utility and pipeline rights-of-way.

Prohibitions of Sites:

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.

Turfgrass tolerance:

- The turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur on certain varieties of hybrid bermuda-

grass. Adverse environmental conditions may reduce the selectivity on the turfgrass. Do not apply this product to stressed turf.

- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury.

Prohibitions:

- Do not apply this product to bentgrass greens or tees, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when temperatures are above 90°F, some injury may be expected with spot treatments when air temperatures exceed 90°F.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Injury can occur if this product is applied under any of these or other stress conditions. Under any of these stress conditions, any turf damage caused by the use of this product is beyond the control of PBI/Gordon Corporation and all risk is assumed by the buyer and/or user.
- New York: Only one application per year of this product is allowed. This product is not allowed to be sold, distributed, or used in Nassau or Suffolk counties.

SPRAY DRIFT MANAGEMENT

Do not apply under circumstances where possible drift to unprotected persons or to food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur.

11/17

HOW MUCH TO USE USE RATES AND SPRAY VOLUMES

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

The maximum application rate to turf is 0.9 pounds 2,4-D acid equivalent per acre per

application per site. The maximum number of broadcast applications per treatment site is 2 per year.

The maximum seasonal rate of sulfentrazone contained in this product with two (2) broadcast applications to turfgrass is 0.074 pounds of sulfentrazone per acre per season. The retreatment interval for sequential broadcast applications of this product on turfgrass is two (2) to six (6) weeks depending upon the growth stages of the target weeds.

TABLE 1. USE RATES FOR SOD FARMS, ORNAMENTAL LAWNS AND TURFGRASS.

	RATE	SPRAY VOLUME
COOL-SEASON TURF		
Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues, creeping bentgrass and colonial bentgrass (excluding golf greens and tees). Mixtures of cool-season species in non-cropland areas established for roadside vegetation management or for low maintenance. (Kentucky bluegrass, tall fescue, smooth brome grass and orchardgrass)	3¼-4 Pints/Acre (1.2 to 1.5 fl. oz./ 1,000 sq. ft.)	20-220 Gallons/Acre (0.5 to 5.0 Gallons/ 1,000 sq. ft.)
WARM-SEASON TURF		
Hybrid bermudagrass, common bermudagrass, zoysiagrass, bahiagrass and buffalograss	2¾-3¼ Pints/Acre (1.0 to 1.2 fl. oz./ 1,000 sq. ft.)	20-220 Gallons/Acre (0.5 to 5.0 Gallons/ 1,000 sq. ft.)
Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy: Dormant turf: This product may be applied to fully dormant bermudagrass, fully dormant zoysiagrass, and fully dormant bahiagrass. New York: Only one application per year of this product is allowed. This product is not allowed to be sold, distributed, or used in Nassau or Suffolk counties.		

12/17

SPOT TREATMENT

WITH HAND-OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS, COMPRESSION SPRAYERS, AND KNAPSACK SPRAYERS)

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over-application or rates above those specified on this label including excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motions result in uneven coverage.
- Follow-up applications as spot treatments at a 2-6 week interval are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **For cool-season turfgrass:** Mix 1.2-1.5 fl. oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **For warm-season turfgrass:** Mix 1.0-1.2 fl. oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

CULTURAL TIPS

Irrigation:

- Do not apply this product through any type of irrigation system.
- Rainfast in as little as 6 hours. Do not apply this product immediately before rainfall or irrigation.
- If possible, do not irrigate or water the turfgrass within 6 to 24 hours after application.
- If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product.

13/17

BROADLEAF WEEDS CONTROLLED

SURGE® Broadleaf Herbicide For Turf will control or suppress the following broadleaf weeds and will control or suppress broadleaf weeds that are susceptible to 2,4-D. Apply any time the emerged broadleaf weeds are susceptible.

– BROADLEAF WEEDS –

- | | | |
|---|--|---|
| Aster, white heath
& white prairie | Field oxeye-daisy
(*creeping oxeye) | Poison oak |
| Bedstraw | Filaree, whitestem & redstem | Prickly lettuce
(*compass plant) |
| Beggarweed, creeping | Florida betony | Puncturevine |
| Bindweed | Florida pusley | Purple cudweed |
| Black medic | Ground ivy | Purslane |
| Broadleaf plantain | Groundsel | Ragweed |
| Buckhorn plantain | Hawkweed | Redweed |
| Bull thistle | Healall | Red sorrel (*sheep sorrel) |
| Burdock, common | Henbit | Shepherdspurse |
| Buttercup, creeping | Innocence (Blue-eyed Mary) | Spurge |
| Carpetweed | Knotweed | Thistle |
| Chickweed, common | Lambsquarters | Virginia buttonweed |
| Chicory | Lawn burweed | White clover (*Dutch clover,
honeysuckle clover,
white trefoil, & purplewort) |
| Cinquefoil | Lespedeza, common | Wild carrot |
| Clover | Mallow, common | Wild garlic |
| Curly dock | Matchweed | Wild geranium |
| Dandelion | Mouseear chickweed | Wild lettuce |
| Dayflower | Nutsedge** (yellow) | Wild mustard |
| Deadnettle | Old world diamond flower | Wild onion |
| Dock | Oxalis (*yellow woodsorrel
& creeping woodsorrel) | Wild strawberry |
| Dogfennel | Parsley-piert | Yarrow |
| Dollarweed (*pennywort) | Pennsylvania smartweed | Yellow rocket
and many more
broadleaf weeds |
| False dandelion
(*spotted catsear &
common catsear) | Pepperweed | |
| Field bindweed
(*morningglory &
creeping jenny) | Pigweed | |
| | Pineappleweed | |
| | Plantain | |
| | Poison ivy | |

*Synonyms

**Suppression only when nutsedge is young and actively growing.

14/17

TOUGH WEED CONTROL IN INDUSTRIAL OR LOW MAINTENANCE AREAS

Mixed stands of Kentucky bluegrass, tall fescue, smooth brome grass, orchardgrass and reed canary-grass may be treated.

Treatments of SURGE® Broadleaf Herbicide For Turf may injure or kill legumes including clovers (sweet, yellow, red, crimson, alsike, hop, white), lespedezas, trefoils and vetches.

TABLE 2. RATES OF PRODUCT PER ACRE FOR SPECIFIC HARD-TO-CONTROL WEEDS WITH GROUND EQUIPMENT.

Weed Types	Troublesome Weeds Such As:	Amount of Product, Pints/Acre ¹	When to Apply
Annual Broadleaf	Ivyleaf morningglory, redroot pigweed, cocklebur, sunflower, velvetleaf (butterprint)	2.5-4.0 pints/acre	Spring or fall during active growth.
Biennial	Bull thistle, musk thistle, common burdock	4.0-5.0 pints/acre	Spring or fall during seedling to rosette stage.
Perennial	Burclover, Canada thistle, field bindweed, English daisy, hoary cress (whitetop), <i>Veronica</i> (corn speedwell), wild violet.	4.0-5.0 pints/acre	Spring or fall during bud to bloom stage.

Footnote 1: Use the lower rate within the range specified for highly favorable plant growing conditions and when broadleaf weeds are less than 6 inches in height.

Use the higher rate within the range specified for tall vegetation, dense canopies, weeds beyond the suggested growth stage, or during adverse conditions.

New York: Only one application per year of this product is allowed. This product is not allowed to be sold, distributed, or used in Nassau or Suffolk counties.

Biennial and perennial weeds may require follow-up or sequential treatments.

The maximum application rate is 0.9 pounds 2,4-D acid equivalent per acre per application per site.

The maximum seasonal rate of sulfentrazone contained in this product with two (2) broadcast applications to turfgrass is 0.074 pounds of sulfentrazone per acre per season. The retreatment interval for sequential broadcast applications of this product on turfgrass is two (2) to six (6) weeks depending upon the growth stages of the target weeds.

15/17

Spray volumes with ground equipment:

- For SURGE® Broadleaf Herbicide For Turf alone, use a minimum spray volume of 10.0 gallons per acre (gpa).
- For tank mixtures of SURGE® Broadleaf Herbicide For Turf, use a minimum spray volume of 20.0 gallons per acre or use the specified spray volume of the companion product(s).

Adjuvants:

The addition of adjuvants (surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents activators, detergents, and drift reducing agents,) is not necessary when using this product. Under extreme environmental conditions such as drought, the addition of a surfactant may improve efficacy. However, if the foliar phytotoxicity (leaf burn and necrosis) occurs too quickly, then the translocation of SURGE® Broadleaf Herbicide For Turf will not occur and the level of performance (control) may be reduced in perennial broadleaf weeds. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use

16/17

of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

PROFORM® and SURGE® are registered trademarks of PBI/Gordon Corporation.

® Checkered Flag/Label Design is a registered trademark of PBI/Gordon Corporation.

655/8-2006 AP041106
EPA REG. NO. 2217-867
EPA EST. NO. 2217-KS-1

Manufactured By



An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com

THIS PAGE BLANK INTENTIONALLY.