

2217-882

07/26/2005

1/4



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
2217-882

Date of Issuance:
JUL 26 2005

Term of Issuance:
Conditional

Name of Pesticide Product:
EH-1415 Professional

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corp.
1217 West 12th Street
P.O. Box 014090
Kansas City, MO 64101-0090

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit/cite all data required for the registration/reregistration review of your product when the Agency requires all registrants of similar products to submit such data.

2. Make the following label changes:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 2217-882.

Signature of Approving Official:

Joanne J. Miller

Date:

JUL 26 2005

7505C
7/19/05

a/14

3. A one year storage stability (830.6317) and corrosion characteristics (830.6317) are required. The one year storage stability study should be generated under warehouse conditions. It is recommended that observations be made at 0, 3, 6, 9, and 12 months. The corrosion characteristics study made be carried out concurrently.

5. Submit two copies of the revised final printed label for the record.

A stamped copy of the label is enclosed for your records.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller
Product Manger (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

3/14

EH-1415 PROFESSIONAL

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester.....	1.153%
Mecoprop-p acid.....	0.174%
Dicamba acid.....	0.073%
Sulfentrazone.....	0.034%
OTHER INGREDIENTS:	98.566%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 15.30 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.765%
- 3.48 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.174%
- 1.46 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.073%
- 0.68 lbs. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per ton or 0.034%

CAS Registry Numbers:

- 2,4-D CAS# 1928-43-4
- Mecoprop-p CAS# 16484-77-8
- Dicamba CAS#1918-00-9
- Sulfentrazone CAS #122836-35-5

Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

[Guaranteed Analysis and other required fertilizer-related statements may be presented on the front panel of the container label. A sample Guaranteed Analysis and related statements are presented in the Appendix of this master label.]

KEEP OUT OF REACH OF CHILDREN:

CAUTION

See back panel for additional Precautionary Statements and First Aid.

NET WEIGHT: 15, 16, 17, 18, 20, 25 or 30 POUNDS

EPA File Symbol No. 2217-xxx

EPA EST. NO. 2217-KS-2

MANUFACTURED BY:



ACCEPTED
with COMMENTS
In EPA Letter Dated:

JUL 26 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2217-882

4/14



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-900-5556 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as MCP-P and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCP-P and 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Paper Bags, Plastic Bags, and Nonrefillable Bulk Bags - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State and local authorities by burning. If burned stay out of smoke.

WHERE TO USE:

EH-1415 Professional is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, dicamba and sulfentrazone. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

EH-1415 Professional is intended to be applied by noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

EH-1415 Professional may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

EH-1415 Professional may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (½ of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass:

Apply 130-174 pounds of EH-1415 Professional per acre of established turfgrass. Or the equivalent application rates are 3.0—4.0 pounds of product per 1,000 square feet of turfgrass.

For New York State only:

Apply 130-166 pounds of EH-1415 Professional per acre of established turfgrass. Or the equivalent application rates are 3.0—3.8 pounds of product per 1,000 square feet of turfgrass.

For St. Augustinegrass, centipedegrass, hybrid bermudagrass, bentgrass, and colonial bentgrass:

Apply 87-109 pounds of EH-1415 Professional per acre of established turfgrass. Or the equivalent application rates are 2.0-2.5 pounds of product per 1,000 square feet of turfgrass.

The maximum application rate to turf is 1.4 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

6/14

APPLICATION SUGGESTIONS:

1. Spring and fall applications are recommended for the best control. However, EH-1415 Professional can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (>½ inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<¼ inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. [Refer to table for appropriate bag sizes.]

- ◆ For 2.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 500 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet.
- ◆ For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.
- ◆ For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- ◆ For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop.....	5
Scotts SpeedyGreen® Rotary.....	3½
GreenView Drop (Spred II).....	4
Central Drop (#205).....	5½
Sears Drop (#19810).....	4
Ortho Drop (#5000).....	5
Precision Drop (Red Devil).....	4

- ◆ For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- ◆ For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- ◆ For 3.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 260 square feet and adjust to apply at the rate of 1 pound of product per 260 square feet.
- ◆ For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

7/14

Spreader	Delivers ___ lbs. /1,000 sq. ft.							
	2.0 lbs.	2.5 lbs	3.0 lbs	3.2 lbs	3.4 lbs	3.6 lbs	3.8 lbs	4.0 lbs
LESCO® Rotary calibration gauge setting #	X		X		X	X		X
Ortho® Drop (5000)	X		X		X	X		X
Ortho® Rotary (6000)	X		X		X	X		X
Precision® Red Devil® Rotary	X		X		X	X		X
Quaker® Rotary	X		X		X	X		X
Republic® EZ Drop	X		X		X	X		X
Republic® EZ Grow	X		X		X	X		X
Republic® EZ Rotary	X		X		X	X		X
Scotts® Drop (PF3)	X		X		X	X		X
Scotts® Easy Green	X		X		X	X		X
Scotts® Speedy Green	X		X		X	X		X
True Temper Rotary	X		X		X	X		X

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
15	2.0	7,500
	2.5	6,000
	3.0	5,000
	3.2	4,700
	3.4	4,400
	3.6	4,200
	4.0	3,750
16	2.0	8,000
	2.5	6,400
	3.0	5,300
	3.2	5,000
	3.4	4,700
	3.6	4,400
	4.0	4,000
17	2.0	8,500
	2.5	6,800
	3.0	5,700
	3.2	5,300
	3.4	5,000
	3.6	4,700
	4.0	4,250
18	2.0	9,000
	2.5	7,200
	3.0	6,000
	3.2	5,600
	3.4	5,300
	3.6	5,000
	4.0	4,500

8/14

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
20	2.0	10,000
	2.5	8,000
	3.0	6,700
	3.2	6,400
	3.4	5,800
	3.6	5,700
	4.0	5,000
24	2.0	12,000
	2.5	9,600
	3.0	8,000
	3.2	7,500
	3.4	7,100
	3.6	6,700
	4.0	6,000
25	2.0	12,500
	2.5	10,000
	3.0	8,300
	3.2	7,800
	3.4	7,400
	3.6	6,900
	4.0	6,250
26	2.0	13,000
	2.5	8,000
	3.0	8,700
	3.2	8,100
	3.4	7,600
	3.6	7,200
	4.0	6,500
28	2.0	14,000
	2.5	11,200
	3.0	9,300
	3.2	8,800
	3.4	8,200
	3.6	7,800
	4.0	7,000
36	2.0	18,000
	2.5	14,400
	3.0	12,000
	3.2	11,300
	3.4	10,500
	3.6	10,000
	4.0	9,000
37	2.0	18,500
	2.5	14,800
	3.0	12,300
	3.2	11,600
	3.4	10,900
	3.6	10,300
	4.0	9,250

9/14

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
39	2.0	19,500
	2.5	15,600
	3.0	13,000
	3.2	12,200
	3.4	11,500
	3.6	10,800
	4.0	9,750
40	2.0	20,000
	2.5	16,000
	3.0	13,300
	3.2	12,500
	3.4	11,800
	3.6	11,100
	4.0	10,000
42	2.0	21,000
	2.5	16,800
	3.0	14,000
	3.2	13,100
	3.4	12,400
	3.6	11,700
	4.0	10,500
45	2.0	22,500
	2.5	18,000
	3.0	15,000
	3.2	14,100
	3.4	13,200
	3.6	12,500
	4.0	11,300
50	2.0	25,000
	2.5	20,000
	3.0	16,700
	3.2	15,600
	3.4	14,700
	3.6	13,900
	4.0	12,500

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One 50 lb. bag covers up to XXXX square feet.

10/14

SINGLE PASS DISTRIBUTION:

For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.

1. For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage.
2. First, spread the borders of the lawn. Then apply this product to the center with straight line passes
3. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

Advisory for Fertilizers Containing Iron:

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may contain iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

Fertilizer Labeling Requirements by Association of American Plant Food Control Officials (AAPFCO):

- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metal.htm>.

OR

- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-xx.com>

Each registrant shall substitute a unique alphanumeric identifier for "xx". This statement may be used only if the licensee establishes and maintains the Internet site: there is a clearly visible, direct hyperlink to a government website; and the Internet site contains no advertising or company specific information

USE PRECAUTIONS:

Do not use on dichondra, carpetgrass or turfgrass where desirable clovers are present.

Do not apply this product to newly seeded grasses until well-established or after the third mowing.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area.

Do not use this product on 'Floritam' St. Augustinegrass.

11/14

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with EH-1415 Professional. A partial list of susceptible weeds is shown below.

- BROADLEAF WEEDS -			
Annual yellow sweetclover	Cinquefoil	Mexicanweed	Spanishneedles
Aster	Clover, crimson	Milk vetch	Speedwell
Austrian fieldcress	Clover, hop	Mouseear hawkweed	Spiny amaranth
Bedstraw	Clover, red	Mugwort	Spiny sowthistle
Beggarticks	Clover, strawberry	Narrowleaf plantain	Spotted catsear
Betony, Florida	Clover, sweet	Nettle	Spotted spurge
Bird vetch	Clover, white	Orange hawkweed	Spurweed
Bitter wintercress	Cockle	Oxalis	Strawberry, India mock
Bittercress, hairy	Creeping jenny	Parsley-piert	Tall nettle
Bitterweed	Cudweed	Parsnip	Tansy ragwort
Black-eyed Susan	Curly dock	Pearlwort	Tansy mustard
Black medic	Daisy, English	Pepperweed	Thistle
Blackseed plantain	Daisy fleabane	Pigweed	Trailing crownvetch
Blessed thistle	Daisy, oxeye	Pineywoods bedstraw	Tumble mustard
Bloodflower milkweed	Dandelion	Plains coreopsis	Tumble pigweed
Blue lettuce	Dichondra	Plantain	Venice mallow
Blue vervain	Dollarweed	Poison ivy	Virginia buttonweed
Bracted plantain	False dandelion	Poison oak	Virginia pepperweed
Bristly oxtongue	Falseflax	Poorjoe	Wavyleaf bullthistle
Broadleaf dock	False sunflower	Prairie sunflower	Western clematis
Broadleaf plantain	Florida pusley	Prostrate knotweed	Wild aster
Buckhorn plantain	Galinsoga	Prostrate pigweed	Wild carrot
Bulbous buttercup	Goldenrod	Prostrate spurge	Wild four-o'clock
Bull thistle	Ground ivy	Puncturevine	Wild garlic
Bullnettle	Gumweed	Purslane, common	Wild geranium
Burclover	Hairy fleabane	Ragweed	Wild lettuce
Burdock	Hawkweed	Red sorrel	Wild marigold
Burning nettle	Healall	Redroot pigweed	Wild onion
Burweed	Heartleaf drymary	Redstem filaree	Wild rape
Buttercup	Heath aster	Rough cinquefoil	Wild strawberry
Buttonweed	Henbit	Russian pigweed	Wild vetch
Canada thistle	Hoary cress	Scarlet pimpernel	Woodsorrel
Carolina geranium	Hoary plantain	Sheep sorrel	Woolly croton
Carpetweed	Hoary vervain	Shepherdspurse	Woolly plantain
Catchweed bedstraw	Knawel	Slender plantain	Wormseed
Catnip	Knotweed	Smallflower galinsoga	Yarrow
Catsear	Lambsquarters	Smooth dock	Yellow rocket
Chickweed, common	Lespedeza	Smooth pigweed	Yellowflower pepperweed
Chickweed, mouseear	Mallow	Sorrel	and other broadleaf weeds
Chicory	Matchweed	Sowthistle	

12/14

LIMITED WARRANTY AND DISCLAIMER.

IMPORTANT: Read this Limited Warranty and Disclaimer before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this Limited Warranty and Disclaimer in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the directions for use, subject to the inherent risks described below. **THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

I. Guaranteed Analysis:

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅ and K₂O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

II. State mandated fertilizer labeling requirements:

Association of American Plant Food Control Officials (AAPFCO):

- Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm> .
- or
- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-xx.com>

Each registrant shall substitute a unique alphanumeric identifier for "xx". This statement may be used only if the licensee establishes and maintains the Internet site: there is a clearly visible, direct hyperlink to a government website; and the Internet site contains no advertising or company specific information

States:

- Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing ___ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.

III. Fertilizer Grades:

Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

19-6-8	23-3-7	26-4-4	30-4-4
22-3-3	25-3-3	26-4-6	32-3-3
22-3-6	25-4-8	28-0-3	32-4-8
22-3-17	26-3-6	29-3-4	24-4-8
22-4-11		30-3-3	