



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460**

**OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION**

June 1, 2023

Eric D. Smith  
Director, Regulatory Affairs  
PBI/Gordon Corporation  
P.O. Box 860350  
Shawnee, KS 66286

Subject: Notification per PRN 98-10 – Updating company address, adding marketing claims, removing an Idaho state restriction, and clarifying that the primary brand name is, “EH-1414 Weed and Feed” not “EH-1414 Weed & Feed.”

Product Name: EH-1414 Weed and Feed  
EPA Registration Number: 2217-884  
Application Date: January 29, 2019  
Decision Number: 551436

Dear Eric Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact Julia Kerr by phone at 202-566-2810 or via email at [kerr.julia@epa.gov](mailto:kerr.julia@epa.gov).

Page 2 of 2  
EPA Reg. No. 2217-884  
Decision No. 551436

Sincerely,

A handwritten signature in black ink, appearing to read "Mindy Ondish".

FOR

Mindy Ondish  
Product Manager 23  
Herbicide Branch  
Registration Division (7505T)  
Office of Pesticide Programs

**NOTIFICATION**

2217-884

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/01/2023

**EH-1414 WEED & AND FEED**

**EPA Reg. No. 2217-884**

**ACTIVE INGREDIENTS:**

2,4-D, 2-ethylhexyl ester .....	1.108%
Mecoprop-p acid .....	0.167%
Dicamba acid .....	0.071%
<b>OTHER INGREDIENTS:</b> .....	<u>98.654%</u>
<b>TOTAL</b> .....	<b>100.000%</b>

**THIS PRODUCT CONTAINS:**

- 14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
  - 3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
  - 1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9
- Isomer Specific by AOAC Methods  
 TRIMEC® is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid.

**Guaranteed Analysis:** See appendix for example.

**Net Contents:** \_\_\_\_

**EPA Est. No.** \_\_\_\_

**Company Name and Address:**

PBI/Gordon Corporation  
1217 West 12<sup>th</sup> Street P.O. Box 860350  
Kansas City, Missouri 64101 Shawnee, Kansas 66286



---

## SUBLABEL 1

**[Note to Reviewer: Residential sites and outdoor domestic use with applications by homeowners]**

---

### **PRECAUTIONARY STATEMENTS**

#### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

#### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

#### **[Environmental Hazards (alternate per PRN 2008-1)]**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

- Do not use on dichondra, carpetgrass, 'Floritam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
- Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

#### **INSTRUCTIONS:**

EH-1414 Weed **8** and Feed is a combination of three proven weed killers--2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, and spurge In addition, a x-x-x (grade percentage) fertilizer provides a supply of plant nutrients.

#### **USE PRECAUTIONS:**

- Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
- Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

#### **HOW MUCH TO APPLY [WHERE TO USE]:**

##### **[option 1a]**

**For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, and zoysiagrass:** Apply 2.5 to 4.0 pounds of product per 1,000 square feet of established turfgrass.

**For St. Augustinegrass (not 'Floritam'), centipedegrass, hybrid bermudagrass, and colonial bentgrass:** Apply 2.0 to 3.0 pounds of product per 1,000 square feet of established turfgrass.

Do not apply more than 4 pounds of product per 1000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 8 pounds of product per 1000 square feet per year. [note to reviewer: the following information may be omitted from the final printed label: 4 pounds of this product contains 0.029 lb 2,4-D (ae), 0.0067 lb mecoprop-p (ae), and 0.0028 lb dicamba (ae). This is equivalent to 1.28 lb 2,4-D (ae), 0.29 lb mecoprop-p (ae), and 0.12 lb dicamba (ae) per acre.]

##### **[Alternate optional maximum rate statement]**

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.28 lb 2,4-D ae, 0.29 lb MCPP-p ae and 0.12 dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (2.56 lb 2,4-D ae, 0.58 lb MCPP-p ae and 0.25 lb dicamba ae per acre.)]

Generally, the lower application rates within the specified range will provide satisfactory control of annual broadleaf weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

Typically a lawn utilizes [xx] pounds of nitrogen per 1,000 sq.ft. per growing season. When applied as directed, this product provides [yy] pounds of nitrogen per 1,000 sq.ft. Contact your local Extension Service for information specific to your geographic region.

**[option 1b—specifically for 14 lb bags]**

**[For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass.]:** Apply 2.8 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

**option 1c—specifically for 13lb bags]**

**[For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass.]:** Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

**WHEN TO APPLY:**

Apply when broadleaf weeds are actively growing in spring or early fall.

Apply when the air is calm (Choose a calm day) to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

**WHAT TO EXPECT [APPLICATION INSTRUCTIONS]:**

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

<b>WEEDS [253]</b>			
Amaranth, purple	Cupid shaving Brush	Mugwort	Speedwell, common
Amaranth, slender	Cudweed	Mustard, ball	Speedwell, corn
Aster, heath	Daisy, English	Mustard, black	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, blue	Speedwell, Persian
Aster, purple	Dandelion, common	Mustard, hedge	Speedwell, purslane
Aster, white prairie	Dandelion, false	Mustard, Indian,	Speedwell, slender
Artichoke	Day flower	Mustard, tansy	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, tumble	Speedwell, thymeleaf
Bedstraw, Catchweed	Dichondra	Mustard, white	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Mustard, wormseed	Spurge, spotted
Bedstraw, Northern	Dock, curly	Nettle, stinging (burning)	Spurge, prostrate
Bedstraw, pineywoods	Dogbane	Nettle, tall	Spurweed
Beggarticks, hairy	Dogfennel	Nettle, wood	Star of Bethlehem
Betony, Florida	Dollarweed	Parsley-piert	Strawberry, India mock
Bindweed, Field	(Pennywort)	Parsnip	Tansy ragwort
Bindweed, Hedge	Dove weed	Pearlwort	Tanweed
Bitter Wintercress	Elderberry	Pennycress (frenchweed)	Thistle, blessed
Bittercress, hairy	Falseflax	Pepperweed, perennial	Thistle, bull
Bitterweed	False sunflower	Pepperweed, Virginia	Thistle, Canada
Black-eyed Susan	Fiddleneck	(Peppergrass)	Thistle, flodman (prairie)
Black Medic	Filaree, redstem	Pigweed, prostrate	Thistle, musk

Blood flower milkweed	Filaree, whitestem	Pigweed, redroot	Thistle, platte
Blue lettuce	Fleabane, daisy	Pigweed, russian	Thistle, plumeless
Brassbuttons	(annual)	Pigweed, smooth	Thistle, Russian
Bristly oxtongue	Fleabane, rough	Pigweed, tumble	Thistle, scotch
Broomweed	Galinsoga, hairy	Pineapple weed	Thistle, wavyleaf
Buckhorn	Galinsoga, smallflower	Plains coreopsis (tickseed)	Thistle, yellow (horrible)
Bullnettle	Goldenrod	Plantain, blackseed	Trailing crownvetch
Burclover	Ground ivy	Plantain, bracted	Velvetleaf
Burdock	Groundsel	Plantain, broadleaf	Vervain, blue
Burweed, lawn (spurweed)	Gumweed	Plantain, buckhorn	Vervain, hoary
Buttercup, bulbous	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Buttercup, creeping	Hawkweed, meadow	Plantain, narrowleaf	Vervain, tall
Buttonweed, common	Hawkweed, mouseear	Plantain, slender	Vetch, bird
(poorjoe)	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Buttonweed, Virginia	Hawkweed, yellow	Poison ivy	Vetch, milk
Carolina geranium (wild)	Healall	Poison oak	Vetch, narrowleaf
Carpetweed	Heartleaf drymary	Pokeweed	Vetch, wild
Catnip	Hemp	Prairie Sunflower	Virginia creeper
Catsear, spotted (false	Henbit	Prickly lettuce (compass plant)	Western clematis
dandelion)	Hoary cress	Prickly sida	Western salsify
Chamber bitter	Horsenettle	Prostrate knotweed	Wild aster
Chickweed, common	Horseweed	Puncturevine	Wild buckwheat
Chickweed, mouseear	Jimsonweed	Purslane, common	Wild carrot
Chickweed, sticky	Johnny-Jumpup violet	Pusley, brazil	Wild four-o'clock
Chicory	Knawel	Pusley, Florida	Wild garlic
Cinquefoil	Knotweed, prostrate	Pusley, large flower	Wild lettuce
Clover, crimson	Kochia	Ragweed, bur	Wild marigold
Clover, alyce	Lambsquarters	Ragweed, common	Wild mustard (yellow rocket)
Clover, hop	Lespedeza	Ragweed, lancleaf	Wild onion
Clover, rabbit foot	Mallow, alkali	Ragweed, western	Wild parsnip
Clover, red	Mallow, bristly	Redstem filaree	Wild radish
Clover, strawberry	Mallow, common	Rough cinquefoil	Wild rape
Clover, yellow sweet	Mallow, dwarf	Scarlet pimpernel	Wild strawberry
Clover, white sweet	Mallow, venice	Shepard's purse	Wild sweet potato
Clover, white	Marcela	Smartweed, ladysthumb	Wild violet
Cockle	Matchweed	Smartweed, pale	Woodsorrel, creeping
Cocklebur	Mexicanweed	Smartweed, Pennsylvania	Woodsorrel, violet
Cocklebur, oriental	Moneywort	Smooth chaff-flower	Woodsorrel, yellow (Oxalis)
Cocklebur, spiny	Morningglory, bigroot	Smoothdock	Woolly croton
Common mullein	Morningglory, ivyleaf	Sorrel, red (sheep)	Wormseed
Corn Spurry	Morningglory, Mexican	Sowthistle, annual	Yarrow, common
Creeping Beggarweed	Morningglory, red	Sowthistle, spiny	Yarrow, western
Creeping Jenny (creeping	Morningglory, tall	Spanish needles	Yarrow, yellow
Charlie)	Morningglory, wolly	Speedwell, bi-lobed	

**[SPREADER SETTINGS] [DISTRIBUTION AND SPREADER PATTERNS]  
[HOW TO APPLY - SPREADERS AND CALIBRATION]:**

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

**Coverage** - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet.

**IMPORTANT:** It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applicators is optional – it may be presented on the container labels at your discretion.]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
14	2.0	7,000	26	2.0	7,500
	2.5	5,600		2.5	8,000
	2.8	5,000		3.0	8,700
	3.0	4,670		3.2	8,100
	3.2	4,375		3.4	7,600
	3.4	4,120		3.6	7,200
	3.6	3,890		4.0	6,500
15	2.0	7,500	28	2.0	14,000
	2.5	6,000		2.5	11,200
	3.0	5,000		3.0	9,300
	3.2	4,700		3.2	8,800
	3.4	4,400		3.4	8,200
	3.6	4,200		3.6	7,800
16	4.0	3,750	36	4.0	7,000
	2.0	8,000		2.0	18,000
	2.5	6,400		2.5	14,400
	3.0	5,300		3.0	12,000
	3.2	5,000		3.2	11,300
	3.4	4,700		3.4	10,500
	3.6	4,400		3.6	10,000
17	4.0	4,000	37	4.0	9,000
	2.0	8,500		2.0	18,500
	2.5	6,800		2.5	14,800
	3.0	5,700		3.0	12,300
	3.2	5,300		3.2	11,600
	3.4	5,000		3.4	10,900
	3.6	4,700		3.6	10,300
18	4.0	4,250	40	4.0	9,250
	2.0	9,000		2.0	20,000
	2.5	7,200		2.5	16,000
	3.0	6,000		3.0	13,300
	3.2	5,600		3.2	12,500
	3.4	5,300		3.4	11,800
	3.6	5,000		3.6	11,100
20	4.0	4,500	42	4.0	10,000
	2.0	10,000		2.0	21,000
	2.5	8,000		2.5	16,800
	3.0	6,700		2.8	15,000
	3.2	6,400		3.0	14,000
	3.4	5,800		3.2	13,100
	3.6	5,700		3.4	12,400
21	4.0	5,000	45	3.6	11,700
	2.0	10,500		4.0	10,500
	2.5	8,400		2.0	22,500
	2.8	7,500		2.5	18,000
	3.0	7,000		3.0	15,000
	3.2	6,550		3.2	14,100
	3.4	6,150		3.4	13,200
3.6	5,800	3.6	12,500		
4.0	5,250	4.0	11,300		



Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
25	2.0	12,500	50	2.0	25,000
	2.5	10,000		2.5	20,000
	3.0	8,300		3.0	16,700
	3.2	7,800		3.2	15,600
	3.4	7,400		3.4	14,700
	3.6	6,900		3.6	13,900
	4.0	6,250		4.0	12,500

**SINGLE PASS DISTRIBUTION:**

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

**CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS:**


Calibrate the spreader according to the manufacturer’s directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.

- For 2.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 500 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet.
- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.
- For 2.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 357 square feet and adjust to apply at the rate of 1 pound of product per 357 square feet.
- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 3.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 260 square feet and adjust to apply at the rate of 1 pound of product per 260 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

**BROADCAST SPREADERS:** For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.

**DROP SPREADERS:** Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.



Spreader	Delivers ___ lbs./1,000 sq.ft.				
	2.0 lb	2.5 lb	2.8 lb	3.0 lb	3.5 lb
LESCO® Rotary calibration gauge setting #	X	X	X	X	X
Ortho® Drop (5000)	X	X	X	X	X

Ortho® Rotary (6000)	X	X	X	X	X
Precision® Red Devil® Rotary	X	X	X	X	X
Quaker® Rotary	X	X	X	X	X
Republic® EZ Drop	X	X	X	X	X
Republic® EZ Grow	X	X	X	X	X
Republic® EZ Rotary	X	X	X	X	X
Scotts® Drop (PF3)	X	X	X	X	X
Scotts® Easy Green	X	X	X	X	X
Scotts® Speedy Green	X	X	X	X	X
True Temper Rotary	X	X	X	X	X

[End optional language]

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

**FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

### [Alternate Warranty and Disclaimer

WARRANTY & DISCLAIMER: To the extent consistent with applicable law, [company name] guarantees this fine product to the extent of the purchase price, only when used according to directions. ]

---

## SUBLABEL 2

### Institutional, ornamental, and residential sites with application by occupational users

---

#### **PRECAUTIONARY STATEMENTS**

##### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

##### **Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks.
- chemical-resistant gloves

##### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

<h5><b>User Safety Recommendations</b></h5> <ul style="list-style-type: none"><li>• Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.</li><li>• Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.</li><li>• Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li></ul>
---



<h5><b>First Aid</b></h5> <table border="1"><tr><td><b>If in eyes:</b></td><td><ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul></td></tr></table> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.</p>	<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>	



##### **Environmental Hazards**

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow,

may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **USE RESTRICTIONS:**

- Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on dichondra, carpetgrass, creeping bentgrass, 'Floritam' St. Augustinegrass, or turf containing desirable clovers.

### **USE PRECAUTIONS:**

- Do not use this product on dichondra, carpetgrass, creeping bentgrass, 'Floritam' St. Augustinegrass, or turf containing desirable clovers.
- Do not apply this product to newly seeded grasses until well-established or after the third mowing.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area.
- This product will not prevent weed seeds from producing new plants.
- Reseeding or overseeding treated areas may occur three (3) weeks after application of this product.
- Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

### **WHERE TO USE:**

EH-1414 Weed & Feed is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

EH-1414 Weed & Feed is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

EH-1414 Weed & Feed may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

EH-1414 Weed & Feed may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.

**HOW MUCH TO APPLY:**

**For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, and zoysiagrass:** Apply 109 to 174 pounds of EH-1414 Weed & Feed per acre of established turfgrass. Or the equivalent application rates are 2.5 to 4.0 pounds of product per 1,000 square feet of turfgrass.

**For St. Augustinegrass (not 'Floritam), centipedegrass, hybrid bermudagrass and colonial bentgrass:** Apply 87 to 130 pounds of EH-1414 Weed & Feed per acre of established turfgrass. Or the equivalent application rates are 2.0 to 3.0 pounds of product per 1,000 square feet of turfgrass.

Generally, the lower application rates within the specified range will provide satisfactory control of annual broadleaf weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for extreme environmental conditions, or for weeds beyond the appropriate growth stages.

Limitations on broadcast applications to ornamental turfgrass.				
Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum seasonal rate
Ornamental turfgrass	174 pounds/A	2	30 Days	348 pounds/A
	(1.28 lb 2,4-D ae/A) (0.29 lb MCP-p ae/A) (0.12 lb dicamba ae/A)			(2.56 lb 2,4-D ae/A) (0.58 lb MCP-p ae/A) (0.25 lb dicamba ae/A)
ae = Acid Equivalent. Do not exceed the maximum seasonal rate of 348 pounds (2.56 lb 2,4-D ae), 0.58 lb MCP-p ae and 0.25 lb dicamba ae) per acre per season.				

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

Typically a lawn utilizes [xx] pounds of nitrogen per 1,000 sq.ft. per growing season. When applied as directed, this product provides [yy] pounds of nitrogen per 1,000 sq.ft. Contact your local Extension Service for information specific to your geographic region.

**APPLICATION DIRECTIONS:**

1. Spring and fall applications are recommended for the best control. However, EH-1414 Weed & Feed can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (> 1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

WEEDS [253]			
Amaranth, purple	Cupid shaving Brush	Mugwort	Speedwell, common

Amaranth, slender	Cudweed	Mustard, ball	Speedwell, corn
Aster, heath	Daisy, English	Mustard, black	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, blue	Speedwell, Persian
Aster, purple	Dandelion, common	Mustard, hedge	Speedwell, purslane
Aster, white prairie	Dandelion, false	Mustard, Indian,	Speedwell, slender
Artichoke	Day flower	Mustard, tansy	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, tumble	Speedwell, thymeleaf
Bedstraw, Catchweed	Dichondra	Mustard, white	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Mustard, wormseed	Spurge, spotted
Bedstraw, Northern	Dock, curly	Nettle, stinging (burning)	Spurge, prostrate
Bedstraw, pineywoods	Dogbane	Nettle, tall	Spurweed
Beggarticks, hairy	Dogfennel	Nettle, wood	Star of Bethlehem
Betony, Florida	Dollarweed	Parsley-piert	Strawberry, India mock
Bindweed, Field	(Pennywort)	Parsnip	Tansy ragwort
Bindweed, Hedge	Dove weed	Pearlwort	Tanweed
Bitter Wintercress	Elderberry	Pennycress (frenchweed)	Thistle, blessed
Bittercress, hairy	Falseflax	Pepperweed, perennial	Thistle, bull
Bitterweed	False sunflower	Pepperweed, Virginia	Thistle, Canada
Black-eyed Susan	Fiddleneck	(Peppergrass)	Thistle, flodman (prairie)
Black Medic	Filaree, redstem	Pigweed, prostrate	Thistle, musk
Blood flower milkweed	Filaree, whitestem	Pigweed, redroot	Thistle, platte
Blue lettuce	Fleabane, daisy	Pigweed, russian	Thistle, plumeless
Brassbuttons	(annual)	Pigweed, smooth	Thistle, Russian
Bristly oxtongue	Fleabane, rough	Pigweed, tumble	Thistle, scotch
Broomweed	Galinsoga, hairy	Pineapple weed	Thistle, wavyleaf
Buckhorn	Galinsoga, smallflower	Plains coreopsis (tickseed)	Thistle, yellow (horrible)
Bullnettle	Goldenrod	Plantain, blackseed	Trailing crownvetch
Burclover	Ground ivy	Plantain, bracted	Velvetleaf
Burdock	Groundsel	Plantain, broadleaf	Vervain, blue
Burweed, lawn (spurweed)	Gumweed	Plantain, buckhorn	Vervain, hoary
Buttercup, bulbous	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Buttercup, creeping	Hawkweed, meadow	Plantain, narrowleaf	Vervain, tall
Buttonweed, common	Hawkweed, mouseear	Plantain, slender	Vetch, bird
(poorjoe)	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Buttonweed, Virginia	Hawkweed, yellow	Poison ivy	Vetch, milk
Carolina geranium (wild)	Healall	Poison oak	Vetch, narrowleaf
Carpetweed	Heartleaf drymary	Pokeweed	Vetch, wild
Catnip	Hemp	Prairie Sunflower	Virginia creeper
Catsear, spotted (false	Henbit	Prickly lettuce (compass plant)	Western clematis
dandelion)	Hoary cress	Prickly sida	Western salsify
Chamber bitter	Horsenettle	Prostrate knotweed	Wild aster
Chickweed, common	Horseweed	Puncturevine	Wild buckweat
Chickweed, mouseear	Jimsonweed	Purslane, common	Wild carrot
Chickweed, sticky	Johnny-Jumpup violet	Pusley, brazil	Wild four-o'clock
Chicory	Knawel	Pusley, Florida	Wild garlic
Cinquefoil	Knotweed, prostrate	Pusley, large flower	Wild lettuce
Clover, crimson	Kochia	Ragweed, bur	Wild marigold
Clover, alyce	Lambsquarters	Ragweed, common	Wild mustard (yellow rocket)
Clover, hop	Lespedeza	Ragweed, lanleaf	Wild onion
Clover, rabbit foot	Mallow, alkali	Ragweed, western	Wild parsnip
Clover, red	Mallow, bristly	Redstem filaree	Wild radish
Clover, strawberry	Mallow, common	Rough cinquefoil	Wild rape
Clover, yellow sweet	Mallow, dwarf	Scarlet pimpernel	Wild strawberry
Clover, white sweet	Mallow, venice	Shepard's purse	Wild sweet potato
Clover, white	Marcela	Smartweed, ladysthumb	Wild violet
Cockle	Matchweed	Smartweed, pale	Woodsorrel, creeping
Cocklebur	Mexicanweed	Smartweed, Pennsylvania	Woodsorrel, violet
Cocklebur, oriental	Moneywort	Smooth chaff-flower	Woodsorrel, yellow (Oxalis)
Cocklebur, spiny	Morningglory, bigroot	Smoothdock	Woolly croton
Common mullein	Morningglory, ivyleaf	Sorrel, red (sheep)	Wormseed
Corn Spurry	Morningglory, Mexican	Sowthistle, annual	Yarrow, common
Creeping Beggarweed	Morningglory, red	Sowthistle, spiny	Yarrow, western

Creeping Jenny (creeping Charlie)	Morningglory, tall Morningglory, woolly	Spanish needles Speedwell, bi-lobed	Yarrow, yellow
-----------------------------------	--	--	----------------

**DISTRIBUTION AND SPREADER PATTERNS:**

Always apply with a rotary or drop type spreader to distribute the granules evenly. Never apply by hand.

**Coverage** - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One y pound bag covers up to XXXX square feet.

**IMPORTANT:** It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft..	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft..	Treated Area for This Bag Size, Sq.Ft.
15	2.0	7,500	28	2.0	14,000
	2.5	6,000		2.5	11,200
	3.0	5,000		3.0	9,300
	3.2	4,700		3.2	8,800
	3.4	4,400		3.4	8,200
	3.6	4,200		3.6	7,800
	4.0	3,750		4.0	7,000
16	2.0	8,000	36	2.0	18,000
	2.5	6,400		2.5	14,400
	3.0	5,300		3.0	12,000
	3.2	5,000		3.2	11,300
	3.4	4,700		3.4	10,500
	3.6	4,400		3.6	10,000
	4.0	4,000		4.0	9,000
17	2.0	8,500	37	2.0	18,500
	2.5	6,800		2.5	14,800
	3.0	5,700		3.0	12,300
	3.2	5,300		3.2	11,600
	3.4	5,000		3.4	10,900
	3.6	4,700		3.6	10,300
	4.0	4,250		4.0	9,250
18	2.0	9,000	39	2.0	19,500
	2.5	7,200		2.5	15,600
	3.0	6,000		3.0	13,000
	3.2	5,600		3.2	12,200
	3.4	5,300		3.4	11,500
	3.6	5,000		3.6	10,800
	4.0	4,500		4.0	9,750
20	2.0	10,000	40	2.0	20,000
	2.5	8,000		2.5	16,000
	3.0	6,700		3.0	13,300
	3.2	6,400		3.2	12,500
	3.4	5,800		3.4	11,800
	3.6	5,700		3.6	11,100
	4.0	5,000		4.0	10,000

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft..	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft..	Treated Area for This Bag Size, Sq.Ft.
24	2.0	12,000	42	2.0	21,000
	2.5	9,600		2.5	16,800
	3.0	8,000		3.0	14,000
	3.2	7,500		3.2	13,100
	3.4	7,100		3.4	12,400
	3.6	6,700		3.6	11,700
	4.0	6,000		4.0	10,500
25	2.0	12,500	45	2.0	22,500
	2.5	10,000		2.5	18,000
	3.0	8,300		3.0	15,000
	3.2	7,800		3.2	14,100
	3.4	7,400		3.4	13,200
	3.6	6,900		3.6	12,500
	4.0	6,250		4.0	11,300
26	2.0	13,000	50	2.0	25,000
	2.5	8,000		2.5	20,000
	3.0	8,700		3.0	16,700
	3.2	8,100		3.2	15,600
	3.4	7,600		3.4	14,700
	3.6	7,200		3.6	13,900
	4.0	6,500		4.0	12,500

**CALIBRATION INSTRUCTIONS:**

To provide proper distribution, calibrate spreader before application.

- For 2.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 500 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet.
- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.
- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 3.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 260 square feet and adjust to apply at the rate of 1 pound of product per 260 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Spreader	Delivers ___ lb/1,000 sq.ft.							
	2.0 lb	2.5 lb	3.0 lb	3.2 lb	3.4 lb	3.6 lb	3.8 lb	4.0 lb
LESCO® Rotary calibration gauge setting #	X	X	X	X	X	X	X	X
Ortho® Drop (5000)	X	X	X	X	X	X	X	X
Ortho® Rotary (6000)	X	X	X	X	X	X	X	X
Precision® Red Devil® Rotary	X	X	X	X	X	X	X	X
Quaker® Rotary	X	X	X	X	X	X	X	X
Republic® EZ Drop	X	X	X	X	X	X	X	X
Republic® EZ Grow	X	X	X	X	X	X	X	X
Republic® EZ Rotary	X	X	X	X	X	X	X	X
Scotts® Drop (PF3)	X	X	X	X	X	X	X	X



Scotts® Easy Green	X	X	X	X	X	X	X	X
Scotts® Speedy Green	X	X	X	X	X	X	X	X
True Temper Rotary	X	X	X	X	X	X	X	X

**SINGLE PASS DISTRIBUTION:**

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

[End optional language]

**STORAGE & DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**LIMITED WARRANTY AND DISCLAIMER**

**FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

[Alternate Warranty for 82898] [Nutrients PLUS, LLC warrants that this product conforms to the analysis on its label. When used in accordance with label directions, under normal conditions, this product is reasonably fit for its intended purposes. Since time, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable, to the extent consistent with applicable law. The user assumes all risks of any such use.]

# **APPENDIX**

## **1. Advertising claims and marketing phrases that may be presented on promotional materials and the container labels of supplemental registrants.**

### **Herbicide Benefit Claims:**

- Controls Dandelions & Other Broadleaf Weeds
- Controls a Wide Spectrum of Broadleaf Weeds
- Controls Dandelion, White Clover, Plantain, Chickweed, Spurge and Many Other Listed Weeds
- Controls '[Kills] clover, dandelion, plantain, chickweed and [other broadleaf,weeds] [others as listed]
- [Kills] [Controls] [dandelions] [any weed listed on label] and [over] [more than] [XXX] other [listed] major lawn weeds [as listed]
- Kills xx+ Weeds
- Kills xx+ Weed; Types, Even the Roots.
- Kills [down to] the root, [not the lawn]
- Kills the root
- Kills weed roots!
- Kills more weeds
- Kills tough lawn weeds
- Kills the weeds, not the lawn
- Kills weeds, not lawn grasses
- Kills [Over] [More Than] XXX [Listed] Weeds [as Listed]
- Kills broadleaf weeds in lawns
- Kills [major] broadleaf weeds [like clover, dandelion, oxalis]
- Kills major broadleaf weeds, even the roots. And [brand name] is formulated to kill only the weeds, not the lawn. Use as directed to kill major broadleaf weeds in your lawn.
- Kills dandelion and many other broadleaf weeds.
- Kills clover, chickweed, spurge and many other [listed] broadleaf weeds [as listed]
- Kills [listed] weeds now for fewer weeds in spring
- Kills even the tough[est] broadleaf weeds [like clover, dandelion, oxalis]
- Kill the toughest weeds like Clover, Dandelion, Henbit, Chickweed, Spurge and many other listed broadleaf weeds
- Kills Over xxx Weed Types [as listed]
- Kills Weeds While Feeding Your Lawn
- Kills Weeds 2 Times [X] Faster!\* (\*compared to our previous formula)
- Kills xx+ Weed Types Including Roots
- Kills major broadleaf weeds including roots. And (brand name) is formulated to kill only the weeds, not the lawn. Use as directed to kill major broadleaf weeds in your lawn.
- Kills Weeds 2x Faster (compared to previous Weed & Feed)
- Kills Weeds Twice as Fast! (compared to previous Weed & Feed)
- Kills Over [XXX] [Listed] Weeds [as Listed], Including Henbit, Dollar Weed, White Clover, Plantain and Chickweed
- Kills tough lawn weeds
- Kills over XXX weeds
- Provides broadleaf control
- Clobbers clover
- Consistently fast on: tough weeds like clover, plantain, ground ivy and spurge [various other weeds listed]
- Use for a beautiful weed-free lawn
- Helps keep lawns weed-free
- Targets weeds, leaves [grasses][the lawn] unharmed
- Our Fastest Weed Control Formula Ever! (compared to our previous Weed & Feed)
- Our Fastest Weed Killing Formula Ever! (compared to our previous Weed & Feed)
- Sta-Green Turf Note – Control Those Pesky Weeds

- Many broadleaf weeds (dandelion, chickweed, henbit) germinate in the fall but are not visible in your lawn until spring. Applying (product name) in the fall can provide effective results, because smaller weeds are easier to control. If needed, you can make a follow-up application in the spring.
- Eliminates and controls more than 250 weeds in your home lawn
- [Brand Name] Weed & Feed: These post-emergent products control a wide variety of existing broadleaf weeds while they feed your desirable grass.
- [Brand Name] Crabgrass Preventers: Use these pre-emergent products to control crabgrass and many other weeds in your lawn before they germinate. Crabgrass preventers are available with or without fertilizer.
- Wipes out tough weeds such as [chickweed, clover, henbit, and/or dandelion]
- Listed weeds die and disappear from your lawn
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion.
- Tough weed killer
- Kills even the toughest weeds!
- Kills over 200 weed types, even the roots
- Kills [Controls] Twice [2x] as many weeds [weed image icon]\* \*compared to our previous product [formula] of 2217-XXXX (insert product name of the previous product)
- Kills [Controls] [2x] more weeds [weed image icon] than our previous product [formula] of 2217-XXXX (insert product name of the previous product)
- This product is designed to promote lush green growth in lawns & control many broadleaf weeds as listed
- Starts working on contact and kills to the roots
- [(Kills over 200 listed weed types)] guaranteed
- One application kills over 250 weeds and feeds [lawn] for [an][the] entire season
- Kills over 250 weeds down to the root
- Fast acting
- Kills and controls tough lawn weeds as listed

#### Fertilizer Benefit Claims:

- For a lush, thick, deep green lawn
- [Greens in 72 Hours &] Promotes Thick, Green Growth [to repair summer damage]
- Promotes root growth for a thicker, greener lawn
- Fast Green-Up & Timed Release for Extended Feeding
- Quick Greening and Extended Feeding
- Encourages Quick Greening & Provides Extended Nitrogen Feeding up to 2 Months
- Pennington Coated Slow Release Technology. For a long lasting green lawn with no burning.\*  
\*When used according to label directions [\*When used as directed]
- Pennington Coated -- Slow Release Technology -- For a Long Lasting Green Lawn With No Burning\*  
\*When used according to label directions.
- Contains [Composite] Timed-Release (CTR) Nitrogen for Extended Feeding!
- (Product Name) is specially formulated to complement the phosphorus already present in some soils and give your turf the additional primary nutrients needed for a good turf feeding. A soil test can help you determine if your soil has an adequate supply of phosphorus and if this is the best product for your soil type. Proper watering and mowing practices along with regular feedings will help promote a beautiful green lawn.
- Feeds up to 4 months
- Feeds for [up to] 3 months
- Feeds For [up to] 3 Months Guaranteed
- Feeds to thicken lawn and crowd out weeds
- Feeds and strengthens to help protect against future problems
- Phosphorus Free – Helping our Water Sources [to be used only on formulations containing 0% phosphorus]

- Encourages Early Spring Green-Up
- Greens quickly, feeds for months
- **Quick greening**
- Helps develop a deep and dense root system [at optimal time]
- Why fertilize the/your lawn?
- While it is easy to see that/a satisfying deep green/greening when you fertilize, regular feedings [can] actually do much more...
- Promote a deep and dense root system that is better able to reach water and find nutrients during tough heat & drought, so lawn stays greener, longer and bounces back faster
- **Promotes [a stronger, deeper root system][and][a quicker][an easier][spring][green-up][start]**
- **[Promotes turf growth][and][deepens [turf][lawn][grass] roots]**
- Thicken lawn with new/additional blades of grass, making it more resistant to bare spots and thinning
- **Special formula that thickens and promotes a lush, deep green lawn**
- Feed every 8 weeks for optimal results
- Phosphorus Free! [to be used only on formulations containing 0% phosphorous]
- Feeds: Combine quick results and a long lasting feeding
- Replenishes: Adding the right balance of nutrient sources helps create a healthy lawn & soil
- Preserves: Using slowly available nutrients reduces run-off potential
- Timed-release nitrogen for a thick, green lawn
- Promotes a strong, deep root system in the fall for a quicker green-up in the spring
- Slow-release nitrogen promotes lush, deep-green growth for up to four months
- Higher nitrogen content for greater coverage
- Nitrogen for lush, green plant growth
- Nitrogen for greening
- Potassium for overall plant vigor
- Potassium for overall plant health and vigor
- Potash is vital for growth. Improves plant hardiness and drought resistance.
- **Enriched with micro-nutrients to revitalize soil**
- **Strengthens lawn against heat and drought**
- **Builds strong, deep roots to protect from drought and heat**
- [product name] is formulated with Polymer Coated Urea as the source of controlled-released Nitrogen providing your lawn with consistent, even feeding of Nitrogen for up to 4-months. Benefits include: Quick Green-Up, Low Burn Potential, Timed-Release Nitrogen for an Even Green, Minimal Runoff & Leaching, Extended Feeding.
- Phosphorus Free Fertilizer [to be used only on formulations containing 0% phosphorous]
- [Brand Name] Lawn Fertilizer: A high nitrogen fertilizer with iron that feeds any established lawn up to 3 months.
- Surfcote – Sustained Uniform Release Fertilizer Continuously Provides Nutrients
- Helps develop a deep, dense root system that encourages early spring green-up
- Quick greening with extended feeding
- Why go phosphorus free? Too much phosphorus in the soil can lead to ground water contamination. Using Sta-Green fertilizers containing no phosphorus helps our water sources. [Only to be included with phosphorus free formulations]
- Helps strengthen lawn against heat and drought
- Slow release nitrogen for continuous greening.
- Provides both fast and long feeding action
- [Icon] Unfed Lawn v.[vs.] [Icon] Regularly fed Lawn/4 annual feedings
- [Icon] Unfed Lawn
- [Icon] [1] [2] [3] [4] annual feeding[s]
- Contains iron for deep greening & micro-nutrients for a balanced blend
- Fast green-up + extended feeding

- For thick, green, rich turf
- Nitrogen Produces lush, dark green plant growth
- Potassium Helps to improve plant's hardiness and stress tolerance
- BASIC Nutrition
- Powered by BASIC Nutrition
- U-Plus
- Powered By U-Plus
- Contains controlled release fertilizer technology
- With controlled release lawn food
- Consistently feeds for up to [16] weeks
- [Nitrogen granule icon] [polymer coating] [urea] [wax coating]
- For Products containing Iron:
- Contains X% iron for [[quick] deep green][lawn]
  - For a [deep] green lawn
  - For a lush, thick deep green lawn
  - Greens quickly
  - Added iron for deeper green

General Marketing Claims:

- Northern and Southern Grasses Bluegrass, Fescue, Zoysia, Bermuda, St. Augustinegrass, Bentgrass, Centipede, Bahia, Perennial Ryegrass
- For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass and colonial bentgrass
- Treats Northern and Southern Grasses
  - Feed[s] Treats [Covers] XXXX sq.ft. [(XXXX square meters)]
- Covers up to XXXX square feet
- Starts working on contact
- Our fastest formula ever
- [Superior][cool][all] weather performance
- Proven [all] [cool] weather performance
- For home lawn care
- For home lawns only
- For use on lawns
- Guaranteed [Results] or Your Money Back\*
- 100% satisfaction guaranteed or your money back
  - If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail you a refund check promptly.
  - If you are not satisfied with this product for any reason, send your written request for a refund to the address at left, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.
  - If you apply [product name] as directed and are not completely satisfied, send proof of purchase from this package and original cash register receipt to the address below. We will refund the purchase price you paid.
  - GUARANTEE: [company name] guarantees this fine product to the extent of the purchase price, only when used according to directions.
  - MONEY BACK GUARANTEED
  - †MONEY BACK GUARANTEE: If you are not completely satisfied with this product, please mail your name, address and proof of purchase (package UPC) and a brief explanation for seeking a refund.
  - Guarantee – Infinity Fertilizers, Inc. guarantees this fine product to the extent of the purchase price, only when used according to directions.

- [Company Name] No-Quibble Guarantee: If for any reason you, the customer, are not satisfied with results after using this product you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.
- Vigoro Guarantee – [ note: To be optionally used in conjunction with the guarantee "If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail you a refund check promptly."]
- [Ace Promise Satisfaction Guaranteed]/[The Ace Promise is in Every Bag] (front & back panel)
- [The Ace Promise] If this Ace product when used for its intended purpose, fails to give you complete satisfaction, return the item to the Ace Hardware store where purchased for free replacement of the same or similar item.
- [Product Name] WILL NOT BURN when used as directed. We're so confident you'll love the results of [Product Name], that it comes with the Pennington Satisfaction Guarantee. †
- † Pennington Satisfaction Guarantee: If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to:  
Pennington Refund  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339
- 100% satisfaction guaranteed or your money back
- Apply When Weeds Are [Actively] Growing
- In Case of Emergency, Call XXX-XXX-XXXX
- Easy to Use
- Easy Application
- Easy to Apply
- Easy to Use
- Lawn Fertilizer and Lawn Weed Killer
- 4 month formula
- With Pennington Coated Technology
- Helps strengthen lawn against heat & drought
- Apply Early Fall When Weeds are Actively Growing
- Beautiful Lawns Since 1924
- Works the first time or your money back! [icon]
- Apply to a wet lawn
- Use on a wet lawn
- Membership is free! [insert web address]
- Determine size of area to be treated
- Measuring/Measure your lawn
- How to Measure Your Lawn – [ICON]
- Icon – computer mouse
- Icon – computer
- Icon – phone
- Distributed/Sold/Manufactured/Guaranteed by/for [insert company information]
- Superscript used is first letter of run code on end of container
- GENERAL TIPS
  1. Determine size of area to be treated – measure length and width in feet, then multiply to determine coverage area in sq ft (length x width). Next, subtract non-lawn areas.
  2. Use the spreader setting chart to determine the appropriate setting.
  3. Treat as directed – thorough coverage is important.
- Controls lawn weeds and prepares your lawn for the stress of winter
- Apply in the Fall
- 25% More Coverage\* \*Than XX lb bag
- Lawn Fertilizer Plus Weed Control
- Questions or comments? [Ace Customer Care] 1-877-465-5162
- Made for Ace Hardware Corp. [insert address]

- Green Turf® is a registered trademark of Ace Hardware Corporation
- Green Turf® is a registered trademark of United Industries Corp
- Controls many broadleaf weeds while providing long-feeding action
- Help prevent weeds by making it difficult for weed(s) [seeds in the air] to find soil to root, [as well as]/and crowd out existing weed(s) [seeds in the soil]
- Visit us on the internet or call our customer satisfaction hotline [insert company address/phone number/website]
- [Brand Name] Centipede & St. Augustine Weed & Feed: This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns.
- High performance weed control with fertilizer specially formulated for your home lawn during the winter months
- Premium grade herbicide and turf food combination
- Alternate emergency contact info: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 8:00 AM-12:00 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.
- Assembled in the USA with foreign and/or domestic parts
- Assembled in the USA
- Factory Certified [Note to Reviewer: For WalMart]
- Apply [in] [late] [spring] [early] [summer] [early] [fall]
- [Note to Reviewer: This statement may be placed on containers to comply with California Prop 65: [Notice][Attention]: This product contains a chemical known to the State of California to cause cancer and birth defects or other reproductive harm.]
- The Pennington® [Product Name] Difference
- (For products with Iron): This premium blended fertilizer is packed with essential plant nutrients to give you the thick, green lawn you desire. It has both fast acting nitrogen for quick greening, as well as slow release nitrogen for extended feeding for up to 3 months. With X% Iron, this fertilizer will give you a deep, dark green lawn.

#### Referrals to Other Products Used Sequentially:

- Project Guide – For a thicker, greener, more beautiful lawn, follow (Product Program). It's easy to follow and can be started at anytime by simply applying the product that corresponds with the current season!
- [Company Name] Annual Feeding Program
  - Plan your feeding schedule. Choose the right product for your lawn.
    - Spring -- Fall
      - Any Season: Lawn Fertilizer, Apply any season if weeds or bugs are not a problem. [bag illustration]
      - Early Spring: Crabgrass Preventer, February - April to prevent crabgrass. [bag illustration]
      - Late Spring: Weed & Feed, April - June if you have weeds [bag illustration]
      - Summer: Lawn Fertilizer, June - August [bag illustration]
      - Fall: Weed & Feed, September - November if you have falls weeds [bag illustration]
- A Beautiful Pennington Lawn, [Pennington UltraGreen Annual Feeding Program] A good fertilization program can keep your lawn green and vigorous. Pennington has a wide selection of fertilizers to help you throughout the year.
- Pennington Lawn Starter, Recommended for use when seeding, sodding, or reseeding lawns to enhance root development.
- Pennington Crabgrass Preventers, Use these pre-emergent products to control crabgrass and many other weeds in your lawn before they germinate. Crabgrass preventers are available with and without fertilizer.

- Pennington Weed & Feed, These post-emergent products control a wide variety of existing broadleaf weeds while they feed your desirable grass.
- Pennington Lawn Fertilizer, A high nitrogen fertilizer with iron that feeds any established lawn up to 4 months.
- Pennington Centipede & St. Augustine Weed & Feed, This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns.
- For additional spreader information, visit pikenursery.com
- Pike Lawn Care Program: A good fertilization program can keep your lawn green and vigorous. Pike has a wide selection of fertilizers specifically formulated for the Southeast to help you throughout the year.
  - Pike Lawn Starter: Recommended for use when seeding, sodding, or reseeding lawns to enhance root development.
  - Pike Crabgrass Preventer: Use pre-emergent product to control crabgrass and many other weeds in your lawn before they germinate.
  - Pike Weed & Feed: Use post-emergent product to control a wide variety of existing broadleaf weeds while feeding desirable grass.
  - Pike Lawn Fertilizer: A high nitrogen fertilizer with iron that feeds any established lawn up to 4 months. Quick greening & long lasting.
- For A Beautiful Sta-Green Lawn: A good fertilization program can keep your lawn green and vigorous. Sta-Green has a wide selection of fertilizers to help you throughout the year.
- STA-GREEN CRABGRASS PREVENTERS: Use these pre-emergent products to control crabgrass and many other weeds in your lawn before they germinate. Crabgrass preventers are available with or without fertilizer. Use these products in early spring before weeds begin to emerge.
- STA-GREEN LAWN FERTILIZER: A high nitrogen fertilizer with iron that feeds any established lawn up to 4 months. Use this product anytime during the active growing season.
- STA-GREEN WEED & FEED: This post-emergent product controls a wide variety of existing broadleaf weeds while it feeds your desirable grass. Use weed and feeds when weeds are actively growing.
- STA-GREEN SOUTHERN WEED & FEED: This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns. Use weed and feeds when weeds are actively growing.
- [Recommended] Next Feeding
- [icon] Vigoro® Lawn Fertilizer [Apply anytime during the growing season]
- [icon] Vigoro Crabgrass Preventer & Lawn Fertilizer [Apply early spring]
- [icon] Vigoro® Weed & Feed for Bahia & Mixed Lawns [Apply anytime weeds are actively growing\* \*maximum of 2 applications per year]
- [icon] Vigoro® Weed & Feed [Apply anytime weeds are actively growing\* \*maximum of 2 applications per year]
- The Nutrients PLUS® Pathway To Solutions
- If you have questions or would like additional information regarding Clarus™ products, call us at 1-888-675-7515.
- All marks are property of Nutrients PLUS. Patents pending.
- [Home Depot logo] Garden Club – [note: Vigoro products are exclusively sold at Home Depot and the Garden Club Claim will only be used on Vigoro branded products]
- [Brand Name] Annual Feeding Program: A good fertilization program can keep your lawn green and vigorous. [Brand Name] has a wide selection of fertilizers to help you throughout the year.
- Ace Green Turf Winterizer Weed & Feed controls more than 250 [listed] weeds [as listed]
- Ace Green Turf Weed & Feed controls more than 250 [listed] weeds [as listed]
- Use this product in the TurfKing™ program for a great lawn

## **2. State mandated fertilizer labeling requirements:**

**[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are**



**relevant to your labeling, please place in the Directions for Use under the heading “State Restrictions”]**

- California, Oregon, Washington: “[When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials]-Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-xx.com>  
[Each registrant shall substitute a unique alphanumeric identifier for “xx”. This statement may be used only if the licensee establishes and maintains the Internet site: there is a clearly visible, direct hyperlink to a government website; and the Internet site contains no advertising or company specific information ]
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
  - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
  - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
  - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida Annual Fertilizer Applications: [Do not exceed] 4 lbs N/1000 sq ft with the following exceptions:
  - On centipedegrass do not exceed 2 lbs N/1000 sq ft in the north and 3 lbs N/1000 sq ft in the central and south.
  - On [bahia grass and] zoysiagrass do not exceed 3 lbs N/1000 sq ft in the north.
  - North Florida is defined as north of a line extending from Cedar Key to Daytona Beach.
  - Central Florida is defined as south of the line extending from Cedar Key to Daytona Beach to a line extending from Tampa to Vero Beach.
  - South Florida includes the remaining southern portion of the state.
- Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices [for feeding your lawn.]
- Check with your county or city government to determine if there are local regulations for fertilizer use.
- Florida: “Apply only to actively growing turf.”
- ~~Idaho: “Apply only as directed”~~
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”

- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: "In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency."
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Advisory for Fertilizers Containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program."
- Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.
- Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label, and referring to the batch number found on this container.

## **DOCUMENT CONTROL INFORMATION**

1. **Unique Label Identifier:** 002217-00884.20220302.notif-proposed-highlighted.doc

2. **Reason for Issue:** ad claims, update corporate address

<b>Internal Usage Notes</b>
Sublabel 1. Residential sites and outdoor domestic use with applications by homeowners Associated Container Label: Distributor Products
Sublabel 2. Institutional, ornamental, and residential sites with application by occupational users Associated Container Label: Distributor Products