



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

SEP - 8 2008

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

James L. Kunstman
PBI/Gordon Corporation
1217 West 12th St.
P.O. Box 014090
Kansas City, Missouri 64101-0090

Dear Dr. Kunstman:


SUBJECT: RED Label Amendment
EH-1422 Herbicide
EPA Registration No. 2217-892

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- 1) Per the acute toxicity review, revise the statement "Avoid contact with eyes or clothing" in the Hazards to Humans and Domestic Animals section to read "Avoid contact with eyes, skin or clothing."
- 2) Per the product chemistry review, the storage and disposal subheading "Pesticide Disposal" and the required text must be added to the label per 40CFR 156.10(i)(2)(ix) and PR Notice 83-3.

Submit one (1) copy of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records. Please note that final product registration cannot be considered until after all active ingredients in this product are eligible for reregistration.

Sincerely yours,



Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

EH-1422 HERBICIDE

EPA Reg. No. 2217-892

ACTIVE INGREDIENT:

Mecoprop-p, dimethylamine salt.....	2.84%
2,4-D, dimethylamine salt.....	5.32%
Dicamba, dimethylamine salt.....	0.96%
INERT INGREDIENTS:	90.88%
TOTAL.....	100.00%

THIS PRODUCT CONTAINS:

- 0.20 lb. (+)-R-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 2.35%
- 0.38 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 4.42%
- 0.07 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.80%

Isomer Specific by AOAC Method 6.DO1-5.
 TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
SEP - 8 2008

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.
2217-892

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	



Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until sprays have dried.

I. SPOT TREATMENTS WITH HAND-OPERATED SPRAYERS:

Spot treat broadleaf weeds as they appear in your lawn with EH-1422 Herbicide using pump-up or pressure type sprayer.

St. Augustinegrass, Bentgrass, Buffalograss and Centipedegrass: Mix 1 fl.oz. (2 Tbs) in 1 gallon water and uniformly spray 333 sq.ft. See table below for additional spray preparation.

Apply any time weeds are actively growing. Do not apply to St. Augustinegrass or centipedegrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage. Do not use on 'Floritam' St. Augustinegrass in Florida.

If necessary, repeat treatment in 2 to 3 weeks if hard-to-kill weeds persist.

Spray volume, gals. of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	1 fl.oz.	2 tbs.	333 sq.ft.
2 gallons	2 fl.oz.	4 tbs.	666 sq.ft.
3 gallons	3 fl.oz.	6 tbs.	1,000 sq.ft.

Bermudagrass, Bahiagrass, Kentucky Bluegrass, Fescues, Perennial Ryegrass and Zoysiagrass: Mix 2 fl.oz. (4 Tbs) in 1 gallon water and uniformly spray 333 sq.ft. See table below for additional spray preparation. Apply any time weeds are actively growing.

If necessary, repeat treatment in 2 to 3 weeks if hard-to-kill weeds persist. On hard-to-kill species or on older weeds, mix 2.5 fl.oz. in 1 gallon water and spray on 333 sq.ft.

Spray volume, gals. of water	Fluid ounces (fl.oz.) of product	Tablespoons (tbs.) of product	Area to be treated, sq.ft.
1 gallon	2 fl.oz.	4 tbs.	333 sq.ft.
2 gallons	4 fl.oz.	8 tbs.	666 sq.ft.
3 gallons	6 fl.oz.	12 tbs.	1,000 sq.ft.

Equal measures: 1 Tablespoon (Tbs.) = 3 teaspoons (tsp.), 1 Fluid Ounce (fl.oz.) = 2 Tablespoons (Tbs.), 1 Cup = 8 Fluid Ounces (fl.oz.)

Limitations on spot treatments:

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 6.0 fl. oz. per 1,000 sq.ft. per application (0.41 lbs MCP-P acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

II. Instructions for hose-end sprayers including Ortho® Dial-N-Spray® or Gilmour Dial-A-Mix® Lawn and Garden Sprayer.

OPEN

Resealable Label for Directions & Precautions

PRESS TO RESEAL

Kills weeds in your lawn - won't harm lawn grasses. EH-1422 Herbicide singles out weeds, enters through leaves and moves inside the weed to the roots.

PRODUCT FACTS:

- Treats up to 5,333 sq ft (approx. size of 3 tennis courts) [1 pint, 16 fl. oz.]
- Treats up to 10,666 sq ft (approx. size of 5 tennis courts) [1 quart, 32 fl. oz.]
- Treats up to 17,920 sq ft (approx. size of 9 tennis courts) [0.42 gallon, 53 fl. oz.]
- Treats up to 21,333 sq ft (approx. size of 10 tennis courts) [one-half gallon, 64 fl. oz.]
- Treats up to 42,667 sq ft (approx. size of 20 tennis courts) [1 gallon, 128 fl. oz.]

KILLS WEEDS:

Dandelion, Clover, Chickweed [illustration] plus over 200 additional [other broadleaf [weeds]

WHERE TO USE:

ON LAWNS [without harming lawn grasses]

AMOUNT TO USE:

Bent, Buffalo, Centipede & St. Augustine (EXCEPT 'Floratam') (Do not use on 'Floratam' in Florida):
 1.0 fl.oz. (2 Tbs) / gal of water / 333 sq ft
 Bahia, Bermuda, Fescues, Kentucky Blue, Perennial Rye & Zoysia: 2.0 fl. oz. (4 Tbs)/gal of water/333 sq ft

QUESTIONS, COMMENTS OR MEDICAL INFORMATION:

Call 1-800-225-2883
<http://www.companyname.com>

MIXING INSTRUCTIONS

FOR SENSITIVE LAWNS:

Bentgrass, Buffalograss, Centipedegrass & St. Augustinegrass (EXCEPT 'Floratam') (Do not use on 'Floratam' St. Augustinegrass in Florida)

Amount to Use: 1 fl.oz. (2 Tbs) per gallon of water for each 333 sq ft.

- For the Ortho® Dial-N-Spray®: Set dial to 1 oz. Add 3 fl.oz. (6 Tbs) to sprayer jar. DO NOT ADD WATER. Spray evenly over 1,000 sq ft until jar is empty.

FOR HARDY LAWNS:

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass

Amount to Use: 2 fl.oz. (4 Tbs) per gallon of water for each 333 sq ft.

- For the Ortho® Dial-N-Spray®: Set dial to 2 oz. Add 6 fl.oz. (12 Tbs) to sprayer jar. DO NOT ADD WATER. Spray evenly over 1,000 sq ft until jar is empty.

FOR BEST RESULTS	
MEASURE [icon of walking man]	<ul style="list-style-type: none"> To treat entire lawn (broadcast application): Use a tank or [Dial-N-Spray®] hose-end sprayer. Measure lawn. Calculate square feet by multiplying length times width. Spray evenly over measured area. To spot treat individual weeds in lawns: Use a tank sprayer.
MOW	<ul style="list-style-type: none"> Do not mow for 1 to 2 days before or after application.
WATER [illustration]	<ul style="list-style-type: none"> If soil is dry, water 1 to 2 days before application. Rain or watering 2 hours after application will not wash away effectiveness.
WHEN TO APPLY [illustration]	<ul style="list-style-type: none"> Spray when weeds are small and actively growing. [Spray when] temperature is below 90°F. [Spray when] air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants. Reapplication can be made in 2 to 3 weeks if necessary. Newly seeded lawn can be sprayed after first-mowing. Bare spots can be planted with grass seed 1 week after application.

IMPORTANT

- If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots.
- St. Augustinegrass, centipedegrass or bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45° F within 72 hours of application.
- This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass if misapplication occurs.
- Do not spray carpetgrass, dichondra or desirable clovers.
- Do not use on 'Floritam' St. Augustinegrass in Florida.

Other Use Precautions: Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

Limitations on 2,4-D applications to residential turfgrass

The maximum application rate is 6.0 fl.oz. of product per 1,000 sq.ft. per application (0.78 lbs 2,4-D ae, 0.41 lb MCP-P ae and 0.14 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 12.0 fl.oz. of product per 1,000 sq.ft. (1.56 lbs 2,4-D ae, 0.82 lb MCP-P ae and 0.28 lb Dicamba ae per acre).

HOW IT WORKS

Because broadleaf weeds and grasses grow differently, EH-1422 Herbicide kills broadleaf weeds without harming lawns. EH-1422 Herbicide singles out weeds, enters through leaves and moves inside the weed to the roots.

WEEDS CONTROLLED (icon)

White heath Aster, Bushy Aster, Heath Aster, Purple Aster, White prairie Aster, Catchweed Bedstraw, Marsh Bedstraw, Northern Bedstraw, Hairy Beggarticks, Creeping Beggarweed, Threeflower Beggarweed, Florida Bellflower, Florida Betony, Field Bindweed, Hairy Bittercress, Whitehead Broom, Burclover, Common Burdock, Great Burdock, Lawn Burweed, Smallflower Buttercup, Hairy Buttercup, Bushy Buttonweed, Virginia Buttonweed, Black Medic, Carolina Geranium, Carpetweed, Wild Carrot, Cat's-ear, Smooth Chaff-flower, Chicory, Common Chickweed, Mouseeared Chickweed, Sticky Chickweed, Chicory, Common Cinquefoil, Oldfield Cinquefoil, Alsike clover, Crimson Clover, Hop Clover,

White Clover, Wandering Cudweed, Purple Cudweed, Shiny Cudweed, Cupids Shaving Brush, English Daisy, Oxeye Daisy, Common Dandelion, False Dandelion, Catsear Dandelion, Dayflowers, Purple Deadnettle, Dichondra, Dogfennel, Doveweed, Broadleaf Dock, Curly Dock, Dollarweed, Doveweed, Heartleaf Drymary, Eclipta, Facelis, Oakleaf Fleabane, Rough Fleabane, Wild Garlic, Carolina Geranium, Clammy Ground Cherry, Smooth Ground Cherry, Ground Ivy, Groundsel, Yellow Hawkweed, Healall, Henbit, Sprawling Horseweed, innocence (blue eyed Mary), Knawel, Prostrate Knotweed, Tufted Knotweed, Kochia, Lambsquarters, Lespedeza, Field Madder, Alkali Mallow, Venice Mallow, Common Mallow, Little Mallow, Dwarf Mallow, Bristly Mallow, Marcela (yellow joyweed), Marestail, Horseweed, Mat Lippia, Milkweed, Moneywort, Mugwort, Common Mullein, Moth Mullein, Mexican Morningglory, Bigroot Morningglory, Ivyleaf Morningglory, Tall Morningglory, Red Morningglory, Ball Mustard, Black Mustard, Blue Mustard, Indian Mustard, Tumble Mustard, Wild Mustard, Tansy Mustard, White Mustard, Wormseed Mustard, Bitter Nightshade, Eastern Black Nightshade, Silverleaf Nightshade, Cutleaf Nightshade, Hairy Nightshade, Wild Onion, Old World Diamond Flower, Birdseye Pearlwort, Florida Pellatory, Pennywort, Asiatic Pennywort, Peppergrass, Virginia Peppered, Prostrate Pigweed, Redroot Pigweed, Smooth Pigweed, Tumble Pigweed, Pineappleweed, Broadleaf Plantain, Buckhorn Plantain, Blackseeded Plantain, Poison Ivy, Poison Oak, Cutleaf Evening Primrose, Showy Evening Primrose, Purple cudweed, Purslane, Brazil Pusley, Largeflower Pusley, Florida Pusley, Giant Ragweed, Common Ragweed, Western Ragweed, Lanceleaf Ragweed, Redweed, Yellow Rocket, Sibara, Red Sorrel, Sheep Sorrel, Shepherdspurse, Pennsylvania Smartweed, Swamp Smartweed, Ladysthumb Smartweed, Pale Smartweed, Bitter Sneezeweed, Bibbed Speedwell, Common Speedwell, Corn Speedwell, Ivyleaf Speedwell, Persian Speedwell, Purslane Speedwell, Slender Speedwell, Snow Speedwell, Thymeleaf Speedwell, Winter Speedwell, Common Spiderwort, Spotted Spurge, Prostrate Spurge, Corn Spurry, Swinecress, Bull Thistle, Canada Thistle, Cotton Thistle, Distaff Thistle, Drummonds Thistle, Elk Thistle, Gray Thistle, Italian Thistle, Leafy Thistle, Meadow Thistle, Milk Thistle, Platte Thistle, Plumeless Thistle, Russian Thistle, Scotch Thistle, Slenderflower Thistle, Tall Thistle, Wavyleaf Thistle, Yellowspine Thistle, Stiff Verbena, Veronica, Common Vetch, Hairy Vetch, Winter Vetch, Woolly Vetch, Violet, Field Violet, Wild Carrot, Wild Lettuce, Woodsorrel, Creeping Woodsorrel, Yellow Woodsorrel, Florida Yellow Woodsorrel, Woollyleaf Bursage, Common Yarrow, Western Yarrow, Yellow Yarrow.

[REFILL INSTRUCTIONS FOR EH-1424 HERBICIDE PACKAGED WITH PULL'N SPRAY BOTTLE]

- a. To refill only an empty 1.33 GAL: [EH-1424 Herbicide Container]
Amount to Use: Add 8 fl. oz. of this product to the empty 1.33 gallon container. Slowly fill with 160 fl. oz. of clean tap water.
- b. To refill only an empty 1 GAL.: [EH-1424 Herbicide Container]
Amount to Use: Add 6 fl. oz. of this product to the empty 1 gallon container. Slowly fill with 120 fl. oz. of clean tap water.
- c. Specially formulated for residential use.
 Equal Measures: 1 Tablespoon (Tbs) = 3 teaspoons (tsp) 1 fl oz = 2 Tbs

III. Instructions for hose-end sprayers including READY-SPRAY® with a package size of 32 fl. oz. (1.0 quart).

WATER BASED

OPEN

Resealable Label for Directions & Precautions

PRESS TO RESEAL

Ready-Spray makes lawn and garden care easy. Simply attach the hose to the attached sprayer.

Eliminates mixing, measuring and messy clean up. Automatically mixes the right amount of product and water.

OR

Kills weeds in your lawn - won't harm lawn grasses. EH 1422 Herbicide singles out weeds enters through leaves and moves inside the weed to the roots.

Makes lawn weed control easy. Simply connect hose to attached sprayer. Eliminates mixing, measuring and messy clean-up.

Eliminates mixing, measuring and messy clean-up.

<p>PRODUCT FACTS: Treats up to 10,666 sq ft (approx. size of 5 tennis courts) (32 oz size)</p> <p>KILLS WEEDS: Dandelion, Clover, Chickweed plus over 80 additional broadleaf weeds [illustration].</p> <p>WHERE TO USE: ON LAWNS [without harming lawn grasses]</p> <p>AMOUNT TO SPRAY: [For Sensitive Lawns] Bentgrass, Buffalograss, Centipedegrass & St. Augustinegrass (EXCEPT 'Floratom') (Do not use on 'Floratom in Florida): 8 fl.oz. (1/4 bottle) treats 2,666 sq.ft.</p> <p>[For Hardy Lawns] Bahagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass: 8 fl.oz. (1/4 bottle) treats 1,333 sq.ft.</p> <p>QUESTIONS, COMMENTS OR MEDICAL INFORMATION: Call 1 -800-xxx-xxxx http://www.companyname.com</p> <p>Specially formulated for residential use.</p>
--

AMOUNT TO SPRAY

- Measure lawn. Calculate square feet by multiplying length times width. [walking man icon]
- Use the amount indicated for your lawn variety:
[FOR SENSITIVE LAWNS:]: Bentgrass, Buffalograss, Centipedegrass & St. Augustinegrass (EXCEPT 'Floratom', a variety of St. Augustinegrass widely grown in Florida.): 8 fl.oz. (1/4 bottle) treats 2,666 sq.ft.
[FOR HARDY LAWNS:]: Bahagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass: 8 fl.oz. (1/4 bottle) treats 1,333 sq.ft.

HOW TO USE THE READY-SPRAY®

CONNECT: Connect sprayer to hose. Be sure on/off switch is set to "OFF". Turn on water. [illustration]

- Use coin or key to rotate small (color) button to "OPEN". [illustration]

SPRAY: To BEGIN spraying, point nozzle towards lawn. Turn on/off switch forward to "ON". Spray evenly over measured area.

FINISH : To STOP spraying, turn on/off switch back to "OFF". Rotate small (color) button to "CLOSED". Turn off water.

- Relieve water pressure by turning switch to "ON" until water slows to a drip. Turn switch back to "OFF". Disconnect from hose.

[illustration]

FOR BEST RESULTS

MOW: Do not mow for 1 to 2 days after application.

[Illustration/ mower]

WATER:

- If soil is dry, water 1 to 2 days before application.
- Rain or watering 6 hours after application will not wash away effectiveness.

WHEN TO APPLY:

- **SPRAY WHEN:**
 - weeds are small and actively growing
 - temperature is below 90°F
 - air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants
 - Reapplication can be made in 2 to 3 weeks if necessary.
 - Newly seeded lawn can be sprayed after first mowing.
 - Bare spots can be planted with grass seed 1 week after application.
- [illustration/ weather]

OR

WHEN TO [MOW & - optional] WATER]

- Do not mow for 1 to 2 days before or after application.
- If soil is dry, water 1 to 2 days before application.
- Rain or watering 6 hours after application will not wash away effectiveness.

WHEN TO APPLY [icon]

- Spray when weeds are small and actively growing.
- [Spray when] temperature is below 90°F.
- [Spray when] air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees and other desirable plants
- Reapplication can be made in 2 to 3 weeks if necessary.
- Newly seeded lawn can be sprayed after first mowing.
- Bare spots can be planted with grass seed 1 week after application.

IMPORTANT

- If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots.
- St. Augustinegrass, centipedegrass or bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45°F within 72 hours of application.
- This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass if misapplication occurs.
- Do not spray carpetgrass, dichondra or desirable clovers.
- Do not use on 'Floritam' St. Augustinegrass in Florida.

Other Use Precautions: Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

Limitations on 2,4-D applications to residential turfgrass

The maximum application rate is 6.0 fl.oz. of product per 1,000 sq.ft. per application (0.78 lbs 2,4-D ae, 0.41 lb MCP-p ae and 0.14 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 12.0 fl.oz. of product per 1,000 sq.ft. (1.56 lbs 2,4-D ae, 0.82 lb MCP-p ae and 0.28 lb Dicamba ae per acre).

HOW IT WORKS

Because broadleaf weeds and grasses grow differently, EH-1422 Herbicide® kills broadleaf weeds without harming lawns. EH-1422 Herbicide singles out weeds enters through leaves and moves inside the weed to the roots.

WEEDS

Annual Aster, Bedstraw, Beggarweed, Bindweed, Bittercress, Black Medic, Brassbuttons, Broadleaf Plantain, Buckhorn, Bull Thistle, Burdock, Bur Clover, Burweed, Buttercup, Buttonweed, Carolina Geranium, Carpetweed, Catsear, Chicory, Chickweed, Cinquefoil, Clover, Cocklebur, Common Chickweed, Cudweed, Curly Dock, Dandelion, Dock, English Daisy, False Dandelion, Florida Pusley, Galinsoga, Geranium, Ground Ivy, Hairy Fleabane, Hawkweed, Healall, Henbit, Knawel, Knotweed, Lambsquarters, Lespedeza, Mallow, Matchweed, Morningglory, Mouseear Chickweed, Musk Thistle, Mustard, Narrowleaf Plantain, Oxalis, Pennsylvania Smartweed, Pennywort (Creeping Charlie), Peppergrass, Pigweed, Piney Woods Bedstraw, Plantain, Poison Ivy, Poison Oak, Prickly Lettuce, Prostrate Spurge, Puncture Vine, Purslane, Ragweed, Red Clover, Red Henbit, Scarlet Pimpernel, Sheep Sorrel, Shepherdspurse, Sowthistle, Speedwell, Spotted Spurge, Spurge, Tansy Mustard, Thistle, Velvetleaf, White Clover, Wild Carrot, Wild Garlic, Wild Lettuce, Wild Onion, Wild Strawberry, Yarrow & other broadleaf weeds

IV. TABULAR FORMAT

Table 1.

Lawn Grass	Sprayer Type	Product To Measure	Area to Treat	Comments
St. Augustinegrass*, Bentgrass, Buffalograss, and Centipede grass	Tank Sprayer	1 fl.oz. (2Tbs.) into 1 gallon of water and spray.	333 sq.ft.	*Do not use on 'Floritam' St. Augustinegrass in Florida
	Dial 'N Spray®	3 fl.ozs. (6 Tbs.) into sprayer jar. DO NOT ADD WATER. Set dial on 1 oz. and spray.	1,000 sq.ft.	Do not apply to St. Augustinegrass and centipede grass emerging from winter dormancy;
	Ready Spray®	32 fl.oz, entire container.	10, 666 sq.ft.	Temperatures below 45°F within 72 hours of application may cause damage

Table 2.

Lawn Grass	Sprayer Type	Product to Measure	Area To Treat
Perennial Ryegrass, Bermudagrass, Bahiagrass, and Zoysiagrass	Tank Sprayer:	2 fl.oz. (4 Tbs.) into 1 gallon of water and spray.	333 sq.ft.
	Dial 'N Spray®	6 fl.ozs. (12 Tbs.) into sprayer jar. DO NOT ADD WATER. Set dial on 2 oz. and spray.	1,000 sq.ft.
	Ready Spray®	32 fl.oz, entire container.	5,333 sq.ft.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Made in USA
- Kills Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf lawn weeds.
- Kills weeds - not lawn grasses.
- Kills weeds — not lawns
- For Lawns Containing Kentucky Bluegrass, Fescues, Bentgrass, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass, Zoysiagrass, St. Augustinegrass, and Buffalograss.
- Kills broadleaf weeds - roots and all.
- For northern and southern grasses.
- Won't harm lawn grasses shown on container label.
- Won't harm lawn grasses
- Kills weeds in your lawn.
- Formulated for use on northern and sensitive southern lawns.
- Kills major broadleaf lawn weeds: List of the weed species.
- Helps provide a beautiful lawn.
- For cool and warm season lawn grasses
- (Brand Name) kills major broadleaf lawn weeds including dandelion, chickweed, knotweed, in lawns, without harming lawn grasses.
- For use on grass lawns especially Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bentgrass, Bermudagrass, Bahiagrass, Centipedegrass, Zoysiagrass, St. Augustinegrass, and Buffalograss.
- A Trimec® Turf Herbicide Formulation.
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf lawn weeds including those shown on the side panel.
- One Quart Covers Up To 10,666 Square Feet

- Now for northern and southern lawns.
- Kills over 2 times/twice as many weeds.
- Water based.
- Low odor formula.
- Low odor.
- Kills major broadleaf weeds including Dandelion, Clover, Oxalis, Chickweed, Plantain, Henbit, Spurge and others.
- Concentrate
- Now faster results
- Now fast acting
- For control that doesn't/won't wash away/off.
- Effective even with rain in 2 hours.
- Does not harm lawn when used properly
- kills [weeds], not [lawn(s) (grass)].
- Trademark name for the lawn.
- For a beautiful Weed-Free Lawn
- Kills weeds down to the root.
- Rainproof (rain-fast) in two [hours]
- Kills broadleaf weeds in your lawn.
- Won't harm most lawn grasses.
- Ideal for spot treating individual lawn weeds;
- Rain proof in hours.
- Kills Dandelion, Clover and Other Broadleaf Weeds—Won't Harm Lawns.
- Kills Weeds—Not Lawn Grasses
- Guaranteed
- Won't harm lawns, guaranteed.
- Guaranteed not to harm the lawn.
- Guaranteed to kill weeds, not grass [lawn].
- Starts working on contact
- Fast results
- Fast-acting formula
- Weeds begin wilting overnight
- Weeds begin to wilt and die overnight
- Rainproof in 2 hours — so it [weed killing power] won't wash away.
- Rainproof in 2 hours so you won't have to respray.
- Kills to the root so weeds don't [won't]ever come [grow] back.
- Kills more weeds
- Complete root [kill]
- Kills major [broadleaf weeds]
- Complete kill
- Kills even tough lawn weeds
- Kills over 200 weeds [weeds]
- EH-1422 Herbicide singles out weeds--leaves [grass(es) alone [unharmd]
- For use on lawns
- How it works — EH-1422 Herbicide's exclusive formula specifically targets [seeks out] weeds, killing them, while leaving the grass unharmed — guaranteed.
- How it works — EH-1422 Herbicide contains a special [unique][exclusive] blend of proven ingredients that seek out [target] and destroy [kill] weeds, while leaving the grass [lawn] unharmed — guaranteed.
- Specially formulated to kill weeds, not the lawn.
- Targets weeds, leaves grass [lawn] unharmed.
- Made [formulated] [special formula] for use on lawns.
- New Formula!

- New
- Helps keep the lawn look[ing] beautiful.
- Easy to use
- Ideal for use in between fertilizer applications to spot treat weeds
- Can be applied with hose-end or pump-up (tank) sprayers.
- "One Quart Covers Up To 10,666 Square Feet" or "One Pint Covers Up To 5,333 Square Feet".
- Can be applied with hose-end and tank sprayers.
- Makes up to ____ gallons of spray (Volume dependent upon package size)
- Use with Ortho® Dial-N-Spray®
- Notice: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.
- Guarantee: If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

A partial listing of broadleaf weeds:			
Annual aster	Clover	Mallow	Red clover
Bedstraw	Cocklebur	Matchweed	Red henbit
Beggarweed	Common chickweed	Morningglory	Scarlet pimpernel
Bindweed	Cudweed	Mouseear chickweed	Sheep sorrel
Bittercress	Curly dock	Musk thistle	Shepherdspurge
Black medic	Dandelion	Mustard	Sowthistle
Brassbuttons	Dock	Narrowleaf plantain	Speedwell
Broadleaf plantain	English daisy	Oxalis	Spotted spurge
Buckhorn	False dandelion	Pennsylvania smartweed	Spurge
Bull thistle	Florida pusley	Pennywort(creeping Charlie)	Tansy mustard
Burdock	Galinsoga	Peppergrass	Thistle
Burclover	Geranium	Pigweed	Velvetleaf
Burweed	Ground ivy	Piney woods bedstraw	White clover
Buttercup	Hairy fleabane	Plantain	Wild carrot
Buttonweed	Hawkweed	Poison ivy	Wild garlic
Carolina geranium	Healall	Poison oak	Wild lettuce
Carpetweed	Henbit	Prickly lettuce	Wild onion
Catsear	Knawel	Prostrate spurge	Wild strawberry
Chicory	Knotweed	Puncturevine	Yarrow
Chickweed	Lambsquarters	Purslane	and other
Cinquefoil	Lespedeza	Ragweed	broadleaf weeds

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00892.20080207.doc
2. Previous Version: AP032206
3. Reason for Issue: Various changes pertaining to re-registration. Re-format label.