

2217-894

12/4/2009

10826



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCE

James L. Kunstman, Ph.D.
Director of Regulatory Services
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, Missouri 64101

DEC 4 2009

Subject: Notification(s) for Label Revisions under PRN 2007-4

Dear Dr. Kunstman:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 2007-4 dated October 16, 2009 for:

- | | |
|--|--------------------------|
| EPA Registration 2217-894[✓] | EH 1432 Herbicide |
| EPA Registration 2217-820 | EH-1149 Herbicide |

The Registration Division (RD) has conducted a review of the request(s) for applicability under PRN 2007-4 and finds that the label changes requested fall within the scope of PRN 2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identify the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20210



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 2217-894	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) EH 1432 Herbicide	PM# Product Manager—Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION DEC - 4 2009
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Pesticide Registration Notice (PRN) 2007-4, Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal: Standards for Pesticide Containers and Containment.

Please refer to cover letter for details and certification statement.

e-mail to jkunstman@pbigordon.com FAX: 816-421-2731

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
		If "Yes" Package wgt.	No. per Container	<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1 pint, 1 quart 2 quarts and 1 gallon		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Other _____		
			<input checked="" type="checkbox"/> Paper glued		
			<input type="checkbox"/> Stenciled		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name James L. Kunstman, Ph.D.		Title Director of Regulatory Services		Telephone No. (Include Area Code) 816-460-6292	
Certification					6. Date Application Received (Stamped)
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					
2. Signature 		3. Title Director of Regulatory Services			
4. Typed Name James L. Kunstman, Ph.D.		5. Date October 16, 2009			

30926



An Employee-Owned Company

1217 WEST 12TH STREET • P.O. BOX 014090
KANSAS CITY, MISSOURI 64101-0090
816-421-4070 • 1-800-821-7925
FAX: 816-474-0462

October 16, 2009

Document Processing Desk (PRN)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attn: Ms. Joanne I. Miller (PM-23)

Dear Ms. Miller:

Subject: Labeling notifications according to Pesticide Registration Notices (PRN) Pesticide Registration Notice (PRN) 2007-4 and (PRN) 98-1. EH 1432 Herbicide (EPA Reg. No. 2217-894)

I. Labeling notification per Pesticide Registration Notice (PRN) 2007-4:

April 2, 2008: An Agency letter with one comment stated that the draft labeling for the RED for 2,4-D was acceptable provided that modifications to the Directions for Use were made.

December 8, 2008: An Agency letter stated that this product was re-registered.

October 8, 2009: We have incorporated or merged the revisions for the draft labeling listed in the previous letter into the current document. Now, we ask to revise the disposal instructions and to adopt the exact language of Pesticide Registration Notice (PRN 2007-4), Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal: Standards for Pesticide Containers and Containment."

Please refer to pages 6, 10, 15 and 21 of the draft labeling.

Certification: This notification is consistent with the guidance PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this amended labeling is not consistent with the terms of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

II. Enclosures:

1. Application for Pesticide Amendment (EPA Form 8570-1)
2. One (1) annotated version of the draft labeling and
3. One (1) version of the draft labeling without notations

If you have any questions, please call me at 816-460-6292 or contact me at jkunstman@pbigordon.com.

Sincerely,



James L. Kunstman, Ph.D.
Director of Regulatory Services

Handwritten scribbles and marks at the bottom right of the page.


50926

EH-1432 HERBICIDE

EPA Reg. No. 2217-894

Three sublabels represent the entire master label of EH-1432 Herbicide. Please refer to the appropriate section of the labeling as shown as follows:

Section
Sublabel 1: Intended for all spray equipment used by homeowners
Sublabel 2: Intended for pre-packed hose-end sprayers (also known as ready-sprays)
Sublabel 3: Intended for pump sprayers and hose-end sprayers that require addition of water
Appendix
Document Control Information



60026

SUBLABEL 1

Intended for all spray equipment used by homeowners

EH-1432 HERBICIDE

NOTIFICATION

EPA Reg. No. 2217-894

DEC - 4 2009

FOR OUTDOOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	4.85%
Quinclorac	1.61%
Dicamba, dimethylamine salt	0.45%
OTHER INGREDIENTS:	93.09%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.343 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.03%
- 0.137 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 1.61%
- 0.031 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.37%

Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: **"See back panel for substatement of ingredient statement"**.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. Wear long sleeved shirt and long pants, shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

70926

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING

<p>PRODUCT FACTS:</p> <p>What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass, foxtail, and yellow nutsedge. (See inside for complete list)</p> <p>Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)</p> <p>When to Use: Apply when daytime temperatures are between 50° and 90°F (except on Bermudagrass - see below).</p> <p>Questions & Comments: (icon phone) (icon computer) Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com</p>

HOW IT WORKS:

EH-1432 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

80426

WHERE TO USE:

EH-1432 Herbicide can be applied to the following established turfgrass species:	
Cool season turfgrass	Warm season turfgrass
Kentucky bluegrass, Ryegrass, perennial and annual, Tall fescue, Bentgrass, Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	Bermudagrass ¹ , common and hybrid Zoysiagrass, Buffalograss,
¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.	

DO NOT apply EH-1432 Herbicide to the following:	
Warm season turfgrass	Other areas
St. Augustinegrass, Bahagrass, Centipedegrass, Carpetgrass, Seashore paspalum	Lawns with desirable clovers or legumes, Dichondra, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Turfgrass species that are not listed on this label, Gardens including vegetables, fruit trees, vines and berries.

WEEDS CONTROLLED

GRASS WEEDS:

Crabgrass, large and smooth, foxtail, (green, yellow and giant), kikuyugrass, and signalgrass (broadleaf).

BROADLEAF WEEDS (A partial list is presented below)			
Annual yellow sweetclover	Cockle	Musk thistle	Spiny amaranth
Aster	Cocklebur	Mustard	Spiny cocklebur
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny sowthistle
Bedstraw	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Beggarticks	Cudweed	Nettle	Spotted spurge
Betony, Florida	Curly dock	Orange hawkweed	Spurweed
Bindweed, field	Daisy, English	Oriental cocklebur	Stinging nettle
Bird vetch	Daisy fleabane	Oxalis	Strawberry, India mock
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall nettle
Bittercress, hairy	Dandelion	Parsnip	Tall vervain
Bitterweed	Dichondra	Pearlwort	Tansy ragwort
Black-eyed Susan	Dogbane	Pennycress	Tansy mustard
Black medic	Dogfennel	Pennywort	Tanweed
Black mustard	Dollarweed	Peppergrass	Thistle
Blackseed plantain	Elderberry	Pepperweed	Trailing crownvetch
Blessed thistle	False dandelion	Pigweed	Tumble mustard
Bloodflower milkweed	Falseflax	Pineywoods bedstraw	Tumble pigweed
Blue lettuce	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue vervain	Fiddleneck	Plantain	Venice mallow
Bracted plantain	Florida pusley	Poison ivy	Virginia buttonweed
Brassbuttons	Frenchweed	Poison oak	Virginia creeper
Bristly oxtongue	Galinsoga	Pokeweed	Virginia pepperweed
Broadleaf dock	Goathead	Poorjoe	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prairie sunflower	Western clematis
Broomweed	Ground ivy	Prickly lettuce	Western salsify
Buckhorn	Gumweed	Prickly sida	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild buckwheat
Bull thistle	Healall	Prostrate spurge	Wild carrot
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild four-o'clock
Burclover	Heathaster	Puncturevine	Wild garlic
Burdock	Hedge bindweed	Purslane, common	Wild geranium
Burning nettle	Hedge mustard	Ragweed	Wild lettuce
Burweed			Wild marigold

BROADLEAF WEEDS (A partial list is presented below)			
Buttercup	Hemp	Red sorrel	Wild mustard
Buttonweed	Henbit	Redroot pigweed	Wild onion *
Canada thistle	Hoary cress	Redstem filaree	Wild parsnip
Carolina geranium	Hoary plantain	Rough cinquefoil	Wild radish
Carpetweed	Hoary vervain	Rough fleabane	Wild rape
Catchweed bedstraw	Horsenettle	Russian pigweed	Wild strawberry
Catnip	Jimsonweed	Russian thistle	Wild sweet potato
Catsear	Knawel	Scarlet pimpernel	Wild vetch
Chickweed, common	Knotweed	Scotch thistle	Wild violet
Chickweed, mouseear	Kochia	Sheep sorrel	Woodsorrel
Chicory	Lambsquarters*	Shepherdspurse	Woolly croton
Cinquefoil	Lespedeza	Slender plantain	Woolly morningglory
Clover, crimson	Mallow	Smallflower galinsoga	Woolly plantain
Clover, hop	Matchweed	Smooth dock	Wormseed
Clover, red	Mexicanweed	Smooth pigweed	Yarrow *
Clover, strawberry	Milk vetch	Sorrel	Yellow rocket
Clover, sweet	Morningglory	Sowthistle	Yellowflower pepperweed
Clover, white	Mouseear hawkweed	Spanishneedles	and other broadleaf
	Mugwort	Speedwell	weeds

WHEN TO USE:

Apply this product when weeds are small and actively growing.

CRABGRASS (large and smooth):

- **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at the same rate at 14 days later for dense populations.

FOXTAIL (GREEN GIANT AND YELLOW):

- **Spring and early summer:** Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at the same rate at 14 days later for dense populations.

BROADLEAF WEEDS:

- **Spring:** Apply when **dandelion, plantain, buttercup, speedwell, knotweed** and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- **Fall:** Apply when **henbit, chickweed, white clover, ground ivy, lawn burweed** and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

HOW MUCH TO USE:

[Limitations on 2,4-D applications to residential turfgrass]

The maximum application rate is 8 fl.oz. of product per 1,000 sq.ft. per application (0.93 lb 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. The maximum seasonal rate is 16 fl.oz. of product per 1,000 sq.ft. (1.87 lb 2,4-D acid equivalent per acre), excluding spot treatments. Do not apply more than 2 applications per year.

1. ENTIRE LAWN TREATMENTS

THREE EASY STEPS TO MIX AND APPLY:

1. Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
2. For hose-end sprayers: Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying. The RATE PER GALLON for this product is 4 fl.oz. of product per gallon. See instructions below.
For hand-operated sprayers (backpack sprayers, compression (pump-up) sprayers, knapsack sprayers or tank type): The RATE PER 1,000 SQ.FT. for this product is 8 fl.oz. of product per 1,000 sq.ft. of lawn. Measure and mix the proper amount of product with water as described below.

100226

3. Spray evenly over the lawn.

For Ortho® Dial'N Spray® Hose-End Sprayer, Hudson® Hose End Liquid Concentrate Sprayer or Gilmour® Dial-a-Mix® Lawn and Garden Sprayer:

Set the dial—

For Ortho® Dial'N Spray® Hose-End Sprayer: Set the dial on 4 oz as the RATE PER GALLON and keep the dial set to this setting while spraying.

Hudson® Hose End Liquid Concentrate Sprayer: Set the dial on 4 oz as the RATE PER GALLON and keep the dial set to this setting while spraying.

Gilmour® Dial-A-Mix® Lawn and Garden Sprayer: Set the dial on 4 oz or 8 TBS or 24 TSP as the RATE PER GALLON and keep the dial set to this setting while spraying.

Fill - Determine the total lawn area to be sprayed.

The application rate for this product is 8 fl.oz. of product per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 8 fl.oz. of product for every 1,000 sq.ft. of lawn area.

Do not add water to the sprayer jar.

Spray - Spray evenly over the lawn.

Example:

Set the dial - Set the dial on 4 oz.

Fill - For a lawn area of 3,000 sq.ft, pour or measure 24 fl.oz of product into sprayer jar. Do not add water.

Spray - Spray uniformly on 3,000 sq.ft. of lawn.

For READY-TO-SPRAY Hose End Sprayers:

This retail sprayer bottle will cover the treated areas described below when this product is applied at 8 fl.oz.of product per 1,000 sq.ft.

Approximate treated areas for four retail container sizes:

Species	Amount of product (to treat approximate area, sq.ft.)				
	fl.oz./1,000 sq.ft.	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container	2.0 qt. (64 fl.oz.) retail container	1 gallon (128 fl.oz.) retail container
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season species Bermudagrass ¹ , common and hybrid Zoysiagrass and Buffalograss, Bentgrass	8 fl.oz./1,000 sq.ft.	2,000 sq.ft.	4,000 sq.ft.	8,000 sq.ft.	16,000 sq.ft.

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Instructions:

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet. Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl.oz. container will treat up to xxxx sq.ft. of turfgrass.
- **FINISH:** To stop spraying, turn [rotate] spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn [rotate] spray nozzle to the "ON" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.


[OR]

CONNECT

1. Shake well before use.
2. Connect sprayer to garden hose.
3. Turn on water.

SPRAY

1. Extend hose to farthest area to be treated, point nozzle in direction to be sprayed and turn on sprayer. Work back toward the faucet to avoid walking over treated grass.
2. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas.



[Operating the Sprayer Top]


Water control knob (closest to hose). < Graphic > Twist counterclockwise [clockwise] for 'On' & clockwise [counter-clockwise] for 'Off'.

Weed killer control knob. < Graphic > Twist knob until flat side is next to protruding plastic tab. This will release the safety feature. Push in for 'On' & out for 'Off'.]

FINISH

1. Turn off sprayer.
2. Turn off water at faucet.
3. Relieve pressure in hose by turning water back on at sprayer until water slows to a drip. Close sprayer. Disconnect sprayer from hose.

[OR]

<p>HOW TO USE</p> <p>Shake Well Before Use</p>	<p>CONNECT</p> <ol style="list-style-type: none"> 1. Shake well before use. 2. Connect sprayer to garden hose. 3. Turn on water. <p>SPRAY</p> <ol style="list-style-type: none"> 1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed. 2. Begin spraying – Push & hold lock tab away from knob, use other hand to turn knob clockwise [counter-clockwise] to ON position. 3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.  <p>FINISH</p> <ol style="list-style-type: none"> 1. To stop spraying, push & hold lock tab away from knob and use other hand to turn knob [clockwise] counter-clockwise to OFF position. 2. Turn off water at faucet. 3. Relieve water pressure in hose by turning sprayer back ON until water slows to a drip, then turn sprayer OFF. [-or-] Relieve water pressure in hose by turning sprayer to WATER position until water slows to a drip, then turn sprayer to OFF position. 4. Disconnect sprayer from hose.
---	---

2. SPOT TREATMENTS FOR SMALL AREAS OR INDIVIDUAL WEEDS:

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up): Mix 4 fl.oz. of product with 1 gallon of water. Note: Approximately one (1) gallon of spray solution will treat 500 sq.ft. See below for additional spray preparations. Shake well before spraying. Apply to actively growing weeds during the early growth stages. Spray coverage should be uniform and complete.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square feet
4 fl.oz.	1 gallon	500 sq.ft.
6 fl.oz.	1.5 gallons	750 sq.ft.
8 fl.oz.	2 gallons	1,000 sq.ft.
12 fl.oz.	3 gallons	1,500 sq.ft.

Equal measures: 1 cup = 8 fl.oz. = 1/2 pint. 4 fl.oz. = 8 Tablespoons

[OR]

Measure indicated amount of product and pour into tank. Dilute with water according to the following chart. Adjust the nozzle to deliver coarse spray droplets and treat the area indicated.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Area To Treat, Square feet
4 fl.oz.	1 gallon	500 sq.ft.
8 fl.oz.	2 gallons	1,000 sq.ft.
12 fl.oz.	3 gallons	1,500 sq.ft.

Equal measures: 1 cup = 8 fl.oz. = 1/2 pint. 4 fl.oz. = 8 Tablespoons

CULTURAL TIPS:

For newly seeded areas:

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas:

- The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Adequate moisture and mild environmental conditions enhance the performance of EH-1432 Herbicide. If dry conditions exist, schedule irrigations (watering) at 24 hours before and after the application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

PROHIBITIONS:

- Apply when daytime temperatures are between 50° and 90°F. Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur. For bermudagrass lawns do not apply when daytime temperatures exceed 85° F.
- Application at rates above specified on this label may cause injury to turfgrass.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

<p>STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage and disposal.</p> <p>PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.</p> <p>PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>
--

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of

14 of 26

purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere

SUBLABEL 2

Intended for pre-packed hose-end sprayers (also known as ready-sprays)

EH-1432 HERBICIDE

EPA Reg. No. 2217-894

**(Pre-packed Hose-end Sprayer Label)
(Front Panel)**

Editorial Note – Marketing claims for the retail container from the Appendix may be positioned here.

ACTIVE INGREDIENT:	
2,4-D, dimethylamine salt	4.85%
Quinclorac	1.61%
Dicamba, dimethylamine salt.....	0.45%
OTHER INGREDIENTS:	<u>93.09%</u>
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Editorial Note – [Bracketed text] is optional language



(Back Panel)

OPEN Resealable Label for Directions & Precautions

PRESS TO RESEAL

EH-1432 Herbicide's exclusive formula kills lawn weeds plus crabgrass in one easy step. And it will not harm your lawn when used as directed. Convenient sprayer top connects to garden hose and automatically dispenses the weed killer. Ideal for large areas – treats 5,000 [4,000] square feet. Rainproof in one hour. No messy cleanup. Controlling lawn weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS	
Editorial Note – Marketing Claims Positioned Here	
KILLS	Broadleaf Weeds: Dandelion, Chickweed, and Clovers Grassy Weeds: Crabgrass and over 200 other weeds.
WHERE TO USE	ON LAWNS without harming lawn grasses – Bermudagrass ¹ , Buffalograss, Kentucky Bluegrass, Ryegrass, Fescue, Zoysia & Bentgrass ¹ Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. (Do not use on Bahiagrass, Carpetgrass, Centipedegrass, St. Augustine [including 'Floritam'], or Dichondra.)
COVERAGE	This bottle treats an average size lawn (5,000 [4,000] sq.ft.). Just hook bottle to hose and you are ready.
 	For questions or comments, call toll-free 1-800-XXX-YYYY or visit our website at www.CompanyName.com


THIS PRODUCT CONTAINS: 4.03% 2,4-D acid equivalent, 1.61% Quinclorac acid, 0.37% Dicamba acid equivalent. Isomer Specific by AOAC Methods.

FOR OUTDOOR RESIDENTIAL USE ONLY

EPA Reg. No. 2217-894
EPA Est. No. 2217-KS-1

DIRECTIONS FOR USE	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.	
HOW MUCH TO USE	Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. This bottle will treat 5,000 sq.ft. in about 20 minutes (actual time will vary depending on water pressure), Editorial Note – Final time value will be adjusted based on final hose-end sprayer selected for market. The maximum application rate is 8 fl.oz. of product per 1,000 sq.ft. per application (0.93 lb 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 applications per year.


170926

<p>HOW TO USE</p> <p>Shake Well Before Use</p>	<p>CONNECT</p> <ol style="list-style-type: none">1. Shake well before use.2. Connect sprayer to garden hose.3. Turn on water. <p>SPRAY</p> <ol style="list-style-type: none">1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.2. Begin spraying – Push & hold lock tab away from knob, use other hand to turn knob clockwise [counter-clockwise] to ON position.3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.  <p>FINISH</p> <ol style="list-style-type: none">1. To stop spraying, push & hold lock tab away from knob and use other hand to turn knob [clockwise] counter-clockwise to OFF position.2. Turn off water at faucet.3. Relieve water pressure in hose by turning sprayer back ON until water slows to a drip then turn sprayer OFF. [-or-] Relieve water pressure in hose by turning sprayer to WATER position until water slows to a drip, then turn sprayer to OFF position.4. Disconnect sprayer from hose.
<p>FOR BEST RESULTS</p>	<p>WHEN TO APPLY</p> <ul style="list-style-type: none">• Spray when<ul style="list-style-type: none">– weeds are small and actively growing– temperature is below 90°F. Do not apply to bermudagrass lawns when daytime temperatures exceed 85° F.• Certain hard-to-kill weeds may be retreated in 2 weeks if necessary. If more than 2 applications are needed, use a spot treatment lawn weed killer such as [the alternate brand name of Herbicide 3D RTU (72155-1)].• Do not apply while grass is emerging from dormancy.• Newly seeded lawn may be sprayed after fourth mowing.• Bare spots may be seeded 4 weeks after application.• Keep product spray off plants growing in or around the lawn such as trees, shrubs, flowers, perennials, groundcovers, vegetables, & herbs. <p>MOWING</p> <ul style="list-style-type: none">• Do not mow for 1 to 2 days after spraying• Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens. <p>WATERING</p> <ul style="list-style-type: none">• If lawn is dry, water 1 to 2 days before using this product.• Rain or watering 3 hours after application will not wash away effectiveness.

HOW IT WORKS	EH-1432 Herbicide contains three proven weed killers to control lawn weeds & crabgrass down to the roots without harming your lawn when used as directed.
---------------------	---

WEEDS

Annual Yellow Sweetclover, Aster, Austrian Fieldcress, Barnyardgrass, Bedstraw, Beggarticks, Florida Betony, Field Bindweed, Bird Vetch, Bitter Wintercress, Hairy Bittercress, Bitterweed, Black-eyed Susan, Black Medic, Black Mustard, Blackseed Plantain, Blessed Thistle, Bloodflower Milkweed, Blue Lettuce, Blue Vervain, Bracted Plantain, Brassbuttons, Bristly Oxtongue, Broadleaf Dock, Broadleaf Plantain, Broadleaf Signalgrass, Broomweed, Buckhorn Plantain, Bulbous Buttercup, Bull Thistle, Bullnettle, Burclover, Burdock, Burning Nettle, Burweed, Buttercup, Buttonweed, Canada Thistle, Carolina Geranium, Carpetweed, Catchweed Bedstraw, Catnip, Catsear, Common Chickweed, Mouseear Chickweed, Chicory, Cinquefoil, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White Clover, Cockle, Cocklebur, Common Mullein, Creeping Charlie, Creeping Jenny, Cudweed, Curly Dock, English Daisy, Fleabane Daisy, Oxeye Daisy, Dandelion, Dichondra, Dogbane, Dogfennel, Dollarweed, Elderberry, False Dandelion, False Flax, False Sunflower, Fiddleneck, Florida Pusley, Frenchweed, Galinsoga, Giant Foxtail, Goathead, Goldenrod, Green Foxtail, Ground Ivy, Gumweed, Hairy Fleabane, Hawkweed, Healall, Drymary Heartleaf, Heath Aster, Hedge Bindweed, Hedge Mustard, Hemp, Henbit, Hoary Cress, Hoary Plantain, Hoary Vervain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Large Crabgrass, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk Vetch, Morningglory, Mouseear Hawkweed, Mugwort, Musk Thistle, Mustard, Narrowleaf Plantain, Narrowleaf Vetch, Nettle, Orange Hawkweed, Oriental Cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods Bedstraw, Plains Coreopsis (Tickseed), Plantain, Poison Ivy, Poison Oak, Pokeweed, Poorjoe, Prairie Sunflower, Prickly Lettuce, Prickly Sida, Prostrate Knotweed, Prostrate Pigweed, Prostrate Spurge, Prostrate Vervain, Puncturevine, Common Purslane, Ragweed, Red Sorrel, Redroot Pigweed, Redstem Filaree, Rough Cinquefoil, Rough Fleabane, Russian Pigweed, Russian Thistle, Scarlet Pimpernel, Scotch Thistle, Sheep Sorrel, Shepherdspurse, Slender Plantain, Smallflower Galinsoga, Smooth Crabgrass, Smooth Dock, Smooth Pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny Amaranth, Spiny Cocklebur, Spiny Sowthistle, Spotted Catsear, Spotted Spurge, Spurweed, Stinging Nettle, India Mock Strawberry, Tall Nettle, Tall Vervain, Tansy Mustard, Tansy Ragwort, Tanweed, Thistle, Trailing Crownvetch, Tumble Mustard, Tumble Pigweed, Velvetleaf, Venice Mallow, Virginia Buttonweed, Virginia Creeper, Virginia Pepperweed, Wavyleaf Bullthistle, Western Clematis, Western Salsify, White Mustard, Wild Aster, Wild Buckwheat, Wild Carrot, Wild Four-o'clock, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Marigold, Wild Mustard, Wild Onion, Wild Parsnip, Wild Radish, Wild Rape, Wild Strawberry, Wild Sweet Potato, Wild Vetch, Wild Violet, Woodsorrel, Woolly Croton, Woolly Morningglory, Woolly Plantain, Wormseed, Yarrow, Yellow Foxtail, Yellowflower Pepperweed, Yellow Rocket, and other broadleaf weeds

	<ul style="list-style-type: none"> Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
---	--


PRECAUTIONARY STATEMENTS

CAUTION
Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. Wear long sleeved shirt and long pants, shoes and socks.
 Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID +	
If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
<p>NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.</p> <p>Active Ingredients: 4.85% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.45% Dicamba, dimethylamine salt (CAS # 2300-66-5), and 1.61% Quinclorac (CAS # 84087-1-4).</p>	

Environmental Hazards
<p>This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.</p> <p>This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.</p>

Important – Directions for Storage and Disposal	
Do not contaminate water, food, or feed by storage and disposal.	
Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
Pesticide Disposal and Container Handling 	Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D-9848 072205

200926

SUBLABEL 3

**Intended for pump sprayers and hose-end sprayers that
require addition of water**

EH-1432 HERBICIDE

EPA Reg. No. 2217-894

**(Pump Sprayer / Hose-end Sprayer Label)
(Front Panel)**

Editorial Note – Marketing claims for the retail container may be positioned here.

ACTIVE INGREDIENT:	
2,4-D, dimethylamine salt	4.85%
Quinclorac	1.61%
Dicamba, dimethylamine salt.....	0.45%
OTHER INGREDIENTS:	<u>93.09%</u>
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Editorial Note – [Bracketed text] is optional language


(Back Panel)

OPEN Resealable Label for Directions & Precautions

PRESS TO RESEAL

EH-1432 Herbicide's exclusive formula kills lawn weeds plus crabgrass in one easy step. And it will not harm your lawn when used as directed. Ideal for large areas – treats 5,000 [4,000] square feet. Rainproof in one hour. Controlling lawn weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS	
Editorial Note – Marketing Claims Positioned Here	
KILLS	Broadleaf Weeds: Dandelion, Chickweed, and Clovers Grassy Weeds: Crabgrass and over 200 other weeds.
WHERE TO USE	ON LAWNS without harming lawn grasses – Bermudagrass ¹ , Buffalograss, Kentucky Bluegrass, Ryegrass, Fescue, Zoysia & Bentgrass ¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. (Do not use on Bahiagrass, Carpetgrass, Centipedegrass, St. Augustine [including 'Floritam'], or Dichondra.)
COVERAGE	This bottle treats an average size lawn (5,000 sq.ft. [4,000])
	For questions or comments, call toll-free 1-800-XXX-YYYY or visit our website at www.CompanyName.com

THIS PRODUCT CONTAINS: 4.03% 2,4-D acid equivalent, 1.61% Quinclorac acid, 0.37% Dicamba acid equivalent. Isomer Specific by AOAC Methods.

FOR OUTDOOR RESIDENTIAL USE ONLY

EPA Reg. No. 2217-894
EPA Est. No. 2217-KS-1


<u>DIRECTIONS FOR USE</u>	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.	
HOW MUCH TO USE	Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. 3.2 fl.oz. (6.5 TBS) [4 fl.oz. (8 TBS)] of this product in one gallon of water treats 500 sq.ft. Food utensils, such as teaspoons and measuring cups, should not be used for food purposes after use with this herbicide. The maximum application rate is 8 fl.oz. of product per 1,000 sq.ft. per application (0.93 lb 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 applications per year.
HOW TO USE PUMP STYLE TANK SPRAYERS	Shake bottle well before use. Best for spraying small areas (spot treatment) such as individual weeds. <ol style="list-style-type: none"> 1. Add the appropriate amounts of water & this product to the sprayer tank. 2. Close sprayer, shake well and pressurize. 3. Adjust nozzle to a coarse spray pattern. Hold tip of wand about 12" from weed(s), aim at center of weed & spray. 4. Occasionally re-pressurize sprayer if needed to maintain a good spray pattern.

<p>DIAL-STYLE HOSE-END SPRAYERS</p>	<p>Best for spraying large areas (broadcast treatment) such as the entire lawn.</p> <ol style="list-style-type: none"> 1. Fill sprayer jar with enough of this product to cover area to be treated. DO NOT ADD WATER. 2. Close sprayer and set dial. 3. Attach sprayer to hose and apply. 4. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.
<p>CONCENTRATE PLUS WATER HOSE-END SPRAYERS</p>	<p>Best for spraying large areas (broadcast treatment) such as the entire lawn.</p> <ol style="list-style-type: none"> 1. Pour this product into sprayer jar to the required 'Fluid Ounce' level. 2. Add the appropriate amount of water to the 'Gallons' level. 3. Replace sprayer top and shake well before applying. 4. Attach sprayer to garden hose and apply. 5. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.

<p>FOR BEST RESULTS</p>	<p>WHEN TO APPLY</p> <ul style="list-style-type: none"> • Spray when <ul style="list-style-type: none"> - weeds are small and actively growing - temperature is below 90°F Do not apply to bermudagrass lawns when daytime temperatures exceed 85° F. ▪ Certain hard-to-kill weeds may be retreated in 2 weeks if necessary. If more than 2 applications are needed, spot treat individual weeds using this product in a tank sprayer. ▪ Do not apply while grass is emerging from dormancy. ▪ Newly seeded lawn may be sprayed after fourth mowing. ▪ Bare spots may be seeded 4 weeks after application. ▪ Keep product spray & drift off plants growing in or around the lawn such as trees, shrubs, flowers, perennials, groundcovers, vegetables, & herbs. <p>MOWING</p> <ul style="list-style-type: none"> • Do not mow for 1 to 2 days after spraying. • Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens. <p>WATERING</p> <ul style="list-style-type: none"> • If lawn is dry, water 1 to 2 days before using this product. • Rain or watering 3 hours after application will not wash away effectiveness.
<p>HOW IT WORKS</p>	<p>EH-1432 Herbicide contains three proven weed killers to control lawn weeds & crabgrass down to the roots without harming your lawn when used as directed.</p>

<p>WEEDS</p>
<p>Annual Yellow Sweetclover, Aster, Austrian Fieldcress, Barnyardgrass, Bedstraw, Beggarticks, Florida Betony, Field Bindweed, Bird Vetch, Bitter Wintercress, Hairy Bittercress, Bitterweed, Black-eyed Susan, Black Medic, Black Mustard, Blackseed Plantain, Blessed Thistle, Bloodflower Milkweed, Blue Lettuce, Blue Vervain, Bracted Plantain, Brassbuttons, Bristly Oxtongue, Broadleaf Dock, Broadleaf Plantain, Broadleaf Signalgrass, Broomweed, Buckhorn Plantain, Bulbous Buttercup, Bull Thistle, Bullnettle, Burclover, Burdock, Burning Nettle, Burweed, Buttercup, Buttonweed, Canada Thistle, Carolina Geranium, Carpetweed, Catchweed Bedstraw, Catnip, Catsear, Common Chickweed, Mouseear Chickweed, Chicory, Cinquefoil, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White Clover, Cockle, Cocklebur, Common Mullein, Creeping Charlie, Creeping Jenny, Cudweed, Curly Dock, English Daisy, Fleabane Daisy, Oxeye Daisy, Dandelion, Dichondra ,</p>

<p>WEEDS</p> <p>Dogbane, Dogfennel, Dollarweed, Elderberry, False Dandelion, False Flax, False Sunflower, Fiddleneck, Florida Pusley, Frenchweed, Galinsoga, Giant Foxtail, Goathead, Goldenrod, Green Foxtail, Ground Ivy, Gumweed, Hairy Fleabane, Hawkweed, Healall, Drymary Heartleaf, Heath Aster, Hedge Bindweed, Hedge Mustard, Hemp, Henbit, Hoary Cress, Hoary Plantain, Hoary Vervain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Large Crabgrass, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk Vetch, Morningglory, Mouseear Hawkweed, Mugwort, Musk Thistle, Mustard, Narrowleaf Plantain, Narrowleaf Vetch, Nettle, Orange Hawkweed, Oriental Cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods Bedstraw, Plains Coreopsis (Tickseed), Plantain, Poison Ivy, Poison Oak, Pokeweed, Poorjoe, Prairie Sunflower, Prickly Lettuce, Prickly Sida, Prostrate Knotweed, Prostrate Pigweed, Prostrate Spurge, Prostrate Vervain, Puncturevine, Common Purslane, Ragweed, Red Sorrel, Redroot Pigweed, Redstem Filaree, Rough Cinquefoil, Rough Fleabane, Russian Pigweed, Russian Thistle, Scarlet Pimpernel, Scotch Thistle, Sheep Sorrel, Shepherdspurse, Slender Plantain, Smallflower Galinsoga, Smooth Crabgrass, Smooth Dock, Smooth Pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny Amaranth, Spiny Cocklebur, Spiny Sowthistle, Spotted Catsear, Spotted Spurge, Spurweed, Stinging Nettle, India Mock Strawberry, Tall Nettle, Tall Vervain, Tansy Mustard, Tansy Ragwort, Tanweed, Thistle, Trailing Crownvetch, Tumble Mustard, Tumble Pigweed, Velvetleaf, Venice Mallow, Virginia Buttonweed, Virginia Creeper, Virginia Pepperweed, Wavyleaf Bullthistle, Western Clematis, Western Salsify, White Mustard, Wild Aster, Wild Buckwheat, Wild Carrot, Wild Four-o'clock, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Marigold, Wild Mustard, Wild Onion, Wild Parsnip, Wild Radish, Wild Rape, Wild Strawberry, Wild Sweet Potato, Wild Vetch, Wild Violet, Woodsorrel, Woolly Croton, Woolly Morningglory, Woolly Plantain, Wormseed, Yarrow, Yellow Foxtail, Yellowflower Pepperweed, Yellow Rocket, and other broadleaf weeds</p>

	<ul style="list-style-type: none"> • Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. • Do not allow people or pets to enter the treated area until sprays have dried.
---	--

<p><u>PRECAUTIONARY STATEMENTS</u></p>
<p>CAUTION Hazards to Humans and Domestic Animals</p>
<p>Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. Wear long sleeved shirt and long pants, shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.</p>

<p>First Aid +</p>	
<p>If Swallowed</p>	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
-------------------	--

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call toll-free 1-877-229-3763 for medical emergency information.**

Active Ingredients: 4.85% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.45% Dicamba, dimethylamine salt (CAS # 2300-66-5), and 1.61% Quinclorac (CAS # 84087-1-4).


Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Important – Directions for Storage and Disposal
Do not contaminate water, food, or feed by storage and disposal.

Storage Conditions	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
---------------------------	---

Pesticide Disposal and Container Handling 	Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.
--	--

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

D-9848 072205

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on the retail containers of supplemental registrants.

- One quart covers up to 4,000 square feet.
- 1 quart treats [up to] 4,000 sq.ft.
- Keep from freezing.
- Controls crabgrass, yellow nutsedge, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills even the tough weeds...roots and all!
- Controls even the tough weeds...roots and all!
- Kills the root.
- Water based.
- Won't harm lawn grass when used according to label directions.
- Kills lawn weeds plus crabgrass.
- Kills [Controls] 200+ broadleaf weeds.
- Kills the weeds not the lawn
- Attach to [garden] hose and spray
- Simple to use
- Easy to use
- Easy to mix
- Proven performance
- For home lawn care
- Won't harm lawn grass when used as directed.
- Controls (Number) weeds!
- Mixes with Water - Concentrate
- Kill crabgrass, yellow nutsedge & [any listed weed(s)] too!
- CONTROLS
 - SENSITIVE WEEDS: Crabgrass, yellow nutsedge, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.
 - HARD-TO-KILL WEEDS: Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and many other broadleaf weeds.
- Kills Lawn Weeds plus Crabgrass in One Step
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grass Weeds
- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed Weeds
- New!
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and certain weed grasses
- Kills Lawn Weeds plus Crabgrass in One Easy Step

- Won't Harm Lawns* (when used as directed)
- Kills 200+ Weeds (to the root)
- Rainproof in 1 Hour
- Treats 5,000 sq.ft.
- Starts Working Immediately
- All Weather Formula (rainproof in one hour).
- Bayer 100% Satisfaction Guaranteed* (or your money back)
- [optional advertising claims when product is sold in "low-water" hose-end sprayers on the order of 40 to 1 dilution ratio or as determined by testing]
 - Water [Saver] [Saving] Formula
 - Water [Saver][Saving] Applicator
 - Water [Saver] [Saving] Sprayer
 - Uses Up To [xx][50]% Less Water*
 - with the footnote disclaimer *vs. leading weed hose end sprayers

3. Trademark acknowledgements

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00894.20091016.notif-proposed-clean.doc

2. **Reason for Issue:** PRN 2007-4