

2217-896

06/26/2008

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

26 JUN 2008

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

James L. Kunstman, Ph.D.  
Manager of Regulatory Services  
PBI/Gordon Corporation  
1217 West 12<sup>th</sup> Street  
Post Office Box 014090  
Kansas City, Missouri 64101

Dear Dr. Kunstman:

Subject: EH-1437 Herbicide

Registration No. 2217-896

Application and Letter Dated November 5, 2007, Request To Amend Registration, in Response to EPA Reregistration Eligibility Decision for 2,4-D (EPA RED), 2,4-D Dimethylamine Salt, 8-month Response to the Reregistration Eligibility Decision, ID#: PDCI-030019-2610; Resubmission Application and Your Letter Dated April 3, 2008, Response to Agency's Letter Dated March 21, 2008

The resubmission application and your letter dated April 3, 2008; and revised labeling have been reviewed. The proposed labeling has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, provided that you:

1. Delete reference to "spot applications". On page 5 reference to "excluding spot treatments" in the table foot note and directions for spot treatments on the following pages of the label must be deleted (labeling claims for mixtures of 2,4-D and Dicamba may not claim "spot treatments", see enclosed "2,4-D and Dicamba Combination Label Table". Note, the limitation of 2 applications of Dicamba per year does not bear the phrase "excluding spot treatments" as allowed for 2,4-D products.
2. The issue of when 2,4-D products may be shipped and the matter of handling existing stocks are addressed in the 2,4-D RED (See pages 111 and 112). The standard statement that goes with Agency's accepted labeling: "Submit one copy of the final printed labeling prior to shipping under the revised labeling.", allows time to determine when to use the accepted labeling.

- 3. Submit one copy of the final printed labeling prior to shipping under the revised labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6 (e). Your release for shipment of this product under the revised labeling constitutes acceptance of these conditions, A stamped copy of the accepted labeling is enclosed for your records.

Please submit a copy of the labeling, submitted with your April 3, 2008 application, showing all revisions (designating deletions and additions) to the final printed label; and highlighted in yellow to allow an expedited review of the final printed labeling.

The enclosed copy of this Agency's "2,4-D and Dicamba Combination Label Table" should be followed in revising the labeling for your products that contain 2,4-D and Dicamba as active ingredients. As you have many such products, we request that you use the language and conditions as described in that Table at this time. Applications to amend the labeling of these products should include a statement that the labeling conforms to conditions described in that Table. Applications that do not bear such a statement may be unacceptable. .

Sincerely yours,



Joanne I. Miller  
 Product Manager (23)  
 Herbicide Branch  
 Registration Division (7505P)

Enclosures (2)

3/12

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:

26 JUN 2008  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

2217-896

# EH-1437 HERBICIDE

## EPA Reg. No. 2217-896

### ACTIVE INGREDIENT:

2,4-D, dimethylamine salt .....	6.42%
Quinclorac .....	2.13%
Dicamba, dimethylamine salt.....	0.60%
INERT INGREDIENTS:.....	90.85%
TOTAL.....	100.00%

### THIS PRODUCT CONTAINS:

- 0.456 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
  - 0.182 lb. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
  - 0.043 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
- Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

### User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.



<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **General Precautions and Restrictions:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

### **SHAKE WELL BEFORE USING**

#### **PRODUCT FACTS:**

**What It Does:** Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)

**Where to Use:** Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)

**When to Use:** Apply when daytime temperatures are below 90°F.

**Questions & Comments:** (icon phone) (icon computer)  
Call 1-800-xxx-xxxx or Visit our Website at [www.companyname.com](http://www.companyname.com)

### **HOW IT WORKS:**

EH-1437 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

5/12

**WHERE TO USE:**

EH-1437 Herbicide can be applied to residential lawns with the following turfgrass species:

<b>Lawns To Be Treated:</b>	
<b>Cool Season Turfgrass</b>	<b>Warm Season Turfgrass</b>
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	Bermudagrass <sup>1</sup> , common and hybrid Zoysiagrass Buffalograss
<sup>1</sup> Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.	

<b>DO NOT apply EH-1437 Herbicide to the following:</b>
Bahiagrass Bentgrass, colonial and creeping Carpetgrass Centipedegrass Dichondra Gardens including vegetables, fruit trees, vines and berries Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs) Seashore paspalum St. Augustinegrass Turfgrass species that are not listed on this label.

6/12

**GRASS AND BROADLEAF WEEDS CONTROLLED:**

A partial list is presented below:

<b>GRASS WEEDS</b>			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	
<b>BROADLEAF WEEDS</b>			
Annual yellow sweetclover	Cocklebur	Mustard	Spiny cocklebur
Aster	Common mullein	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Bedstraw	Cudweed	Nettle	Spotted spurge
Beggarticks	Curly dock	Orange hawkweed	Spurweed
Betony, Florida	Daisy, English	Oriental cocklebur	Stinging nettle
Bindweed, field	Daisy fleabane	Oxalis	Strawberry, India mock
Bird vetch	Daisy, oxeye	Parsley-piert	Tall nettle
Bitter wintercress	Dandelion	Parsnip	Tall vervain
Bittercress, hairy	Dichondra	Pearlwort	Tansy ragwort
Bitterweed	Dogbane	Pennycress	Tansy mustard
Black-eyed Susan	Dogfennel	Pennywort	Tanweed
Black medic	Dollarweed	Peppergrass	Thistle
Black mustard	Elderberry	Pepperweed	Trailing crownvetch
Blackseed plantain	False dandelion	Pigweed	Tumble mustard
Blessed thistle	Falseflax	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	Fiddleneck	Plantain	Venice mallow
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Water pennywort
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly lettuce	Western clematis
Broomweed	Gumweed	Prickly sida	Western salsify
Buckhorn	Hairy fleabane	Prostrate knotweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate pigweed	Wild aster
Bulbous buttercup	Healall	Prostrate spurge	Wild buckwheat
Bull thistle	Heartleaf drymary	Prostrate vervain	Wild carrot
Bullnettle	Heathaster	Puncturevine	Wild four-o'clock
Burclover	Hedge bindweed	Purslane, common	Wild garlic
Burdock	Hedge mustard	Ragweed	Wild geranium
Burning nettle	Hemp	Red sorrel	Wild lettuce
Burweed	Henbit	Redroot pigweed	Wild marigold
Buttercup	Hoary cress	Redstem filaree	Wild mustard
Buttonweed	Hoary plantain	Rough cinquefoil	Wild onion
Canada thistle	Hoary vervain	Rough fleabane	Wild parsnip
Carolina geranium	Horsenettle	Russian pigweed	Wild radish
Carpetweed	Jimsonweed	Russian thistle	Wild rape
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild strawberry
Catnip	Knotweed	Scotch thistle	Wild sweet potato
Catsear	Kochia	Sheep sorrel	Wild vetch
Chickweed, common	Lambsquarters	Shepherdspurse	Wild violet
Chickweed, mouseear	Lespedeza	Slender plantain	Woodsorrel
Chicory	Mallow	Smallflower galinsoga	Woolly croton
Cinquefoil	Matchweed	Smooth dock	Woolly morningglory
Clover, crimson	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, hop	Milk vetch	Sorrel	Wormseed
Clover, red	Morningglory	Sowthistle	Yarrow
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, sweet	Mugwort	Speedwell	Yellowflower pepperweed and other broadleaf weeds
Clover, white	Musk thistle	Spiny amaranth	
Cockle			

**WHEN TO USE:**

Apply this product when weeds are small and actively growing.

Weed species	Season	Timing	Maximum number of applications <sup>1</sup>	Application interval, days	Comments
<b>Crabgrass</b> (large and smooth):	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height	2	14	Re-treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence.
<b>Foxtail</b> (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations.
<b>Broadleaf signalgrass</b>	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	14	Re-treat at the same rate at 14 days later if needed.
<b>Dandelion, plantain, buttercup, speedwell, knotweed</b> and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
<b>Henbit, chickweed, white clover, ground ivy, lawn burweed</b> and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

<sup>1</sup>The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application (1.0 lb 2,4-D acid equivalent per acre per application).  
 The maximum number of broadcast applications is limited to 2 per year.  
 Do not apply more than 2 applications per year.  
 The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. (2.0 lbs 2,4-D acid equivalent per acre), excluding spot treatments.

**HOW MUCH TO USE:**

**I. ENTIRE LAWN TREATMENTS WITH DIAL TYPE HOSE-END SPRAYERS**

Three Easy Steps To Mix And Apply:

**1. FILL**

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- The application rate for this product is 6.4 fl.oz. of product per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 6.4 fl.oz. of product for every 1,000 sq.ft. of lawn area. Over-application or rates above those recommended on this label can cause turf injury.
- Do not add water to the sprayer jar.

**2. SET THE DIAL**

For hose-end sprayers: Set the dial to the rate per gallon and keep the dial set to this setting while spraying.

The dial setting or rate per gallon for this product is 2.5 oz. of product per gallon of water.

Types of hose-end sprayers	Dial setting
Ortho® Dial'N Spray® Hose-End Sprayer	2.5 oz.
Hudson® Hose End Liquid Concentrate Sprayer	2.5 fl. oz.
Gilmour® Dial-A-Mix® Lawn and Garden Sprayer	15 Teaspoons or 5 tablespoons

**3. SPRAY**— Apply evenly over the lawn.

**II. ENTIRE LAWN TREATMENTS WITH READY-TO-USE HOSE-END SPRAYERS :**

This product may be packaged as a ready-to-use product with no mixing. The retail plastic container is fitted with a sprayer body that delivers spray droplets at a fixed rate.

**For READY-TO-USE Hose End Sprayers:**

- This retail sprayer bottle will cover the treated areas described below when this product is applied at 6.4 fl.oz.of product per 1,000 sq.ft. Over-application or rates above those recommended on this label can cause turf injury.

Approximate treated areas for four retail container sizes:

Species	Amount of product to treat approximate area, sq. ft.				
	Fl.oz./1,000 sq.ft.	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container	2.0 qt. (64 fl.oz.) retail container	1 gallon (128 fl.oz.) retail container
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season species Bermudagrass <sup>1</sup> common and hybrid Zoysiagrass and Buffalograss	6.4 fl.oz. /1,000 sq.ft.	2,500 sq. ft.	5,000 sq.ft.	10,000 sq.ft.	20,000 sq.ft.

<sup>1</sup>Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

**Instructions:**

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.

Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl.oz. container will treat up to xxx sq.ft. of turfgrass.

- **FINISH:** To stop spraying, turn [rotate] spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn [rotate] spray nozzle to the "ON" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

**III. SPOT TREATMENTS WITH HAND-OPERATED SPRAYERS FOR SMALL AREAS OR INDIVIDUAL WEEDS:**

**For hand-operated sprayers (backpack, knapsack, tank-type or pump-up):**

- Apply to actively growing grass and broadleaf weeds during the early growth stages.
- Measure and mix the proper amount of product with water as described below. Mix 2.5 fl.oz. of product with 1 gallon of water to treat 400 sq.ft. of lawn. See below for additional spray preparations
- Spray evenly over the lawn. Spray coverage should be uniform and complete. Over-application or rates above those recommended on this label can cause turf injury.



9/12

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square feet
2.5 fl.oz.	1 gallon	400 sq.ft.
3.75 fl.oz.	1.5 gallons	600 sq.ft.
5.0 fl.oz.	2 gallons	800 sq.ft.
7.5 fl.oz.	3 gallons	1,200 sq.ft.

Equal measures: 2.5 fl.oz. = 5 Tablespoons. 1 cup = 8 fl.oz. = 1/2 pint.

OR

Measure indicated amount of product and pour into tank. Dilute with water according to the following chart. Adjust the nozzle to deliver coarse spray droplets and treat the area indicated.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Area To Treat, Square feet
2.5 fl.oz.	1 gallon	400 sq.ft.
5 fl. fl.oz.	2 gallons	800 sq.ft.
7.5 fl.oz.	3 gallons	1,200 sq.ft.

Equal measures: 2.5 fl.oz. = 5 Tablespoons. 1 cup = 8 fl.oz. = 1/2 pint.

**IV. ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® SYSTEM:**

**HOW TO USE THE READY-SPRAY® SYSTEM**

**Connect:**

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

**Spray:**

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small plastic tab back and turn knob clockwise to ON position.
3. Spray evenly over measured area.

**Finish:**

1. To STOP spraying, rotate the knob counterclockwise to the OFF position.
2. Turn off water.
4. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.
5. Disconnect sprayer from hose.

**V. FOR SPOT TREATMENT: TANK SPRAYER:**

- Add 2.5 fl.oz. per gallon of water for each 400 sq.ft.

For broadcast application: Ortho Dial 'N Spray® Applicator

1. Add 16 fl.oz. of concentrate directly into sprayer jar. DO NOT ADD WATER.
2. Set dial to 2.5 oz
3. Spray evenly over 2,500 sq.ft.
4. Any unused portion should be poured back into its original container.

**VI. FOR ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL [KILLER] SINGLES:**

**For Spot Treatment: (Use a) Tank Sprayer:**

- Add entire contents (of 1 bottle) ( 5 oz.) to an empty 2-gallon tank sprayer. Slowly fill with 2 gallons of clean tap water.

10/12

**(Instructions) For Spot Weed Treatment**

- Add entire contents (5 oz.) to an empty (2-gallon) tank sprayer. Slowly fill with 2 Gallons of clean tap water

<b>CULTURAL TIPS AND NOTES:</b>		
<b>When to Apply:</b>	<b>For established lawns For newly seeded areas For newly sodded, sprigged, or plugged areas</b>	<ul style="list-style-type: none"> <li>• Apply when daytime temperatures are below 90°F.</li> <li>• Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.</li> <li>• Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.</li> </ul>
<b>Seeding:</b>		<ul style="list-style-type: none"> <li>• The treated area can be seeded at 4 weeks after the application of this product.</li> </ul>
<b>Irrigation:</b>	<b>Non-stressed turfgrass Stress from high temperatures, drought, insects or diseases.</b>	<ul style="list-style-type: none"> <li>• Adequate soil moisture and favorable growing conditions enhance the performance of this product.</li> <li>• Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.</li> </ul>
<b>Mowing:</b>		<ul style="list-style-type: none"> <li>• Delay mowing 2 days before and until 2 days after the application of this product.</li> <li>• Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.</li> </ul>
<b>Nontarget plants:</b>		<ul style="list-style-type: none"> <li>• Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.</li> <li>• Do not spray exposed roots of ornamentals and trees.</li> </ul>

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Avoid contamination of food or feedstuffs.

**DISPOSAL: If Empty**—Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled**—Call your local solid waste agency or toll free number (1-800-XXX-XXXX) for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

[OPTIONAL WARRANTY STATEMENT]

The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

# **APPENDIX**

## **1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

## **2. Advertising claims that may be presented on the retail containers of supplemental registrants.**

- One quart covers up to 5,000 square feet.
- 1 quart treats [up to] 5,000 sq.ft.
- Keep from freezing.
- Controls crabgrass, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills even the tough weeds...roots and all!
- Controls even the tough weeds...roots and all!
- Kills the root.
- Water based.
- Won't harm lawn grass when used according to label directions.
- Kills lawn weeds plus crabgrass.
- Kills [Controls] 200+ broadleaf weeds.
- Kills the weeds not the lawn
- Attach to [garden] hose and spray
- Simple to use
- Easy to use
- Easy to mix
- Proven performance
- For home lawn care
- Won't harm lawn grass when used as directed.
- Controls (Number) weeds!
- Mixes with Water - Concentrate
- Kill crabgrass & [any listed weed(s)] too!

### **CONTROLS**

**SENSITIVE WEEDS:** Crabgrass, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.

**HARD-TO-KILL WEEDS:** Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and other broadleaf weeds.

- Kills Lawn Weeds plus Crabgrass in One Step
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grass Weeds
- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed Weeds
- New!
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Certain Weed Grasses

**For any of the ORTHO Products:**

- Makes more than XX Gallon!
- Kills Even Tough Weeds – Roots & All
- Lawn Weed & Crabgrass Killer
- Kills Dandelions, Clover, and Other Broadleaf Weeds! Plus Kills Crabgrass and Other grass Weeds!
- Kills the Toughest Weeds in these Lawns (any lawn listed on label)
- Open Resealable Label for Directions & Precautions (Press to Reseal)
- People or pets may reenter treatment area after spray has dried.
- Ready-Spray
- Do Not Use on St. Augustine Lawns
  - Ortho Weed-B-Gon MAX Concentrate Plus Crabgrass Control [Killer]
  - Kills Weeds, Not Lawns(TM)
  - For [Use on] Lawns
  - ORTHO & Your Environment - Your yard is a place for your family and pets to enjoy. That's why ORTHO products are designed with care to provide effective solutions to weed problems. For best results and a healthy environment, please follow instructions for appropriate usage, storage and disposal.
- Kills Over 200 Lawn Weeds plus [including] Crabgrass [in one easy step.
- Guaranteed [Results]
- See Ortho.com for complete Spanish Translation.
- DO NOT use on St. Augustine, including Floratam species.

**For ORTHO Weed-B-Gon MAX Plus Crabgrass Control [Killer] Singles:**

- NET 5 FL.OZ.
- Pre-Measured Concentrate Bottles
- Treats 5,000 sq.ft.
- No Measuring, No Mess
- Weed-B-Gon® Max(TM) Singles makes using concentrates easier than ever. No more measuring and no more mess. Mix (contents) (each pre-measured bottle) with two gallons of water to deliver consistent results. And its' revolutionary formula won't harm your lawn - Guaranteed.

**3. Trademark acknowledgements**

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

**DOCUMENT CONTROL INFORMATION**

**1. Unique Label Identifier:** 002217-00896.20080403.proposed.doc

**2. Reason for Issue:** Modify per EPA comments. Environmental Hazards, Warranty, Ad Claims.