



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D C 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

James L Kunstman Ph D
PBI/Gordon Corporation
Post Office Box 014090
Kansas City, Missouri 64101

OCT - 4 2012

Subject Amendment with minor revisions to Directions for Use applied through Sub-labeling
Product name EH-1437 Herbicide
EPA Reg No 2217-896
Application Dated July 5 2012

Dear Dr Kunstman

The labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide, and Rodenticide Act (FIFRA) as amended is acceptable

Please note that marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available then those claims must be removed.

40 CFR 156 10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of the label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Grant Rowland at 703-347-0254 or at Rowland.Grant@epa.gov


Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

EH-1437 HERBICIDE

EPA Reg. No. 2217-896



Two sublabels represent the entire or master label of EH-1437 HERBICIDE. Please refer to the appropriate section of the labeling as shown as follows:

Section
Label language common to all sublabels
Sublabel 1 Traditional layout
Sublabel 2 Concentrate
Sublabel 3 Ready Spray
Appendix
Document Control Information

Label Language Common To All Sublabels

EH-1437 HERBICIDE

EPA Reg. No. 2217-896

Alternate Brand Names

**Tractor Supply Co GroundWork Concentrate Crabgrass & Broadleaf Weed Killer
Gordon s Trimec Crabgrass Plus Lawn Weed Killer Ready Spray
Gordon s Trimec Crabgrass Plus Lawn Weed Killer Concentrate**

ACTIVE INGREDIENT

2 4-D dimethylamine salt 6 42%
Quinclorac 2 13%
Dicamba dimethylamine salt 0 60%

INERT INGREDIENTS

TOTAL 90 85%
100 00%

THIS PRODUCT CONTAINS

0 456 lb 2 4-dichlorophenoxyacetic acid equivalent per gallon or 5 33%
0 182 lb 3 7-dichloro-8-quinolinecarboxylic acid per gallon or 2 13%
0 043 lb 3 6-dichloro-o-anisic acid equivalent per gallon or 0 50%
Isomer Specific By AOAC Methods

Note If the design format or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel then add a referral statement and present this substatement on the back panel **“See back panel for substatement of ingredient statement”**

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION Causes moderate eye irritation Avoid contact with eyes skin or clothing Harmful if swallowed When using this product wear long-sleeved shirt long pants socks shoes and rubber gloves

Wash thoroughly with soap and water after handling and before eating drinking chewing gum or using tobacco After using this product rinse gloves before removing remove clothing and launder separately before reuse and promptly and thoroughly wash hands and exposed skin with soap and water Remove saturated clothing as soon as possible and shower

User Safety Recommendations

- Users should wash hands before eating drinking chewing gum using tobacco or using the toilet
- Users should remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing If pesticide gets on skin wash immediately with soap and water
- Users should remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

First Aid	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses if present after the first minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to by a poison control center or doctor • Do not give anything by mouth to an unconscious person
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1-877-800-5556 for emergency medical treatment advice	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas Do not contaminate water when disposing of equipment wash waters or rinsate

This chemical has properties and characteristics associated with chemicals detected in groundwater The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination Application around a cistern or well may result in contamination of drinking water or groundwater

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates To protect the environment do not allow pesticide to enter or run off into storm drains drainage ditches gutters or surface waters Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems

This chemical has properties and characteristics associated with chemicals detected in groundwater The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination Application around a cistern or well may result in contamination of drinking water or groundwater]

DIRECTIONS FOR USE

STORAGE AND DISPOSAL
 Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Keep from freezing Store in original container in a locked storage area inaccessible to children and pets Avoid contamination of food or feedstuffs

PESTICIDE DISPOSAL AND CONTAINER HANDLING Nonrefillable container Do not reuse or refill

this container **If empty** Place in trash or offer for recycling if available **If partly filled** Call your local solid waste agency or www website com for disposal instructions Never place unused product down any indoor or outdoor drain

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE HEREBY EXPRESSLY DISCLAIMED IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT If these terms are not acceptable return this product unopened immediately to the point of purchase and the purchase price will be refunded in full The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere

Alternate Warranty

NOTICE To the extent consistent with applicable law buyer assumes all responsibility for lack of performance or safety if not used according to the label directions

[OPTIONAL WARRANTY STATEMENT]

The ORTHO Guarantee

If for any reason you are not satisfied with this product mail us proof of purchase to obtain a full refund of your purchase price


6/24

SUBLABEL 1 Traditional layout

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

GENERAL PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet either directly or through drift
Keep people and pets out of the area during application
-  Optional Icon] Do not allow people or pets to enter the treated area until sprays have dried

SHAKE WELL BEFORE USING

PRODUCT FACTS

What It Does Kills major broadleaf and troublesome grass weeds including dandelion chickweed and clover as well as crabgrass and foxtail (See inside for complete list)

Where to Use Lawns Kills the weeds not the lawn when used as directed (icon weeds)

When to Use Apply when daytime temperatures are below 90°F

Questions & Comments (icon phone) (icon computer)
Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com

HOW IT WORKS

EH 1437 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE

EH-1437 Herbicide can be applied to residential lawns with the following turfgrass species

Lawns To Be Treated	
Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass	Bermudagrass ¹ common and hybrid
Ryegrass perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues (red hard and chewings)	

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F

DO NOT apply EH 1437 Herbicide to the following

- Bahiagrass
- Bentgrass colonial and creeping
- Carpetgrass
- Centipedegrass
- Dichondra
- Gardens including vegetables fruit trees vines and berries
- Ornamental plants (flowers trees groundcovers landscape beds and shrubs)
- Seashore paspalum

St Augustinegrass
 Turfgrass species that are not listed on this label

GRASS AND BROADLEAF WEEDS CONTROLLED

A partial list is presented below

GRASS WEEDS			
Crabgrass large and smooth	Foxtail green yellow and giant	Signalgrass broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Mustard	Spiny cocklebur
Aster	Common mullein	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Bedstraw	Cudweed	Nettle	Spotted spurge
Beggarticks	Curly dock	Orange hawkweed	Spurweed
Betony Florida	Daisy English	Oriental cocklebur	Stinging nettle
Bindweed field	Daisy fleabane	Oxalis	Strawberry India mock
Bird vetch	Daisy oxeye	Parsley piert	Tail nettle
Bitter wintercress	Dandelion	Parsnip	Tall vervain
Bittercress hairy	Dichondra	Pearlwort	Tansy ragwort
Bitterweed	Dogbane	Pennycress	Tansy mustard
Black eyed Susan	Dogfennel	Pennywort	Tanweed
Black medic	Dollarweed	Peppergrass	Thistle
Black mustard	Elderberry	Pepperweed	Trailing crownvetch
Blackseed plantain	False dandelion	Pigweed	Tumble mustard
Blessed thistle	False flax	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	Fiddleneck	Plantain	Venice mallow
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Water pennywort
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prairie sunflower	Western clematis
Broomweed	Gumweed	Prickly lettuce	Western salsify
Buckhorn	Hairy fleabane	Prickly sida	White mustard
Buckhorn plantain	Hawkweed	Prostrate knotweed	Wild aster
Bulbous buttercup	Healall	Prostrate pigweed	Wild buckwheat
Bull thistle	Heartleaf drymary	Prostrate spurge	Wild carrot
Bullnettle	Heathaster	Prostrate vervain	Wild four-o clock
Burclover	Hedge bindweed	Puncturevine	Wild garlic
Burdock	Hedge mustard	Purslane common	Wild geranium
Burning nettle	Hemp	Ragweed	Wild lettuce
Burweed	Henbit	Red sorrel	Wild mangold
Buttercup	Hoary cress	Redroot pigweed	Wild mustard
Buttonweed	Hoary plantain	Redstem filaree	Wild onion
Canada thistle	Hoary vervain	Rough cinquefoil	Wild parsnip
Carolina geranium	Horsenettle	Rough fleabane	Wild radish
Carpetweed	Jimsonweed	Russian pigweed	Wild rape
Catchweed bedstraw	Knawel	Russian thistle	Wild strawberry
Catnip	Knotweed	Scarlet pimpernel	Wild sweet potato
Catsear	Kochia	Scotch thistle	Wild vetch
Chickweed common	Lambsquarters	Sheep sorrel	Wild violet
Chickweed mouseear	Lespedeza	Shepherdspurse	Woodsor cl
Chicory	Mallow	Slender plantain	Woolly croctc
Cinquefoil	Matchweed	Smallflower galinsoga	Woolly morningglory
Clover crimson	Mexicanweed	Smooth dock	Woolly plantain
Clover hop	Milk vetch	Smooth pigweed	Wormseed
Clover red	Morningglory	Sorrel	Yarrow
Clover strawberry		Sowthistle	Yellow rocket

BROADLEAF WEEDS			
Clover sweet	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover white	Mugwort	Speedwell	and other broadleaf
Cockle	Musk thistle	Spiny amaranth	weeds

WHEN TO USE

Apply this product when weeds are small and actively growing

Weed species	Season	Timing	Maximum number of applications ¹	Application interval days	Comments
Crabgrass (large and smooth)	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height	2	14	Re treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence
Foxtail (green giant and yellow)	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height	2	14	Re treat at the same rate at 14 days later for dense populations
Broadleaf signalgrass	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage	2	14	Re treat at the same rate at 14 days later if needed
Dandelion plantain buttercup speedwell knotweed and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture
Henbit chickweed white clover ground ivy lawn burweed and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

¹The maximum application rate is 6.4 fl oz of product per 1 000 sq ft per application [1.0 lb 2.4 D acid equivalent per acre per application]. The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 applications per year. The maximum seasonal rate is 12.8 fl oz of product per 1 000 sq ft [2.0 lb 2.4 D acid equivalent per acre].

HOW MUCH TO USE

I ENTIRE LAWN TREATMENTS WITH DIAL TYPE HOSE-END SPRAYERS

Three Easy Steps To Mix And Apply

1 FILL

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas multiply the length by the width to obtain total square feet.
- The application rate for this product is 6.4 fl oz of product per 1 000 sq ft of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area or add 6.4 fl oz of product for every 1 000 sq ft of lawn area. Over application or rates above those recommended on this label can cause turf injury.
- Do not add water to the sprayer jar.

2 SET THE DIAL

For hose-end sprayers Set the dial to the rate per gallon and keep the dial set to this setting while spraying.

The dial setting or rate per gallon for this product is 2.5 oz of product per gallon of water.

Types of hose-end sprayers	Dial setting
Ortho® Dial N Spray® Hose End Sprayer	2.5 oz
Hudson® Hose End Liquid Concentrate Sprayer	2.5 fl oz
Gilmour® Dial A Mix® Lawn and Garden Sprayer	15 Teaspoons or 5 tablespoons

3 **SPRAY** - Apply evenly over the lawn

II ENTIRE LAWN TREATMENTS WITH READY-TO-USE HOSE-END SPRAYERS

This product may be packaged as a ready-to-use product with no mixing. The retail plastic container is fitted with a sprayer body that delivers spray droplets at a fixed rate.

For READY-TO-USE Hose End Sprayers

- This retail sprayer bottle will cover the treated areas described below when this product is applied at 6.4 fl oz of product per 1,000 sq ft. Over-application or rates above those recommended on this label can cause turf injury.

Approximate treated areas for four retail container sizes

Species	Amount of product to treat approximate area sq ft				
	Fl oz /1 000 sq ft	1.0 pt (16 fl oz) sprayer jar or retail container	1.0 qt (32 fl oz) sprayer jar or retail container	2.0 qt (64 fl oz) retail container	1 gallon (128 fl oz) retail container
Kentucky bluegrass Ryegrass perennial and annual Tall fescue Mixtures of cool season species Bermudagrass ¹ common and hybrid Zoysiagrass and Buffalograss	6.4 fl oz /1 000 sq ft	2 500 sq ft	5 000 sq ft	10 000 sq ft	20 000 sq ft

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Instructions

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- CONNECT** Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the OFF position.
- SPRAY** Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the SPRAY position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.

Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl oz container will treat up to xxx sq ft of turfgrass.

- FINISH** To stop spraying, turn [rotate] spray nozzle lever to the OFF position. Turn off water at faucet. To relieve pressure, turn [rotate] spray nozzle to the ON position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

III HAND-OPERATED SPRAYERS

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up)

- Apply to actively growing grass and broadleaf weeds during the early growth stages
- Measure and mix the proper amount of product with water as described below Mix 2.5 fl oz of product with 1 gallon of water to treat 400 sq ft of lawn See below for additional spray preparations
- Spray evenly over the lawn Spray coverage should be uniform and complete Over-application or rates above those recommended on this label can cause turf injury

Amount of Product to Use Fluid Ounces	Amount of Water in Sprayer Gallons	Approximate Treated Area Square feet
2.5 fl oz	1 gallon	400 sq ft
3.75 fl oz	1.5 gallons	600 sq ft
5.0 fl oz	2 gallons	800 sq ft
7.5 fl oz	3 gallons	1,200 sq ft

Equal measures 2.5 fl oz = 5 Tablespoons 1 cup = 8 fl oz = 1/2 pint

OR

Measure indicated amount of product and pour into tank Dilute with water according to the following chart Adjust the nozzle to deliver coarse spray droplets and treat the area indicated

Amount of Product to Use Fluid Ounces	Amount of Water in Sprayer Gallons	Area To Treat, Square feet
2.5 fl oz	1 gallon	400 sq ft
5 fl oz	2 gallons	800 sq ft
7.5 fl oz	3 gallons	1,200 sq ft

Equal measures 2.5 fl oz = 5 Tablespoons 1 cup = 8 fl oz = 1/2 pint

IV ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® SYSTEM

HOW TO USE THE READY-SPRAY® SYSTEM

Connect

- 1 Shake well before using
- 2 Connect sprayer to hose
- 3 Turn on water
- 4 Extend hose to farthest area of the lawn to be treated

Spray

- 1 To begin spraying point nozzle in the direction you want to spray
- 2 Bend small plastic tab back and turn knob clockwise to ON position
- 3 Spray evenly over measured area

Finish

- 1 To STOP spraying rotate the knob counterclockwise to the OFF position
- 2 Turn off water
- 3 Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip Then turn knob back to OFF position
- 4 Disconnect sprayer from hose

V FOR TANK SPRAYER

- Add 2.5 fl oz per gallon of water for each 400 sq ft

For broadcast application Ortho Dial N Spray® Applicator

- 1 Add 16 fl oz of concentrate directly into sprayer jar DO NOT ADD WATER
- 2 Set dial to 2.5 oz
- 3 Spray evenly over 2,500 sq ft
- 4 Any unused portion should be poured back into its original container

[Over-application or rates above those recommended on this label can cause turf injury]

VI FOR ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL [KILLER] SINGLES

For (Use a) Tank Sprayer

- Add entire contents (of 1 bottle) (5 oz) to an empty 2-gallon tank sprayer Slowly fill with 2 gallons of clean tap water
- **For Broadcast Application – Ortho® Dial N Spray® Applicator**
 - 1 Set dial to 2.5 oz
 - 2 Add entire contents of 1 bottle (5 oz) directly into sprayer jar DO NOT ADD WATER
 - 3 Spray evenly over 800 sq ft

[Over application or rates above those recommended on this label can cause turf injury]

VII ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® II [2]/APPLICATOR NAME SYSTEM

HOW TO USE THE READY-SPRAY® II/[READY-SPRAY 2]/[APPLICATOR NAME] SYSTEM

[Optional]

- 1 [Measure the total [entire] area to be sprayed For rectangular areas multiply the length by width to obtain total square feet
- 2 See below directions for the ready-to-use hose-end sprayer
- 3 Spray evenly over the entire measured area]

[End of optional section]

Connect

- 1 Shake well before using [optional step]
- 2 Connect sprayer to hose
- 3 Turn on water
- 4 Extend hose to farthest area of the measured area to be treated [optional step]

Spray

- 1 To begin spraying [disengage safety lock and] point nozzle in the direction you want to spray
- 2 Slide thumb activation switch forward to ON position
- 3 Spray evenly over measured area

Finish

- 1 To STOP spraying slide thumb activation switch backward to OFF position
- 2 Turn off water
- 3 Relieve water pressure by sliding thumb switch to ON /WATER/ WATER ONLY position until water slows to a drip Then slide switch back to OFF position [and engage safety lock]
- 4 Disconnect sprayer from hose

[Optional illustrations may be used in conjunction with the above steps]

CULTURAL TIPS AND NOTES		
When to Apply	For established lawns For newly seeded areas For newly sodded sprigged or plugged areas	<ol style="list-style-type: none"> 1 Apply when daytime temperatures are below 90 F 2 Apply this product to newly seeded grasses when well established (approximately 6 weeks after seedling emergence) or after the third mowing 3 Apply this product 3 to 4 weeks after the sodding sprigging or plugging operations
Seeding		<ul style="list-style-type: none"> • The treated area can be seeded at 4 weeks after the application of this product

Irrigation	Non stressed turfgrass Stress from high temperatures drought insects or diseases	<ul style="list-style-type: none"> • Adequate soil moisture and favorable growing conditions enhance the performance of this product • Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing		<ul style="list-style-type: none"> • Delay mowing 2 days before and until 2 days after the application of this product. • Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
Nontarget plants		<ul style="list-style-type: none"> • Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. • Do not spray exposed roots of ornamentals and trees.

22 4
69 2

SUBLABEL 2 Concentrate

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

1 Before You Use Read & follow these directions when using

- Do not apply this product in a way that will contact any person or pet either directly or through drift Keep people and pets out of the area during application Do not allow people or pets to enter the treated area until sprays have dried
- Use on established lawns Will not harm lawn grasses when used as directed Do not use on Bahiagrass Bentgrass (colonial and creeping) Carpetgrass Centipedegrass Dichondra Gardens (including vegetables fruit trees vines and berries) Ornamental plants (flowers trees groundcovers landscape beds and shrubs) Seashore Paspalum St Augustinegrass and turfgrass species that are not listed on this label)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected Do not apply to bermudagrass when daytime temperatures exceed 85° F

2 How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings) Ryegrass (perennial and annual) Tall Fescue, and Zoysiagrass

Spot Treatments Mix 2.5 fl oz (5 Tbs) in 1 gallon of water to treat 400 sq ft

Entire Lawn Treatments

- Pump Style Sprayer Mix 6.4 fl oz in 0.5 to 1 gallon of water to treat 1,000 sq ft
- Hose-End Sprayer Apply 6.4 fl oz per 1,000 sq ft
- [12 volt &] Tow-Behind Sprayers Mix 6.4 fl oz in 0.5 to 1 gallon of water to treat 1,000 sq ft

Over-application or rates above those specified on this label can cause turf injury

3 When To Use

Apply this product when weeds are small and actively growing

Weed species	Timing	Comments
Crabgrass (large and smooth)	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re treat at the same rate at 14 days later for dense populations or fo continued crabgrass emergence
Foxtail (green giant and yellow)	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height	Re treat at the same rate at 14 days later for dense populations
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage	Re treat at the same rate at 14 days later if needed
Dandelion plantain buttercup speedwell knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture

Weed species	Timing	Comments
Henbit chickweed white clover ground ivy lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture
The maximum application rate is 6.4 fl oz of product per 1,000 sq ft per application [1.0 lb 2.4 D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 14 days between applications. The maximum seasonal rate is 12.8 fl oz of product per 1,000 sq ft [2.0 lb 2.4 D acid equivalent per acre].		

4 How To Apply

4.1 Pump Style Sprayers [sprayer icon]

Ideal for spraying individual weeds

- 1 Add the appropriate amount of water and concentrate to the sprayer tank
- 2 Close sprayer, shake well, and pump handle to pressurize
- 3 Adjust nozzle to deliver a coarse spray pattern
- 4 To broadcast over large areas, apply evenly to treated area
- 5 To treat individual weeds, spray to wet weed leaf surfaces
- 6 Re-pressurize the sprayer as needed to maintain a good spray pattern

4.2 [12 Volt &] Tow-Behind Boom Sprayers [sprayer icon]

Ideal for treating large country and suburban lawns

- 1 To calibrate your application equipment, consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq ft. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq ft at 40 psi.
- 2 To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
- 3 Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Tank Size	Spray volume	Coverage Area	Concentrate to add to tank
15 gal	0.5 gal/1000 sq ft	30,000 sq ft	192 fl oz (1.5 gal)
	1 gal/1000 sq ft	15,000 sq ft	96 fl oz (0.75 gal)
25 gal	0.5 gal/1000 sq ft	50,000 sq ft	320 fl oz (2.5 gal)
	1 gal/1000 sq ft	25,000 sq ft	160 fl oz (1.25 gal)

For spot treatments with [12 Volt &] Pull Behind Sprayers

See rates under How Much To Apply Spot Treatments

4.3 Dial Style Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

- 1 Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER**
- 2 Close sprayer and set dial to 2.5 oz
- 3 Connect sprayer to hose
- 4 Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
- 5 Pour any unused product back into this container.

4.4 Concentrate Plus Water Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar.
3. Close sprayer, connect to garden hose, shake well.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

5. For Best Results

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

6. Grass and Broadleaf Weeds Killed

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Mustard	Spiny cocklebur
Aster	Common mullein	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Bedstraw	Cudweed	Nettle	Spotted spurge
Beggarticks	Curly dock	Orange hawkweed	Spurweed
Betony, Florida	Daisy, English	Oriental cocklebur	Stinging nettle
Bindweed, field	Daisy fleabane	Oxalis	Strawberry, India mock
Bird vetch	Daisy, oxeye	Parsley-piert	Tall nettle
Bitter wintercress	Dandelion	Parsnip	Tall vervain
Bittercress, hairy	Dichondra	Pearlwort	Tansy ragwort
Bitterweed	Dogbane	Pennycress	Tansy mustard
Black-eyed Susan	Dogfennel	Pennywort	Tanweed
Black medic	Dollarweed	Peppergrass	Thistle
Black mustard	Elderberry	Pepperweed	Trailing crownvetch
Blackseed plantain	False dandelion	Pigweed	Tumble mustard
Blessed thistle	Falseflax	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	Fiddleneck	Plantain	Venice mallow
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Water pennywort
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain			Western clematis

BROADLEAF WEEDS			
Broomweed	Ground ivy	Prickly lettuce	Western salsify
Buckhorn	Gumweed	Prickly sida	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild buckwheat
Bull thistle	Healall	Prostrate spurge	Wild carrot
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild four o'clock
Burclover	Heathaster	Puncturevine	Wild garlic
Burdock	Hedge bindweed	Purslane common	Wild geranium
Burning nettle	Hedge mustard	Ragweed	Wild lettuce
Burweed	Hemp	Red sorrel	Wild margold
Buttercup	Henbit	Redroot pigweed	Wild mustard
Buttonweed	Hoary cress	Redstem filaree	Wild onion
Canada thistle	Hoary plantain	Rough cinquefoil	Wild parsnip
Carolina geranium	Hoary vervain	Rough fleabane	Wild radish
Carpetweed	Horsenettle	Russian pigweed	Wild rape
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild strawberry
Catnip	Knawel	Scarlet pimpernel	Wild sweet potato
Catsear	Knotweed	Scotch thistle	Wild vetch
Chickweed common	Kochia	Sheep sorrel	Wild violet
Chickweed mouseear	Lambsquarters	Shepherdspurse	Woodsorrel
Chicory	Lespedeza	Slender plantain	Woolly croton
Cinquefoil	Mallow	Smallflower galinsoga	Woolly morningglory
Clover crimson	Matchweed	Smooth dock	Woolly plantain
Clover hop	Mexicanweed	Smooth pigweed	Wormseed
Clover red	Milk vetch	Sorrel	Yarrow
Clover strawberry	Momingglory	Sowthistle	Yellow rocket
Clover sweet	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover white	Mugwort	Speedwell	and other broadleaf
Cockle	Musk thistle	Spiny amaranth	weeds

17/24

[Back Panel] [Restatement of information in sections 1 and 2]

Getting Started

Kills

Crabgrass foxtail and most troublesome broadleaf weeds including dandelion clover thistle and over
XXX other weeds listed

Where To Use

On established lawns without harming lawn grasses when used as directed (Do not use on Bahiagrass
Bentgrass (colonial and creeping) Carpetgrass Centipede grass Dichondra Seashore Paspalum St
Augustinegrass and turfgrass species that are not listed on this label) [See Inside Label for Complete
Directions]

Amount To Use

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass

Spot Treatments Mix 2.5 fl oz (5 Tbs) in 1 gallon of water to treat 400 sq ft

Entire Lawn Treatments

- Pump Style Sprayer Mix 6.4 fl oz in 0.5 to 1 gallon of water to treat 1,000 sq ft
- Hose-End Sprayer Apply 6.4 fl oz per 1,000 sq ft [See Inside Label for Sprayer Settings]
- [12 volt &] Tow-Behind Sprayers Mix 6.4 fl oz in 0.5 to 1 gallon of water to treat 1,000 sq ft

Coverage

This [size] bottle treats XXXX sq ft

Telephone Icon

Computer Icon

For questions or comments call toll-free 1-800-821 7925 or visit our website at www.pbigordon.com

SUBLABEL 3 Ready Spray

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

1 Before You Use Read & follow these directions when using

- Do not apply this product in a way that will contact any person or pet either directly or through drift Keep people and pets out of the area during application Do not allow people or pets to enter the treated area until sprays have dried
- Use on established lawns Will not harm lawn grasses when used as directed Do not use on Bahiagrass Bentgrass (colonial and creeping) Carpetgrass Centipedegrass Dichondra Gardens (including vegetables fruit trees vines and berries) Ornamental plants (flowers trees groundcovers landscape beds and shrubs) Seashore Paspalum St Augustinegrass and turfgrass species that are not listed on this label)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected Do not apply to bermudagrass when daytime temperatures exceed 85° F

2 How Much To Apply

For Bermudagrass (common and hybrid) Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red hard and chewings), Ryegrass (perennial and annual) Tall Fescue and Zoysiagrass

Entire Lawn Treatments This container covers XXX sq ft Measure the size of your yard If the area to be treated is larger than XXX sq ft you may need to buy more containers

Over-application or rates above those specified on this label can cause turf injury

3 When To Use

Apply this product when weeds are small and actively growing

Weed species	Timing	Comments
Crabgrass (large and smooth)	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence
Foxtail (green giant and yellow)	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height	Re treat at the same rate at 14 days later for dense populations
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage	Re treat at the same rate at 14 days later if needed
Dandelion plantain buttercup speedwell knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture
Henbit chickweed white clover ground ivy lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture

The maximum application rate is 6.4 fl oz of product per 1 000 sq ft per application [1.0 lb 2.4 D acid equivalent per acre per application] The maximum number of applications is limited to 2 per year with a minimum of 14 days between applications The maximum seasonal rate is 12.8 fl oz of product per 1 000 sq ft [2.0 lb 2.4 D acid equivalent per acre]

4 How To Apply

- 1 Connect your garden hose to the hose-end sprayer With your thumb snap off the safety locking tab that prevents accidental discharge of spray solution Leave control valve in the OFF position
- 2 To BEGIN spraying Turn on water at faucet Extend hose to the furthest point from the faucet Point nozzle toward the lawn to be treated and turn the control valve to the ON position Water will automatically mix with the product Walk backwards as you apply in order to avoid contact with the treated area For best results spray evenly over the measured area
- 3 To STOP spraying Turn control valve to the OFF position Turn off water at faucet To relieve pressure turn control valve to ON position and wait until water stops spraying before removing sprayer from hose

5 For Best Results

- Apply when weeds are young and actively growing
- Delay mowing 2 days before and until 2 days after the application of this product
- If the lawn is dry water 24 hours before using this product
- Do not water lawn within 24 hours of treatment
- Certain hard to-kill weeds may need to be spot treated in 14 days
- For new lawns apply only when grass has reached a height of at least 2 inches
- Bare spots may be seeded 3 to 4 weeks after application
- Spray when air temperatures are cooler than 90°F
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures
- Do not use clippings from the treated areas as mulch or compost around flowers ornamentals trees or in vegetable gardens

6 Grass and Broadleaf Weeds Killed

A partial list is presented below

GRASS WEEDS			
Crabgrass large and smooth	Foxtail green yellow and giant	Signalgrass broadleaf	

BROADLEAF WEEDS			
Amaranth purple	Cupid shaving Brush	Mustard ball	Speedwell common
Amaranth slender	Cudweed	Mustard black	Speedwell corn
Aster heath	Daisy English	Mustard blue	Speedwell ivyleaf
Aster white heath	Daisy oxeye	Mustard hedge	Speedwell persian
Aster purple	Dandelion common	Mustard Indian	Speedwell purslane
Aster white prairie	Dandelion false	Mustard tansy	Speedwell slender
Artichoke	Day flower	Mustard tumble	Speedwell snow
Austrian fieldcress	Deadnettle	Mustard white	Speedwell thymeleaf
Bedstraw Catchweed	Dichondra	Mustard wormseed	Spiny amaranth
Bedstraw smooth	Dock broadleaf	Nettle stinging (burning)	Spurge spotted
Bedstraw Northern	Dock curly	Nettle tall	Spurge prostrate
Bedstraw pineywoods	Dogbane	Nettle wood	Spurweed
Beggarticks hairy	Dogfennel	Parsley piert	Star of Bethlehem
Betony Florida	Dollarweed (Pennywort)	Parsnip	Strawberry leaf mock
Bindweed Field	Dove weed	Pearlwort	Tansy ragwort
Bindweed Hedge	Elderberry	Pennycress (frenchweed)	Tanweed
Bitter Wintercress	Falseflax	Pepperweed perennial	Thistle blessed
Bittercress hairy	False sunflower	Pepperweed virginia	Thistle bull
Bitterweed	Fiddleneck	(Peppergrass)	Thistle canada
Black eyed Susan	Filaree redstem	Pigweed prostate	Thistle floodman (prairie)

BROADLEAF WEEDS			
Black Medic	Filaree whitestem	Pigweed redroot	Thistle musk
Blood flower milkweed	Fleabane daisy (annual)	Pigweed russian	Thistle platte
Blue lettuce	Fleabane rough	Pigweed smooth	Thistle plumeless
Brassbuttons	Galinsoga hairy	Pigweed tumble	Thistle russian
Bristly oxtongue	Galinsoga smallflower	Pineapple weed	Thistle scotch
Broomweed	Goldenrod	Plains coreopsis (tickseed)	Thistle wavyleaf
Buckhorn	Ground ivy	Plantain blackseed	Thistle yellow (horrible)
Bullnettle	Groundsel	Plantain bracted	Trailing crownvetch
Burclover	Gumweed	Plantain broadleaf	Velvetleaf
Burdock	Hairy fleabane	Plantain buckhorn	Vervain blue
Burweed lawn (spurweed)	Hawkweed meadow	Plantain hoary	Vervain hoary
Buttercup bulbous	Hawkweed mouseear	Plantain narrowleaf	Vervain prostrate
Buttercup creeping	Hawkweed orange	Plantain slender	Vervain tall
Buttonweed common (poorjoe)	Hawkweed yellow	Plantain woolly	Vetch bird
Buttonweed virginia	Healall	Poison ivy	Vetch hairy
Carolina geranium (wild)	Heartleaf drymary	Poison oak	Vetch milk
Carpetweed	Hemp	Pokeweed	Vetch narrowleaf
Catnip	Henbit	Prairie Sunflower	Vetch wild
Catsear spotted (false dandelion)	Hoary cress	Prickly lettuce (compass plant)	Virginia creeper
Chamber bitter	Horsenettle	Prickly sida	Western clematis
Chickweed common	Horseweed	Prostrate knotweed	Western salsify
Chickweed mouseear	Jimsonweed	Puncturevine	Wild aster
Chickweed sticky	Johnny Jumpup violet	Purslane common	Wild buckweat
Chicory	Knawel	Pusley brazil	Wild carrot
Cinquefoil	Knotweed prostrate	Pusley florida	Wild four o clock
Clover crimson	Kochia	Pusley large flower	Wild garlic
Clover alyce	Lambsquarters	Ragweed bur	Wild lettuce
Clover hop	Lespedeza	Ragweed common	Wild mangold
Clover rabbit foot	Mallow alkali	Ragweed lanleaf	Wild mustard (yellow rocket)
Clover red	Mallow bristly	Ragweed western	Wild onion
Clover strawberry	Mallow common	Redstem filaree	Wild parsnip
Clover yellow sweet	Mallow dwarf	Rough cinquefoil	Wild radish
Clover white sweet	Mallow venice	Scarlet pimpernel	Wild rape
Clover white	Marcela	Shepardspurse	Wild strawberry
Cockle	Matchweed	Smartweed ladysthumb	Wild sweet potato
Cocklebur	Mexicanweed	Smartweed pale	Wild violet
Cocklebur oriental	Moneywort	Smartweed pennsylvania	Woodsorrel creeping
Cocklebur spiny	Morningglory bigroot	Smooth chaff flower	Woodsorrel violet
Common mullein	Morningglory ivyleaf	Smoothdock	Woodsorrel yellow (Oxalis)
Corn Spurry	Morningglory mexican	Sorrel red (sheep)	Woolly croton
Creeping Beggarweed	Morningglory red	Sowthistle annual	Wormseed
Creeping Jenny (creeping charlie)	Morningglory tall	Sowthistle spiny	Yarrow common
	Morningglory wolly	Spanish needles	Yarrow western
	Mugwort	Speedwell bi lobed	Yarrow yellow

[Back Panel] [Restatement of information in sections 1 and 2]

Getting Started

Kills

Crabgrass foxtail and most troublesome broadleaf weeds including dandelion clover thistle and over XXX other weeds listed

Where To Use

On established lawns without harming lawn grasses when used as directed (Do not use on Bahiagrass Bentgrass (colonial and creeping) Carpetgrass Centipedegrass Dichondra Seashore Paspalum St Augustinegrass and turfgrass species that are not listed on this label) [See Inside Label for Complete Directions]

Amount To Use

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass Perennial Ryegrass & Zoysiagrass
Entire Lawn Treatments This container covers XXX sq ft

Coverage

This [size] bottle treats XXXX sq ft

Telephone Icon

Computer Icon

For questions or comments call toll-free 1-800 821-7925 or visit our website at www.pbigordon.com

APPENDIX

1 Statements which may appear on different label components depending on packaging configuration

- See next panel for additional Precautionary Statements and First Aid
- Net Contents _____
- EPA Est No _____

2 Advertising claims that may be presented on the retail containers of supplemental registrants

- One quart covers up to 5 000 square feet
- 1 quart treats [up to] 5 000 sq ft
- Covers [XXXX] sq ft
- Keep from freezing
- Controls crabgrass dandelion chickweed knotweed henbit plantain clover spurge and many other broadleaf lawn weeds including those shown on the side panel
- Controls crabgrass and broadleaf weeds in lawns
- Kills even the tough weeds roots and all!
- Controls even the tough weeds roots and all!
- Kills the root
- Water based
- Won t harm lawn grass when used according to label directions
- Kills lawn weeds plus crabgrass
- Kills [Controls] 200+ broadleaf weeds
- Kills the weeds not the lawn
- Attach to [garden] hose and spray
- Simple to use
- Easy to use
- Easy to mix
- Proven performance
- For home lawn care
- Won t harm lawn grass when used as directed
- Controls (Number) weeds!
- Mixes with Water Concentrate
- Kill crabgrass & [any listed weed(s)] too!
- **CONTROLS**
 - SENSITIVE WEEDS** Crabgrass Bedstraw Black Medic Buckhorn Chickweed Clover Dandelion Knotweed Lambsquarters Peppergrass Plantain Purslane Ragweed and Wild Lettuce
 - HARD-TO-KILL WEEDS** Chicory Creeping Charlie Creeping Jenny Dock Ground Ivy Heal-All Lespedeza Mallow Morningglory Oxalis Pigweed Poison Ivy Poison Oak Sheep Sorrel Speedwell Spurge Wild Carrot Wild Onion Yarrow and other broadleaf weeds
- Kills Lawn Weeds plus Crabgrass in One Step
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grass Weeds
- [Also] Kills [Dandelion] [Crabgrass] [Clover] plus 200 Other [Listed] Weeds [As Listed]
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Certain Weed Grasses

- [XX] [Pint Quart etc] Treats Up To [YYYY] Sq Ft
- Local Solutions for Local Problems
- For more Lawn and Garden Products Visit [www website com]
- Contains Trimec Herbicide
- Kills Crabgrass and Broadleaf weeds roots and all!
- XX covers up to XXXX square feet
- Starts working overnight
- Won t harm lawn grass when used according to label directions
- Won t harm lawn grass [when used as directed]
- Won t harm lawns* *When used as directed
- For outdoor use
- Bonus Size
- 25% More Free!
- Kills Crabgrass and over XXX Weeds in Country & Suburban Lawns* *See inside booklet for a complete list of weeds controlled
- This product kills Crabgrass PLUS broadleaf weeds The special Trimec® dual-action formula goes beyond crabgrass and foxtail to kill over XXX other broadleaf weeds listed in country and suburban lawns
- Before/After
- Convenient [Simple] [Design]
- Easily covers large areas
- [Easy] [Ergonomic] Comfort Grip [Design]
- Easy grip [control]
- Easy [to use]
- Economical – use to cover large areas
- Ergonomic [design]
- Gentle shower [applicator] [spray pattern]
- Hand doesn t get wet
- Hand stays dry
- Includes water [only] setting/option
- Instant/Easy [One-handed] on/off applicator/switch/action
- No mixing required
- One-handed operation
- Ready-Spray®/Ready-to-Spray/Applicator Name
- Simple [and easy] [to use]
- [Simply] attach to [garden] hose and spray/go

For any of the ORTHO Products

- Makes more than XX Gallon!
- Kills Even Tough Weeds – Roots & All
- Lawn Weed & Crabgrass Killer
- Kills Dandelions Clover and Other Broadleaf Weeds! Plus Kills Crabgrass and Other grass Weeds!
- Kills the Toughest Weeds in these Lawns (any lawn listed on label)
- Open Resealable Label for Directions & Precautions (Press to Reseal)
- People or pets may reenter treatment area after spray has dried
- Ready-Spray
- Do Not Use on St Augustine Lawns
 - Ortho Weed-B-Gon MAX Concentrate Plus Crabgrass Control [Killer]
 - Kills Weeds Not Lawns(TM)
 - For [Use on] Lawns

- ORTHO & Your Environment - Your yard is a place for your family and pets to enjoy That s why ORTHO products are designed with care to provide effective solutions to weed problems For best results please follow instructions for appropriate usage storage and disposal
- Kills Over 200 Lawn Weeds plus [including] Crabgrass [in one easy step
- Guaranteed [Results]
- See Ortho com for complete Spanish Translation
- DO NOT use on St Augustine including Floratam species
- NOTICE Buyer assumes all risks of use storage or handling of this product not in accordance with directions
- Rainfast in 1 Hour
- Kills Major [Lawn] Broadleaf Weeds
- Rainproof in 1 Hour
- Easy-Measure [Cap] [Top] [Lid] [Cup]
- Drain-Back [Cap] [Top] [Lid] [Cup]

For ORTHO Weed-B-Gon MAX Plus Crabgrass Control [Killer] Singles

- NET 5 FL OZ
- Pre-Measured Concentrate Bottles
- Treats 5 000 sq ft
- No Measuring No Mess
- Weed-B-Gon® Max(TM) Singles makes using concentrates easier than ever No more measuring and no more mess Mix (contents) (each pre-measured bottle) with two gallons of water to deliver consistent results And its revolutionary formula won t harm your lawn - Guaranteed
- Designed for tank sprayers
- 1 bottle makes 2 gallons

3 Trademark acknowledgements

- Ortho® Dial N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H D Hudson Manufacturing Company
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company
- Trimec® is registered trademark of PBI/Gordon Corporation

DOCUMENT CONTROL INFORMATION

1 Unique Label Identifier 002217-00896 20120705 amend-proposed-clean doc

2 Reason for Issue new sublabels alternate warranty ad claims