

2217-8916

11/13/2013

11/27



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

November 13, 2013

James L. Kunstman, Ph.D.
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, MO 64101

Subject: Notification per PR Notice 98-10 (ABNs, marketing claim, contact info)
EH-1437 Herbicide
EPA Reg. No. 2217-896
Application Dated June 27, 2013

Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product. The Registration Division (RD) has conducted a review of this request and finds that the action falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped 'Notification' and will be placed in our records.

In addition, the following alternate brand names have been added to our records:

"Trimec Crabgrass Plus Lawn Weed Killer Ready Spray"
"Trimec Crabgrass Plus Lawn Weed Killer Concentrate"


If you have any questions, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Mindy Ondish, for".

Kathryn V. Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

2/27

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
----------------------------------------------------------------------------------	---------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 2217-896	2. EPA Product Manager Kathryn V. Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) EH-1437 Herbicide	PM# Product Manager—Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	NOV 13 2013
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments. Section II(A),(N), Section IV(F)7.



Please refer to cover letter for details and certification statement.

e-mail to jkunstman@pbigordon.com FAX: 816-421-2731

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per Container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 1 pint, 1 quart, 2 quarts, 1 gallon	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name James L. Kunstman, Ph.D.	Title Director of Regulatory Services	Telephone No. (Include Area Code) 816-460-6292
Certification <i>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</i>		6. Date Application Received (Stamped) 
2. Signature 	3. Title Director of Regulatory Services	
4. Typed Name James L. Kunstman, Ph.D.	5. Date June 27, 2013	



1217 W. 12th STREET · P.O. BOX 014090
KANSAS CITY, MISSOURI 64101-0090
816-421-4070 · 1-800-821-7925
FAX: 816-474-0462

June 27, 2013

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Kathryn V. Montague (PM-23)

NOTIFICATION: Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments. Section II(A),(N) and IV(F)7 - for EH-1437 Herbicide (EPA Reg. No. 2217-896).

I. Labeling notification:

Section II (A): Brand Names

- We ask to add two new alternate brand names.
 - Trimec Crabgrass Plus Lawn Weed Killer Ready Spray
 - Trimec Crabgrass Plus Lawn Weed Killer Concentrate

Section II (N): Other Revisions

- We ask to add advertising claims that are consistent with the Directions for Use previously approved and re-state an existing advertising claim to make it more explicit. Please refer to pages 21 and 22 of the draft labeling.

Section IV (F) 7: Consumer Access Numbers

- Update contact information as seen on pages 16 and 20 of the enclosed draft labeling.

II. Certification Statement

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA:

6/27

Label Language Common To All Sublabels

EH-1437 HERBICIDE

EPA Reg. No. 2217-896

Alternate Brand Names:

- Tractor Supply Co. GroundWork Concentrate Crabgrass & Broadleaf Weed Killer
- Gordon's Trimec Crabgrass Plus Lawn Weed Killer Ready Spray
- Gordon's Trimec Crabgrass Plus Lawn Weed Killer Concentrate
- Trimec Crabgrass Plus Lawn Weed Killer Ready Spray
- Trimec Crabgrass Plus Lawn Weed Killer Concentrate

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	6.42%
Quinclorac	2.13%
Dicamba, dimethylamine salt	0.60%
INERT INGREDIENTS:	90.85%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.456 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
 - 0.182 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
 - 0.043 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
- Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately.

before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations	
<ul style="list-style-type: none"> • Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)]

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

DIRECTIONS FOR USE

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area

inaccessible to children and pets. Avoid contamination of food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.


Alternate Warranty:

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for lack of performance or safety if not used according to the label directions.

[OPTIONAL WARRANTY STATEMENT]

The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.




SUBLABEL 1 Traditional layout

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
-  Optional Icon] Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)

When to Use: Apply when daytime temperatures are below 90°F.

Questions & Comments: (icon phone) (icon computer)
Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com

HOW IT WORKS:

EH-1437 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE:

EH-1437 Herbicide can be applied to residential lawns with the following turfgrass species:

Lawns To Be Treated:	
Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass	Bermudagrass ¹ , common and hybrid
Ryegrass, perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. ^cDc not apply to bermudagrass when daytime temperatures exceed 85° F.

DO NOT apply EH-1437 Herbicide to the following:

Bahiagrass	
Bentgrass, colonial and creeping	
Carpetgrass	
Centipedegrass	
Dichondra	
Gardens including vegetables, fruit trees, vines and berries	
Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs)	
Seashore paspalum	

St. Augustinegrass
Turfgrass species that are not listed on this label.

GRASS AND BROADLEAF WEEDS CONTROLLED:

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Mustard	Spiny cocklebur
Aster	Common mullein	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Bedstraw	Cudweed	Nettle	Spotted spurge
Beggarticks	Curly dock	Orange hawkweed	Spurweed
Betony, Florida	Daisy, English	Oriental cocklebur	Stinging nettle
Bindweed, field	Daisy fleabane	Oxalis	Strawberry, India mock
Bird vetch	Daisy, oxeye	Parsley-piert	Tall nettle
Bitter wintercress	Dandelion	Parsnip	Tall vervain
Bittercress, hairy	Dichondra	Pearlwort	Tansy ragwort
Bitterweed	Dogbane	Pennycress	Tansy mustard
Black-eyed Susan	Dogfennel	Pennywort	Tanweed
Black medic	Dollarweed	Peppergrass	Thistle
Black mustard	Elderberry	Pepperweed	Trailing crownvetch
Blackseed plantain	False dandelion	Pigweed	Tumble mustard
Blessed thistle	Falseflax	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	Fiddleneck	Plantain	Venice mallow
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Water pennywort
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly lettuce	Western clematis
Broomweed	Gumweed	Prickly sida	Western salsify
Buckhorn	Hairy fleabane	Prostrate knotweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate pigweed	Wild aster
Bulbous buttercup	Healall	Prostrate spurge	Wild buckwheat
Bull thistle	Heartleaf drymary	Prostrate vervain	Wild carrot
Bullnettle	Heathaster	Puncturevine	Wild four-o'clock
Burclover	Hedge bindweed	Purslane, common	Wild garlic
Burdock	Hedge mustard	Ragweed	Wild geranium
Burning nettle	Hemp	Red sorrel	Wild lettuce
Burweed	Henbit	Redroot pigweed	Wild marigold
Buttercup	Hoary cress	Redstem filaree	Wild mustard
Buttonweed	Hoary plantain	Rough cinquefoil	Wild onion
Canada thistle	Hoary vervain	Rough fleabane	Wild parsnip
Carolina geranium	Horsenettle	Russian pigweed	Wild radish
Carpetweed	Jimsonweed	Russian thistle	Wild rape
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild strawberry
Catnip	Knotweed	Scotch thistle	Wild sweet potato
Catsear	Kochia	Sheep sorrel	Wild vetch
Chickweed, common	Lambsquarters	Shepherdspurse	Wild violet
Chickweed, mouseear	Lespedeza	Slender plantain	Woodsorrel
Chicory	Mallow	Smallflower galinsoga	Woolly cotton
Cinquefoil	Matchweed	Smooth dock	Woolly morningglory
Clover, crimson	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, hop	Milk vetch	Sorrel	Wormseed
Clover, red	Morningglory	Sowthistle	Yarrow
Clover, strawberry			Yellow rocket

BROADLEAF WEEDS			
Clover, sweet Clover, white Cockle	Mouseear hawkweed Mugwort Musk thistle	Spanishneedles Speedwell Spiny amaranth	Yellowflower pepperweed and other broadleaf weeds

WHEN TO USE:

Apply this product when weeds are small and actively growing.

Weed species	Season	Timing	Maximum number of applications ¹	Application interval, days	Comments
Crabgrass (large and smooth):	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height	2	14	Re-treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations.
Broadleaf signalgrass	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	14	Re-treat at the same rate at 14 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

¹The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 applications per year. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre]

HOW MUCH TO USE:

I. ENTIRE LAWN TREATMENTS WITH DIAL TYPE HOSE-END SPRAYERS

Three Easy Steps To Mix And Apply:

1. FILL

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- The application rate for this product is 6.4 fl.oz. of product per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 6.4 fl.oz. of product for every 1,000 sq.ft. of lawn area. Over-application or rates above those recommended on this label can cause turf injury.
- Do not add water to the sprayer jar.

2. SET THE DIAL

For hose-end sprayers: Set the dial to the rate per gallon and keep the dial set to this setting while spraying.

The dial setting or rate per gallon for this product is 2.5 oz. of product per gallon of water.

Types of hose-end sprayers	Dial setting
Ortho® Dial'N Spray® Hose-End Sprayer	2.5 oz.
Hudson® Hose End Liquid Concentrate Sprayer	2.5 fl.oz.
Gilmour® Dial-A-Mix® Lawn and Garden Sprayer	15 Teaspoons or 5 tablespoons

3. **SPRAY** - Apply evenly over the lawn.

II. ENTIRE LAWN TREATMENTS WITH READY-TO-USE HOSE-END SPRAYERS :

This product may be packaged as a ready-to-use product with no mixing. The retail plastic container is fitted with a sprayer body that delivers spray droplets at a fixed rate.

For READY-TO-USE Hose End Sprayers:

- This retail sprayer bottle will cover the treated areas described below when this product is applied at 6.4 fl.oz. of product per 1,000 sq.ft. Over-application or rates above those recommended on this label can cause turf injury.

Approximate treated areas for four retail container sizes:

Species	Amount of product to treat approximate area, sq. ft.				
	Fl.oz./1,000 sq.ft.	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container	2.0 qt. (64 fl.oz.) retail container	1 gallon (128 fl.oz.) retail container
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season species Bermudagrass ¹ common and hybrid Zoysiagrass and Buffalograss	6.4 fl.oz. /1,000 sq.ft.	2,500 sq.ft.	5,000 sq.ft.	10,000 sq.ft.	20,000 sq.ft.

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Instructions:

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.

Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl.oz. container will treat up to xxxx sq.ft. of turfgrass.

- **FINISH:** To stop spraying, turn [rotate] spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn [rotate] spray nozzle to the "ON" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

III. HAND-OPERATED SPRAYERS:

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up):

- Apply to actively growing grass and broadleaf weeds during the early growth stages.
- Measure and mix the proper amount of product with water as described below. Mix 2.5 fl.oz. of product with 1 gallon of water to treat 400 sq.ft. of lawn. See below for additional spray preparations
- Spray evenly over the lawn. Spray coverage should be uniform and complete. Over-application or rates above those recommended on this label can cause turf injury.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square feet
2.5 fl.oz.	1 gallon	400 sq.ft.
3.75 fl.oz.	1.5 gallons	600 sq.ft.
5.0 fl.oz.	2 gallons	800 sq.ft.
7.5 fl.oz.	3 gallons	1,200 sq.ft.

Equal measures: 2.5 fl.oz. = 5 Tablespoons. 1 cup = 8 fl.oz. = 1/2 pint.

OR

Measure indicated amount of product and pour into tank. Dilute with water according to the following chart. Adjust the nozzle to deliver coarse spray droplets and treat the area indicated.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Area To Treat, Square feet
2.5 fl.oz.	1 gallon	400 sq.ft.
5 fl. fl.oz.	2 gallons	800 sq.ft.
7.5 fl.oz.	3 gallons	1,200 sq.ft.

Equal measures: 2.5 fl.oz. = 5 Tablespoons. 1 cup = 8 fl.oz. = 1/2 pint.

IV. ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® SYSTEM:

HOW TO USE THE READY-SPRAY® SYSTEM

Connect:

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.
4. Extend hose to farthest area of the lawn to be treated

Spray:

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small plastic tab back and turn knob clockwise to ON position.
3. Spray evenly over measured area.

Finish:

1. To STOP spraying, rotate the knob counterclockwise to the OFF position.
2. Turn off water.
3. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.
4. Disconnect sprayer from hose.

HOW TO USE THE ORTHO® DIAL-N-SPRAY® APPLICATOR WITH DIRECT CONNECT [Applicator Name]

[Prepare Applicator:] [illustration]

1. Set dial to [X] fl.oz./Tbsp./tsp.

[Connect:] [illustration]

1. Remove cap from the concentrate bottle. [Keep cap for storage.]
2. Insert suction tube into bottle opening and twist the Ortho® Dial 'N Spray® applicator [applicator name] onto bottle to attach.
3. Connect hose to the sprayer and turn on water.

[Spray:] [illustration] [directions will vary by product type]

1. Hold sprayer in upright position and apply evenly to area to be treated.
2. Spray only until the surface of the lawn is wet.

[Finish:] [illustration]

1. After use, untwist to remove the [insert product name] concentrate bottle from the Ortho® Dial 'N Spray® applicator [applicator name].
2. Replace cap on the [insert product name] concentrate bottle.

Refer to the Ortho® Dial 'N Spray applicator [applicator name] booklet for additional information. Important: To prevent sprayer from clogging, always clean after use as directed.

V. FOR TANK SPRAYER:

- Add 2.5 fl.oz. per gallon of water for each 400 sq.ft.

For broadcast application: Ortho Dial 'N Spray® Applicator

1. Add 16 fl.oz. of concentrate directly into sprayer jar. DO NOT ADD WATER.
2. Set dial to 2.5 oz.
3. Spray evenly over 2,500 sq.ft.
4. Any unused portion should be poured back into its original container.

[Over-application or rates above those recommended on this label can cause turf injury]

VI. FOR ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL [KILLER] SINGLES:

For (Use a) Tank Sprayer:

- Add entire contents (of 1 bottle) (5 oz.) to an empty 2-gallon tank sprayer. Slowly fill with 2 gallons of clean tap water.
- **For Broadcast Application – Ortho® Dial 'N Spray® Applicator**
 1. Set dial to 2.5 oz.
 2. Add entire contents of 1 bottle (5 oz) directly into sprayer jar. DO NOT ADD WATER.
 3. Spray evenly over 800 sq.ft.

[Over-application or rates above those recommended on this label can cause turf injury]

VII. ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® II [2]/APPLICATOR NAME SYSTEM:

HOW TO USE THE READY-SPRAY® II/[READY-SPRAY 2]/[APPLICATOR NAME] SYSTEM

[Optional]

1. [Measure the total [entire] area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.]

[End of optional section]

Connect:


1. Shake well before using. [optional step]
2. Connect sprayer to hose.
3. Turn on water.
4. Extend hose to farthest area of the measured area to be treated. [optional step]

Spray:

1. To begin spraying, [disengage safety lock and] point nozzle in the direction you want to spray.
2. Slide thumb activation switch forward to ON position.
3. Spray evenly over measured area.

Finish:


1. To STOP spraying, slide thumb activation switch backward to OFF position.
2. Turn off water.



3. Relieve water pressure by sliding thumb switch to ON /WATER/ WATER ONLY position until water slows to a drip. Then, slide switch back to OFF position [and engage safety lock].
4. Disconnect sprayer from hose.

[Optional illustrations may be used in conjunction with the above steps]

CULTURAL TIPS AND NOTES:		
When to Apply:	For established lawns For newly seeded areas For newly sodded, sprigged, or plugged areas	<ol style="list-style-type: none"> 1. Apply when daytime temperatures are below 90°F. 2. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing. 3. Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
Seeding:		<ul style="list-style-type: none"> • The treated area can be seeded at 4 weeks after the application of this product.
Irrigation:	Non-stressed turfgrass Stress from high temperatures, drought, insects or diseases.	<ul style="list-style-type: none"> • Adequate soil moisture and favorable growing conditions enhance the performance of this product. • Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing:		<ul style="list-style-type: none"> • Delay mowing 2 days before and until 2 days after the application of this product. • Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
Nontarget plants:		<ul style="list-style-type: none"> • Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. • Do not spray exposed roots of ornamentals and trees.



**SUBLABEL 2
Concentrate**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Before You Use. Read & follow these directions when using.

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Will not harm lawn grasses when used as directed. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Spot Treatments: Mix 2.5 fl.oz. (5 Tbs) in 1 gallon of water to treat 400 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Over-application or rates above those specified on this label can cause turf injury.

3. When To Use

Apply this product when weeds are small and actively growing.

Weed species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re-treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 14 days later for dense populations.
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 14 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

Weed species	Timing	Comments
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture
The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 14 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre]		

4. How To Apply

4.1 Pump Style Sprayers [sprayer icon]

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

4.2 [12 Volt &] Tow-Behind Boom Sprayers [sprayer icon]

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq.ft. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Tank Size	Spray volume	Coverage Area	Concentrate to add to tank
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft.	192 fl.oz. (1.5 gal)
	1 gal/1000 sq.ft.	15,000 sq.ft.	96 fl.oz. (0.75 gal)
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft.	320 fl.oz. (2.5 gal)
	1 gal/1000 sq.ft.	25,000 sq.ft.	160 fl.oz. (1.25 gal)

For spot treatments with [12 Volt &] Pull Behind Sprayers

See rates under "How Much To Apply," Spot Treatments.

4.3 Dial Style Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER.**
2. Close sprayer and set dial to 2.5 oz.
3. Connect sprayer to hose
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying; avoiding contact with the treated area.
5. Pour any unused product back into this container.

4.4 Concentrate Plus Water Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar.
3. Close sprayer, connect to garden hose, shake well.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

5. For Best Results

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.


6. Grass and Broadleaf Weeds Killed

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Mustard	Spiny cocklebur
Aster	Common mullein	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear
Bedstraw	Cudweed	Nettle	Spotted spurge
Beggarticks	Curly dock	Orange hawkweed	Spurweed
Betony, Florida	Daisy, English	Oriental cocklebur	Stinging nettle
Bindweed, field	Daisy fleabane	Oxalis	Strawberry, India mockc
Bird vetch	Daisy, oxeye	Parsley-piert	Tall nettle c c c c c
Bitter wintercress	Dandelion	Parsnip	Tall vervain c c c c c
Bittercress, hairy	Dichondra	Pearlwort	Tansy ragwort c c c c c
Bitterweed	Dogbane	Pennycress	Tansy mustard c c c c c
Black-eyed Susan	Dogfennel	Pennywort	Tanweed c c c c c
Black medic	Dollarweed	Peppergrass	Thisle c c c c c
Black mustard	Elderberry	Pepperweed	Trailing crownvetch c c c c c
Blackseed plantain	False dandelion	Pigweed	Tumble mustard c c c c c
Blessed thistle	Falseflax	Pineywoods bedstraw	Tumble pigweed c c c c c
Bloodflower milkweed	False sunflower	Plains coreopsis (tickseed)	Velvetleaf c c c c c
Blue lettuce	Fiddleneck	Plantain	Venice mallow c c c c c
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed c c c c c
Bracted plantain	Frenchweed	Poison oak	Virginia creeper c c c c c
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed c c c c c
Bristly oxtongue	Goathead	Poorjoe	Water pennywort c c c c c
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle c c c c c
Broadleaf plantain			Western clematis c c c c c

BROADLEAF WEEDS			
Broomweed	Ground ivy	Prickly lettuce	Western salsify
Buckhorn	Gumweed	Prickly sida	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild buckwheat
Bull thistle	Healall	Prostrate spurge	Wild carrot
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild four-o'clock
Burclover	Heathaster	Puncturevine	Wild garlic
Burdock	Hedge bindweed	Purslane, common	Wild geranium
Burning nettle	Hedge mustard	Ragweed	Wild lettuce
Burweed	Hemp	Red sorrel	Wild marigold
Buttercup	Henbit	Redroot pigweed	Wild mustard
Buttonweed	Hoary cress	Redstem filaree	Wild onion
Canada thistle	Hoary plantain	Rough cinquefoil	Wild parsnip
Carolina geranium	Hoary vervain	Rough fleabane	Wild radish
Carpetweed	Horsenettle	Russian pigweed	Wild rape
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild strawberry
Catnip	Knawel	Scarlet pimpernel	Wild sweet potato
Catsear	Knotweed	Scotch thistle	Wild vetch
Chickweed, common	Kochia	Sheep sorrel	Wild violet
Chickweed, mouseear	Lambsquarters	Shepherdspurse	Woodsorrel
Chicory	Lespedeza	Slender plantain	Woolly croton
Cinquefoil	Mallow	Smallflower galinsoga	Woolly morningglory
Clover, crimson	Matchweed	Smooth dock	Woolly plantain
Clover, hop	Mexicanweed	Smooth pigweed	Wormseed
Clover, red	Milk vetch	Sorrel	Yarrow
Clover, strawberry	Morningglory	Sowthistle	Yellow rocket
Clover, sweet	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, white	Mugwort	Speedwell	and other broadleaf
Cockle	Musk thistle	Spiny amaranth	weeds



[Back Panel] [Restatement of information in sections 1 and 2.]

Getting Started

Kills:

Crabgrass, foxtail and most troublesome broadleaf weeds including dandelion, clover, thistle and over XXX other weeds listed.

Where To Use:

On established lawns without harming lawn grasses when used as directed (Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipede grass, Dichondra, Seashore Paspalum, St. Augustine grass, and turfgrass species that are not listed on this label.) [See Inside Label for Complete Directions]

Amount To Use:

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:

Spot Treatments: Mix 2.5 fl.oz. (5 Tbs) in 1 gallon of water to treat 400 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft. [See Inside Label for Sprayer Settings]
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.


Coverage:

This [size] bottle treats XXXX sq.ft.

Telephone Icon

Computer Icon

For questions or comments call toll-free 1-800-821-7925 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST or visit [our website at] www.pbigordon.com [GordonsUSA.com]



SUBLABEL 3
Ready Spray

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Before You Use. Read & follow these directions when using.

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Will not harm lawn grasses when used as directed. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewing), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Entire Lawn Treatments: This container covers XXX sq.ft. Measure the size of your yard. If the area to be treated is larger than XXX sq.ft., you may need to buy more containers.

Over-application or rates above those specified on this label can cause turf injury.

3. When To Use

Apply this product when weeds are small and actively growing.

Weed species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re-treat at the same rate at 14 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 14 days later for dense populations.
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 14 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture

The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 14 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre]

4. How To Apply

1. Connect your garden hose to the hose-end sprayer. With your thumb, snap off the safety locking tab that prevents accidental discharge of spray solution. Leave control valve in the OFF position.
2. To BEGIN spraying: Turn on water at faucet. Extend hose to the furthest point from the faucet. Point nozzle toward the lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Walk backwards as you apply in order to avoid contact with the treated area. For best results, spray evenly over the measured area.
3. To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

5. For Best Results

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- Certain hard-to-kill weeds may need to be spot treated in 14 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

6. Grass and Broadleaf Weeds Killed

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Amaranth, purple	Cupid shaving Brush	Mustard, ball	Speedwell, common
Amaranth, slender	Cudweed	Mustard, black	Speedwell, corn
Aster, heath	Daisy, English	Mustard, blue	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, hedge	Speedwell, persian
Aster, purple	Dandelion, common	Mustard, Indian,	Speedwell, purslane
Aster, white prairie	Dandelion, false	Mustard, tansy	Speedwell, slender
Artichoke	Day flower	Mustard, tumble	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, white	Speedwell, thymeleaf
Bedstraw, Catchweed	Dichondra	Mustard, wormseed	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Nettle, stinging (burning)	Spurge, spotted
Bedstraw, Northern	Dock, curly	Nettle, tall	Spurge, prostrate
Bedstraw, pineywoods	Dogbane	Nettle, wood	Spurweed
Beggarticks, hairy	Dogfennel	Parsley-piert	Star of Bethlehem
Betony, Florida	Dollarweed (Pennywort)	Parsnip	Strawberry, and a flock
Bindweed, Field	Dove weed	Pearlwort	Tansy ragwort
Bindweed, Hedge	Elderberry	Pennycress (frenchweed)	Tanweed
Bitter Wintercress	Falseflax	Pepperweed, perennial	Thistle, blessed
Bittercress, hairy	False sunflower	Pepperweed, virginia	Thistle, bull
Bitterweed	Fiddleneck	(Peppergrass)	Thistle, canada
Black-eyed Susan	Filaree, redstem	Pigweed, prostate	Thistle, flodman (prairie)

BROADLEAF WEEDS			
Black Medic	Filaree, whitestem	Pigweed, redroot	Thistle, musk
Blood flower milkweed	Fleabane, daisy (annual)	Pigweed, russian	Thistle, platte
Blue lettuce	Fleabane, rough	Pigweed, smooth	Thistle, plumeless
Brassbuttons	Galinsoga, hairy	Pigweed, tumble	Thistle, russian
Bristly oxtongue	Galinsoga, smallflower	Pineapple weed	Thistle, scotch
Broomweed	Goldenrod	Plains coreopsis (tickseed)	Thistle, wavyleaf
Buckhorn	Ground ivy	Plantain, blackseed	Thistle, yellow (horrible)
Bullnettle	Groundsel	Plantain, bracted	Trailing crownvetch
Burclover	Gumweed	Plantain, broadleaf	Velvetleaf
Burdock	Hairy fleabane	Plantain, buckhorn	Vervain, blue
Burweed, lawn (spurweed)	Hawkweed, meadow	Plantain, hoary	Vervain, hoary
Buttercup, bulbous	Hawkweed, mouseear	Plantain, narrowleaf	Vervain, prostrate
Buttercup, creeping	Hawkweed, orange	Plantain, slender	Vervain, tall
Buttonweed, common	Hawkweed, yellow	Plantain, woolly	Vetch, bird
(poorjoe)	Healall	Poison ivy	Vetch, hairy
Buttonweed, virginia	Heartleaf drymary	Poison oak	Vetch, milk
Carolina geranium (wild)	Hemp	Pokeweed	Vetch, narrowleaf
Carpetweed	Henbit	Prairie Sunflower	Vetch, wild
Catnip	Hoary cress	Prickly lettuce (compass plant)	Virginia creeper
Catsear, spotted (false dandelion)	Horsenettle	Prickly sida	Western clematis
Chamber bitter	Horseweed	Prostrate knotweed	Western salsify
Chickweed, common	Jimsonweed	Puncturevine	Wild aster
Chickweed, mouseear	Johnny-Jumpup violet	Purslane, common	Wild buckweat
Chickweed, sticky	Knawel	Pusley, brazil	Wild carrot
Chicory	Knotweed, prostrate	Pusley, florida	Wild four-o'clock
Cinquefoil	Kochia	Pusley, large flower	Wild garlic
Clover, crimson	Lambsquarters	Ragweed, bur	Wild lettuce
Clover, alyce	Lespedeza	Ragweed, common	Wild marigold
Clover, hop	Mallow, alkali	Ragweed, lancleaf	Wild mustard (yellow rocket)
Clover, rabbit foot	Mallow, bristly	Ragweed, western	Wild onion
Clover, red	Mallow, common	Redstem filaree	Wild parsnip
Clover, strawberry	Mallow, dwarf	Rough cinquefoil	Wild radish
Clover, yellow sweet	Mallow, venice	Scarlet pimpernel	Wild rape
Clover, white sweet	Marcela	Shepardspur	Wild strawberry
Clover, white	Matchweed	Smartweed, ladysthumb	Wild sweet potato
Cockle	Mexicanweed	Smartweed, pale	Wild violet
Cocklebur	Moneywort	Smartweed, pennsylvania	Woodsorrel, creeping
Cocklebur, oriental	Morningglory, bigroot	Smooth chaff-flower	Woodsorrel, violet
Cocklebur, spiny	Morningglory, ivyleaf	Smoothdock	Woodsorrel, yellow (Oxalis)
Common mullein	Morningglory, mexican	Sorrel, red (sheep)	Woolly croton
Corn Spurry	Morningglory, red	Sowthistle, annual	Wormseed
Creeping Beggarweed	Morningglory, tall	Sowthistle, spiny	Yarrow, common
Creeping Jenny (creeping charlie)	Morningglory, wolly	Spanish needles	Yarrow, western
	Mugwort	Speedwell, bi-lobed	Yarrow, yellow

[Back Panel] [Restatement of information in sections 1 and 2.]

Getting Started

Kills:

Crabgrass, foxtail and most troublesome broadleaf weeds including dandelion, clover, thistle and over XXX other weeds listed.

Where To Use:

On established lawns without harming lawn grasses when used as directed (Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.) [See Inside Label for Complete Directions]

Amount To Use:

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:

Entire Lawn Treatments: This container covers XXX sq.ft.


Coverage:

This [size] bottle treats XXXX sq.ft.

Telephone Icon

Computer Icon

For questions or comments call toll-free 1-800-821-7925 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST or visit [our website at] www.pbigordon.com [GordonsUSA.com]



APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on the retail containers of supplemental registrants.


- One quart covers up to 5,000 square feet.
- 1 quart treats [up to] 5,000 sq.ft.
- Covers [XXXX] sq.ft.
- Keep from freezing.
- Controls crabgrass, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Controls crabgrass and broadleaf weeds in lawns
- Kills even the tough weeds...roots and all!
- Controls even the tough weeds...roots and all!
- Kills the root.
- Water based.
- Won't harm lawn grass when used according to label directions.
- Kills lawn weeds plus crabgrass.
- Kills [Controls] 200+ broadleaf weeds.
- Kills the weeds not the lawn
- Attach to [garden] hose and spray
- Just connect to garden hose & spray
- Simple to use
- Easy to use
- Easy to mix
- Proven performance
- For home lawn care
- Won't harm lawn grass when used as directed.
- Controls (Number) weeds!
- Mixes with Water - Concentrate
- Kill crabgrass & [any listed weed(s)] too!

CONTROLS

SENSITIVE WEEDS: Crabgrass, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.

HARD-TO-KILL WEEDS: Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and other broadleaf weeds.


- Kills Lawn Weeds plus Crabgrass in One Step
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grass Weeds
- [Also] Kills [Dandelion], [Crabgrass], [Clover], plus 200 Other [Listed] Weeds [As Listed]
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds



- Effectively Kills Both Broadleaf and Certain Weed Grasses
- [XX] [Pint, Quart, etc] Treats Up To [YYYY] Sq.Ft.
- Local Solutions for Local Problems
- For more Lawn and Garden Products Visit [www.website.com]
- Contains Trimec Herbicide
- Kills Crabgrass and Broadleaf weeds...roots and all!
- XX covers up to XXXX square feet.
- Starts working overnight
- Won't harm lawn grass when used according to label directions.
- Won't harm lawn grass [when used as directed]
- Won't harm lawns* *When used as directed
- For outdoor use
- Bonus Size
- 25% More Free!
- Kills Crabgrass and over XXX Listed Weeds in Country & Suburban Lawns* *See inside booklet for a complete list of weeds controlled
- This product kills Crabgrass PLUS broadleaf weeds. The special Trimec® dual-action formula goes beyond crabgrass and foxtail to kill over XXX other broadleaf weeds listed in country and suburban lawns.
- Before/After
- Convenient [Simple] [Design]
- Easily covers large areas
- [Easy] [Ergonomic] Comfort Grip [Design]
- Easy grip [control]
- Easy [to use]
- Economical – use to cover large areas
- Ergonomic [design]
- Gentle shower [applicator] [spray pattern]
- Hand doesn't get wet
- Hand stays dry
- Includes water [only] setting/option
- Instant/Easy [One-handed] on/off applicator/switch/action
- No mixing required
- One-handed operation
- Ready-Spray®/Ready-to-Spray/Applicator Name
- Simple [and easy] [to use]
- [Simply] attach to [garden] hose and spray/go

For any of the ORTHO Products:

- Makes more than XX Gallon!
- Kills Even Tough Weeds – Roots & All
- Lawn Weed & Crabgrass Killer
- Kills Dandelions, Clover, and Other Broadleaf Weeds! Plus Kills Crabgrass and Other grass Weeds!
- Kills the Toughest Weeds in these Lawns (any lawn listed on label)
- Open Resealable Label for Directions & Precautions (Press to Reseal)
- People or pets may reenter treatment area after spray has dried.
- Ready-Spray
- Do Not Use on St. Augustine Lawns
 - Ortho Weed-B-Gon MAX Concentrate Plus Crabgrass Control [Killer]
 - Kills Weeds, Not Lawns(TM)
 - For [Use on] Lawns



- ORTHO & Your Environment - Your yard is a place for your family and pets to enjoy. That's why ORTHO products are designed with care to provide effective solutions to weed problems. For best results please follow instructions for appropriate usage, storage and disposal.
- Kills Over 200 Lawn Weeds plus [including] Crabgrass [in one easy step.
- Guaranteed [Results]
- See Ortho.com for complete Spanish Translation.
- DO NOT use on St. Augustine, including Floratam species.
- NOTICE: Buyer assumes all risks of use, storage, or handling of this product not in accordance with directions
- Rainfast in 1 Hour
- Kills Major [Lawn] Broadleaf Weeds
- Rainproof in 1 Hour
- Easy-Measure [Cap] [Top] [Lid] [Cup]
- Drain-Back [Cap] [Top] [Lid] [Cup]
- Direct connect to the Ortho® Dial 'N Spray®
- Direct connect to Ortho® Dial 'N Spray® means no more measuring
- No more measuring concentrates with Ortho® Dial 'N Spray®
- No concentrate contact with Ortho® Dial 'N Spray®
- No over/under application with Ortho® Dial 'N Spray®
- Exact measurement and improved efficacy with Ortho® Dial 'N Spray®
- Simple, direct connect capability with Ortho® Dial 'N Spray®
- No measure, direct connect capability with Ortho® Dial 'N Spray®
- No over/under application, direct connect capability with Ortho® Dial 'N Spray®
- Quick and easy to attach to Ortho® Dial 'N Spray®
- Easily connects to the Ortho® Dial 'N Spray®
- No measuring! Attaches directly to the Ortho® Dial 'N Spray®
- With a quick twist, attach directly to the Ortho® Dial 'N Spray®

For ORTHO Weed-B-Gon MAX Plus Crabgrass Control [Killer] Singles:

- NET 5 FL.OZ.
- Pre-Measured Concentrate Bottles
- Treats 5,000 sq.ft.
- No Measuring, No Mess
- Weed-B-Gon® Max(TM) Singles makes using concentrates easier than ever. No more measuring and no more mess. Mix (contents) (each pre-measured bottle) with two gallons of water to deliver consistent results. And its' revolutionary formula won't harm your lawn - Guaranteed.
- Designed for tank sprayers
- 1 bottle makes 2 gallons

3. Trademark acknowledgements

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00896.20130627.notif-proposed-highlighted.doc
2. Reason for Issue: ad claims, ABN