

2217-896

12/20/2013

1/32



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

James L. Kunstman, Ph.D.
PBI/Gordon Corporation
1217 W. 12th Street,
P.O. Box 014090
Kansas City, Missouri 64101-0090

DEC 20 2013

Subject: Label amendment to add a sub-label and new marketing claims to the master label
Product Name: EH-1437 Herbicide
EPA Reg. No: 2217-896
Decision Number(s): 482985

Dear Dr. Kunstman:

The labeling referred to above, submitted in connection with registration in accordance with FIFRA section 3(C)(5), as amended, is acceptable, provided that you submit and/or cite all data required for reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions. If you have questions or concerns regarding this letter, please contact Beth Benbow at (703) 347-8072 or email at benbow.bethany@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn V. Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

EH-1437 HERBICIDE

EPA Reg. No. 2217-896

Four sublabels represent the entire or master label of EH-1437 HERBICIDE. Please refer to the appropriate section of the labeling as shown as follows:

Section
Label language common to all sublabels
Sublabel 1. Traditional layout
Sublabel 2. Concentrate
Sublabel 3. Ready Spray
Sublabel 4. Alternate to Traditional layout
Appendix
Document Control Information

ACCEPTED
DEC 20 2013
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 2217-896

Label Language Common To All Sublabels

EH-1437 HERBICIDE

EPA Reg. No. 2217-896

ACCEPTED
DEC 20 2013
Under the Federal Insecticide, Fungicide, and Rodenticide Act, it is hereby ordered, for the pesticide labeled under EPA Reg. No. 2217-896

Alternate Brand Names:

- Tractor Supply Co. GroundWork Concentrate Crabgrass & Broadleaf Weed Killer
- Gordon's Trimec Crabgrass Plus Lawn Weed Killer Ready Spray
- Gordon's Trimec Crabgrass Plus Lawn Weed Killer Concentrate
- Trimec Crabgrass Plus Lawn Weed Killer Ready Spray
- Trimec Crabgrass Plus Lawn Weed Killer Concentrate

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	6.42%
Quinclorac	2.13%
Dicamba, dimethylamine salt	0.60%
OTHER INGREDIENTS:	90.85%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 0.456 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
 - 0.182 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
 - 0.043 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
- Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately

before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

DIRECTIONS FOR USE

[Note to Reviewer: The following section is for nonrefillable plastic containers and residential use only.]

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Alternate Warranty:

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for lack of performance or safety if not used according to the label directions.

[OPTIONAL WARRANTY STATEMENT]

The ORTHO Guarantee

If for any reason you [, the consumer,] are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.


6/32

SUBLABEL 1 Traditional layout

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
-  Optional Icon] Do not allow people or pets to enter the treated area until sprays have dried.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

SHAKE WELL BEFORE USING

PRODUCT FACTS:

What it Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)

When to Use: Apply when daytime temperatures are below 90°F.

Questions & Comments: (icon phone) (icon computer)
Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com

HOW IT WORKS:

EH-1437 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE:

EH-1437 Herbicide can be applied to residential lawns with the following turfgrass species:

Lawns To Be Treated:	
Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	Bermudagrass ¹ , common and hybrid Zoysiagrass Buffalograss
¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.	

DO NOT apply EH-1437 Herbicide to the following:

- Bahiagrass
- Bentgrass, colonial and creeping
- Carpetgrass
- Centipedegrass

Dichondra
 Gardens including vegetables, fruit trees, vines and berries
 Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs)
 Seashore paspalum
 St. Augustinegrass
 Turfgrass species that are not listed on this label.

GRASS AND BROADLEAF WEEDS CONTROLLED:

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Nettle	Spotted catsear
Bedstraw	Cudweed	Orange hawkweed	Spotted spurge
Beggarticks	Curly dock	Oriental cocklebur	Spurweed
Betony, Florida	Daisy, English	Oxalis	Stinging nettle
Bindweed, field	Daisy fleabane	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy, oxeye	Parsnip	Tall nettle
Bitter wintercress	Dandelion	Pearlwort	Tall vervain
Bittercress, hairy	Dichondra	Pennycress	Tansy ragwort
Bitterweed	Dogbane	Pennywort	Tansy mustard
Black-eyed Susan	Dogfennel	Peppergrass	Tanweed
Black medic	Dollarweed	Pepperweed	Thistle
Black mustard	False dandelion	Pigweed	Trailing crownvetch
Blackseed plantain	Falseflax	Pineywoods bedstraw	Tumble mustard
Blessed thistle	False sunflower	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed	Fiddleneck	Plantain	Velvetleaf
Blue lettuce	Florida pusley	Poison ivy	Venice mallow
Blue vervain	Frenchweed	Poison oak	Virginia buttonweed
Bracted plantain	Galinsoga	Pokeweed	Virginia creeper
Brassbuttons	Goathead	Poorjoe	Virginia pepperweed
Bristly oxtongue	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf dock	Ground ivy	Prickly lettuce	Western clematis
Broadleaf plantain	Gumweed	Prickly sida	Western salsify
Broomweed	Hairy fleabane	Prostrate knotweed	White mustard
Buckhorn	Hawkweed	Prostrate pigweed	Wild aster
Buckhorn plantain	Healall	Prostrate spurge	Wild buckwheat
Bulbous buttercup	Heartleaf drymary	Prostrate vervain	Wild carrot
Bull thistle	Heathaster	Puncturevine	Wild four-o'clock
Bullnettle	Hedge bindweed	Purslane, common	Wild garlic
Burclover	Hedge mustard	Ragweed	Wild geranium
Burdock	Hemp	Red sorrel	Wild lettuce
Burning nettle	Henbit	Redroot pigweed	Wild marigold
Burweed	Hoary cress	Redstem filaree	Wild mustard
Buttercup	Hoary plantain	Rough cinquefoil	Wild onion
Buttonweed	Hoary vervain	Rough fleabane	Wild parsnip
Canada thistle	Horsenettle	Russian pigweed	Wild radish
Carolina geranium	Jimsonweed	Russian thistle	Wild rape
Carpetweed	Knawel	Scarlet pimpernel	Wild strawberry
Catchweed bedstraw	Knotweed	Scotch thistle	Wild sweet potato
Catnip	Kochia	Sheep sorrel	Wild vetch
Catsear	Lambsquarters	Shepherd's purse	Wild violet
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil			Woolly plantain

BROADLEAF WEEDS			
Clover, crimson	Mexicanweed	Smooth pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Morningglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, sweet	Mugwort	Speedwell	and other broadleaf
Clover, white	Musk thistle	Spiny amaranth	weeds
Cockle	Mustard		

WHEN TO USE:

Apply this product when weeds are small and actively growing.

Weed species	Season	Timing	Maximum number of applications ¹	Application interval, days	Comments
Crabgrass (large and smooth):	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height	2	21	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	21	Re-treat at the same rate at 21 days later for dense populations.
Broadleaf signalgrass	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	21	Re-treat at the same rate at 21 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

¹The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. The maximum number of broadcast applications is limited to 2 per year. A second application may be made 21 days after the initial application. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre]

HOW MUCH TO USE:

I. ENTIRE LAWN TREATMENTS WITH DIAL TYPE HOSE-END SPRAYERS

Three Easy Steps To Mix And Apply:

1. FILL

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- The application rate for this product is 6.4 fl.oz. of product per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 6.4 fl.oz. of product for every 1,000 sq.ft. of lawn area. Over-application or rates above those listed on this label can cause turf injury.
- Do not add water to the sprayer jar.

2. SET THE DIAL

For hose-end sprayers: Set the dial to the rate per gallon and keep the dial set to this setting while spraying.

The dial setting or rate per gallon for this product is 2.5 oz. of product per gallon of water.

9/32

Types of hose-end sprayers	Dial setting
Ortho® Dial'N Spray® Hose-End Sprayer	2.5 oz.
Hudson® Hose End Liquid Concentrate Sprayer	2.5 fl.oz.
Gilmour® Dial-A-Mix® Lawn and Garden Sprayer	15 Teaspoons or 5 tablespoons

3. **SPRAY** - Apply evenly over the lawn.

II. ENTIRE LAWN TREATMENTS WITH READY-TO-USE HOSE-END SPRAYERS :

This product may be packaged as a ready-to-use product with no mixing. The retail plastic container is fitted with a sprayer body that delivers spray droplets at a fixed rate.

For READY-TO-USE Hose End Sprayers:

- This retail sprayer bottle will cover the treated areas described below when this product is applied at 6.4 fl.oz. of product per 1,000 sq.ft. Over-application or rates above those listed on this label can cause turf injury.

Approximate treated areas for four retail container sizes:

Species	Amount of product to treat approximate area, sq. ft.				
	Fl.oz./1,000 sq.ft.	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container	2.0 qt. (64 fl.oz.) retail container	1 gallon (128 fl.oz.) retail container
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season species Bermudagrass ¹ , common and hybrid Zoysiagrass and Buffalograss	6.4 fl.oz. /1,000 sq.ft.	2,500 sq.ft.	5,000 sq.ft.	10,000 sq.ft.	20,000 sq.ft.

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Instructions:

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.

Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl.oz. container will treat up to xxx sq.ft. of turfgrass.

- **FINISH:** To stop spraying, turn [rotate] spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn [rotate] spray nozzle to the "ON" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

III. HAND-OPERATED SPRAYERS:

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up):

- Apply to actively growing grass and broadleaf weeds during the early growth stages.
- Measure and mix the proper amount of product with water as described below. Mix 2.5 fl.oz. of product with 1 gallon of water to treat 400 sq.ft. of lawn. See below for additional spray preparations.
- Spray evenly over the lawn. Spray coverage should be uniform and complete. Over-application or rates above those recommended on this label can cause turf injury.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square feet
2.5 fl.oz.	1 gallon	400 sq.ft.
3.75 fl.oz.	1.5 gallons	600 sq.ft.
5.0 fl.oz.	2 gallons	800 sq.ft.
7.5 fl.oz.	3 gallons	1,200 sq.ft.

Equal measures: 2.5 fl.oz. = 5 Tablespoons. 1 cup = 8 fl.oz. = 1/2 pint.

OR

Measure indicated amount of product and pour into tank. Dilute with water according to the following chart. Adjust the nozzle to deliver coarse spray droplets and treat the area indicated.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Area To Treat, Square feet
2.5 fl.oz.	1 gallon	400 sq.ft.
5 fl. fl.oz.	2 gallons	800 sq.ft.
7.5 fl.oz.	3 gallons	1,200 sq.ft.

Equal measures: 2.5 fl.oz. = 5 Tablespoons. 1 cup = 8 fl.oz. = 1/2 pint.

**IV. ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® SYSTEM:
HOW TO USE THE READY-SPRAY® SYSTEM**

Connect:

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.
4. Extend hose to farthest area of the lawn to be treated

Spray:

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small plastic tab back and turn knob clockwise to ON position.
3. Spray evenly over measured area.

Finish:

1. To STOP spraying, rotate the knob counterclockwise to the OFF position.
2. Turn off water.
3. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.
4. Disconnect sprayer from hose.

HOW TO USE THE ORTHO® DIAL-N-SPRAY® APPLICATOR WITH DIRECT CONNECT [Applicator Name]

[Prepare Applicator:] [illustration]

1. Set dial to [X] fl.oz./Tbsp./tsp.

[Connect:] [illustration]

1. Remove cap from the concentrate bottle. [Keep cap for storage.]
2. Insert suction tube into bottle opening and twist the Ortho® Dial 'N Spray® applicator [applicator name] onto bottle to attach.
3. Connect hose to the sprayer and turn on water.

[Spray:] [illustration] [directions will vary by product type]

1. Hold sprayer in upright position and apply evenly to area to be treated.
2. Spray only until the surface of the lawn is wet.

[Finish:] [illustration]

1. After use, untwist to remove the [insert product name] concentrate bottle from the Ortho® Dial 'N Spray® applicator [applicator name].
2. Replace cap on the [insert product name] concentrate bottle.

Refer to the Ortho® Dial 'N Spray applicator [applicator name] booklet for additional information. Important: To prevent sprayer from clogging, always clean after use as directed.

V. FOR TANK SPRAYER:

- Add 2.5 fl.oz. per gallon of water for each 400 sq.ft.

For broadcast application: Ortho Dial 'N Spray® Applicator

1. Add 16 fl.oz. of concentrate directly into sprayer jar. DO NOT ADD WATER.
2. Set dial to 2.5 oz.
3. Spray evenly over 2,500 sq.ft.
4. Any unused portion should be poured back into its original container.

[Over-application or rates above those recommended on this label can cause turf injury]

VI. FOR ORTHO WEED-B-GON MAX PLUS CRABGRASS CONTROL [KILLER] SINGLES:

For (Use a) Tank Sprayer:

- Add entire contents (of 1 bottle) (5 oz.) to an empty 2-gallon tank sprayer. Slowly fill with 2 gallons of clean tap water.
- **For Broadcast Application – Ortho® Dial 'N Spray® Applicator**
 1. Set dial to 2.5 oz.
 2. Add entire contents of 1 bottle (5 oz) directly into sprayer jar. DO NOT ADD WATER.
 3. Spray evenly over 800 sq.ft.

[Over-application or rates above those recommended on this label can cause turf injury]

VII. ENTIRE LAWN TREATMENTS WITH THE ORTHO READY-SPRAY® II [2]/APPLICATOR NAME SYSTEM:

HOW TO USE THE READY-SPRAY® II/[READY-SPRAY 2]/[APPLICATOR NAME] SYSTEM

[Optional]

1. [Measure the total [entire] area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.]

[End of optional section]

Connect:

1. Shake well before using. [optional step]
2. Connect sprayer to hose.
3. Turn on water.
4. Extend hose to farthest area of the measured area to be treated. [optional step]

Spray:

1. To begin spraying, [disengage safety lock and] point nozzle in the direction you want to spray.
2. Slide thumb activation switch forward to ON position.
3. Spray evenly over measured area.

Finish:

1. To STOP spraying, slide thumb activation switch backward to OFF position.
2. Turn off water.

3. Relieve water pressure by sliding thumb switch to ON /WATER/ WATER ONLY position until water slows to a drip. Then, slide switch back to OFF position [and engage safety lock].
4. Disconnect sprayer from hose.

[Optional illustrations may be used in conjunction with the above steps]

CULTURAL TIPS AND NOTES:		
When to Apply:	<p>For established lawns</p> <p>For newly seeded areas</p> <p>For newly sodded, sprigged, or plugged areas</p>	<ol style="list-style-type: none"> 1. Apply when daytime temperatures are below 90°F. 2. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing. 3. Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
Seeding:		<ul style="list-style-type: none"> • The treated area can be seeded at 4 weeks after the application of this product.
Irrigation:	<p>Non-stressed turfgrass</p> <p>Stress from high temperatures, drought, insects or diseases.</p>	<ul style="list-style-type: none"> • Adequate soil moisture and favorable growing conditions enhance the performance of this product. • Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing:		<ul style="list-style-type: none"> • Delay mowing 2 days before and until 2 days after the application of this product.

13/32

SUBLABEL 2 Concentrate

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Spot Treatments: Mix 2.5 fl.oz. (5 Tbs) in 1 gallon of water to treat 400 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Over-application or rates above those specified on this label can cause turf injury.

3. When To Use

Apply this product when weeds are small and actively growing.

Weed species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 21 days later for dense populations.
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 21 days later if needed.

Weed species	Timing	Comments
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture
The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre]		

4. How To Apply

4.1 Pump Style Sprayers [sprayer icon]

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

4.2 [12 Volt &] Tow-Behind Boom Sprayers [sprayer icon]

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq.ft.. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Tank Size	Spray volume	Coverage Area	Concentrate to add to tank
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft.	192 fl.oz. (1.5 gal)
	1 gal/1000 sq.ft.	15,000 sq.ft.	96 fl.oz. (0.75 gal)
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft.	320 fl.oz. (2.5 gal)
	1 gal/1000 sq.ft.	25,000 sq.ft.	160 fl.oz. (1.25 gal)

For spot treatments with [12 Volt &] Pull Behind Sprayers
See rates under "How Much To Apply," Spot Treatments.

4.3 Dial Style Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER.**
2. Close sprayer and set dial to 2.5 oz.
3. Connect sprayer to hose

15/34

4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

4.4 Concentrate Plus Water Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar.
3. Close sprayer, connect to garden hose, shake well.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

5. Use Precautions:

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected

6. Grass and Broadleaf Weeds Killed

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Nettle	Spotted catsear
Bedstraw	Cudweed	Orange hawkweed	Spotted spurge
Beggarticks	Curly dock	Oriental cocklebur	Spurweed
Betony, Florida	Daisy, English	Oxalis	Stinging nettle
Bindweed, field	Daisy fleabane	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy, oxeye	Parsnip	Tall nettle
Bitter wintercress	Dandelion	Pearlwort	Tall vervain
Bittercress, hairy	Dichondra	Pennycress	Tansy ragwort
Bitterweed	Dogbane	Pennywort	Tansy mustard
Black-eyed Susan	Dogfennel	Peppergrass	Tanweed
Black medic	Dollarweed	Pepperweed	Thistle
Black mustard	False dandelion	Pigweed	Trailing crownvetch
Blackseed plantain	Falseflax	Pineywoods bedstraw	Tumble mustard
Blessed thistle	False sunflower	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed	Fiddleneck	Plantain	Velvetleaf
Blue lettuce	Florida pusley	Poison ivy	Venice mallow
Blue vervain			Virginia buttonweed

BROADLEAF WEEDS			
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Wavyleaf bullthistle
Broadleaf dock	Goldenrod	Prairie sunflower	Western clematis
Broadleaf plantain	Ground ivy	Prickly lettuce	Western salsify
Broomweed	Gumweed	Prickly sida	White mustard
Buckhorn	Hairy fleabane	Prostrate knotweed	Wild aster
Buckhorn plantain	Hawkweed	Prostrate pigweed	Wild buckwheat
Bulbous buttercup	Healall	Prostrate spurge	Wild carrot
Bull thistle	Heartleaf drymary	Prostrate vervain	Wild four-o'clock
Bullnettle	Heathaster	Puncturevine	Wild garlic
Burclover	Hedge bindweed	Purslane, common	Wild geranium
Burdock	Hedge mustard	Ragweed	Wild lettuce
Burning nettle	Hemp	Red sorrel	Wild marigold
Burweed	Henbit	Redroot pigweed	Wild mustard
Buttercup	Hoary cress	Redstem filaree	Wild onion
Buttonweed	Hoary plantain	Rough cinquefoil	Wild parsnip
Canada thistle	Hoary vervain	Rough fleabane	Wild radish
Carolina geranium	Horsenettle	Russian pigweed	Wild rape
Carpetweed	Jimsonweed	Russian thistle	Wild strawberry
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild sweet potato
Catnip	Knotweed	Scotch thistle	Wild vetch
Catsear	Kochia	Sheep sorrel	Wild violet
Chickweed, common	Lambsquarters	Shepherd's purse	Woodsorrel
Chickweed, mouseear	Lespedeza	Slender plantain	Woolly croton
Chicory	Mallow	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Matchweed	Smooth dock	Woolly plantain
Clover, crimson	Mexicanweed	Smooth pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Morningglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, sweet	Mugwort	Speedwell	and other broadleaf
Clover, white	Musk thistle	Spiny amaranth	weeds
Cockle	Mustard		

17/32

[Back Panel] [Restatement of information in sections 1 and 2.]

Getting Started

Kills:

Crabgrass, foxtail and most troublesome broadleaf weeds including dandelion, clover, thistle and over XXX other weeds listed.

Where To Use:

On established lawns without harming lawn grasses when used as directed (Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.) [See Inside Label for Complete Directions]

Amount To Use:

Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:

Spot Treatments: Mix 2.5 fl.oz. (5 Tbs) in 1 gallon of water to treat 400 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft. [See Inside Label for Sprayer Settings]
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Coverage:

This [size] bottle treats XXXX sq.ft.

Telephone Icon

Computer Icon

For questions or comments call toll-free 1-800-821-7925 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST or visit [our website at] www.pbigordon.com [GordonsUSA.com]

SUBLABEL 3 Ready Spray

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Entire Lawn Treatments: This container covers XXX sq.ft. Measure the size of your yard. If the area to be treated is larger than XXX sq.ft., you may need to buy more containers.

Over-application or rates above those specified on this label can cause turf injury.

3. When To Use

Apply this product when weeds are small and actively growing.

Weed species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 21 days later for dense populations.
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 21 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture

19/34

Weed species	Timing	Comments
The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre]		

4. How To Apply

1. Connect your garden hose to the hose-end sprayer. With your thumb, snap off the safety locking tab that prevents accidental discharge of spray solution. Leave control valve in the OFF position.
2. To BEGIN spraying: Turn on water at faucet. Extend hose to the furthest point from the faucet. Point nozzle toward the lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Walk backwards as you apply in order to avoid contact with the treated area. For best results, spray evenly over the measured area.
3. To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

5. Use Precautions:

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- Certain hard-to-kill weeds may need to be spot treated in 14 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

6. Grass and Broadleaf Weeds Killed

A partial list is presented below:

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Amaranth, purple	Cupid shaving Brush	Mustard, blue	Speedwell, corn
Amaranth, slender	Cudweed	Mustard, hedge	Speedwell, ivyleaf
Aster, heath	Daisy, English	Mustard, Indian,	Speedwell, persian
Aster, white heath	Daisy, oxeye	Mustard, tansy	Speedwell, purslane
Aster, purple	Dandelion, common	Mustard, tumble	Speedwell, slender
Aster, white prairie	Dandelion, false	Mustard, white	Speedwell, snow
Artichoke	Day flower	Mustard, wormseed	Speedwell, thymeleaf
Austrian fieldcress	Deadnettle	Nettle, stinging (burning)	Spiny amaranth
Bedstraw, Catchweed	Dichondra	Nettle, tall	Spurge, spotted
Bedstraw, smooth	Dock, broadleaf	Nettle, wood	Spurge, prostrate
Bedstraw, Northern	Dock, curly	Parsley-piert	Spurweed
Bedstraw, pineywoods	Dogbane	Parsnip	Star of Bethlehem
Beggarticks, hairy	Dogfennel	Pearlwort	Strawberry, India mock
Betony, Florida	Dollarweed (Pennywort)	Pennycress (frenchweed)	Tansy ragwort
Bindweed, Field	Dove weed	Pepperweed, perennial	Tanweed

BROADLEAF WEEDS			
Bindweed, Hedge	Falseflax	Pepperweed, virginia (Peppergrass)	Thistle, blessed
Bitter Wintercress	False sunflower	Pigweed, prostate	Thistle, bull
Bittercress, hairy	Fiddleneck	Pigweed, redroot	Thistle, canada
Bitterweed	Filaree, redstem	Pigweed, russian	Thistle, flodman (prairie)
Black-eyed Susan	Filaree, whitestem	Pigweed, smooth	Thistle, musk
Black Medic	Fleabane, daisy (annual)	Pigweed, tumble	Thistle, platte
Blood flower milkweed	Fleabane, rough	Pineapple weed	Thistle, plumeless
Blue lettuce	Galinsoga, hairy	Plains coreopsis (tickseed)	Thistle, russian
Brassbuttons	Galinsoga, smallflower	Plantain, blackseed	Thistle, scotch
Bristly oxtongue	Goldenrod	Plantain, bracted	Thistle, wavyleaf
Broomweed	Ground ivy	Plantain, broadleaf	Thistle, yellow (horrible)
Buckhorn	Groundsel	Plantain, buckhorn	Trailing crownvetch
Bullnettle	Gumweed	Plantain, hoary	Velvetleaf
Burclover	Hairy fleabane	Plantain, narrowleaf	Vervain, blue
Burdock	Hawkweed, meadow	Plantain, slender	Vervain, hoary
Burweed, lawn (spurweed)	Hawkweed, mouseear	Plantain, woolly	Vervain, prostrate
Buttercup, bulbous	Hawkweed, orange	Poison ivy	Vervain, tall
Buttercup, creeping	Hawkweed, yellow	Poison oak	Vetch, bird
Buttonweed, common (poorjoe)	Healall	Pokeweed	Vetch, hairy
Buttonweed, virginia	Heartleaf drymary	Prairie Sunflower	Vetch, milk
Carolina geranium (wild)	Hemp	Prickly lettuce (compass plant)	Vetch, narrowleaf
Carpetweed	Henbit	Prickly sida	Vetch, wild
Catnip	Hoary cress	Prostrate knotweed	Virginia creeper
Catsear, spotted (false dandelion)	Horsenettle	Puncturevine	Western clematis
Chamber bitter	Horseweed	Purslane, common	Western salsify
Chickweed, common	Jimsonweed	Pusley, brazil	Wild aster
Chickweed, mouseear	Johnny-Jumpup violet	Pusley, florida	Wild buckweat
Chickweed, sticky	Knawel	Pusley, large flower	Wild carrot
Chicory	Knotweed, prostrate	Ragweed, bur	Wild four-o'clock
Cinquefoil	Kochia	Ragweed, common	Wild garlic
Clover, crimson	Lambsquarters	Ragweed, lancleaf	Wild lettuce
Clover, alyce	Lespedeza	Ragweed, western	Wild marigold
Clover, hop	Mallow, alkali	Redstem filaree	Wild mustard (yellow rocket)
Clover, rabbit foot	Mallow, bristly	Rough cinquefoil	Wild onion
Clover, red	Mallow, common	Scarlet pimpernel	Wild parsnip
Clover, strawberry	Mallow, dwarf	Shepherd's purse	Wild radish
Clover, yellow sweet	Mallow, venice	Smartweed, ladythumb	Wild rape
Clover, white sweet	Marcela	Smartweed, pale	Wild strawberry
Clover, white	Matchweed	Smartweed, pennsylvania	Wild sweet potato
Cockle	Mexicanweed	Smooth chaff-flower	Wild violet
Cocklebur	Moneywort	Smoothdock	Woodsorrel, creeping
Cocklebur, oriental	Morningglory, bigroot	Sorrel, red (sheep)	Woodsorrel, violet
Cocklebur, spiny	Morningglory, ivyleaf	Sowthistle, annual	Woodsorrel, yellow (Oxalis)
Common mullein	Morningglory, mexican	Sowthistle, spiny	Woolly croton
Corn Spurry	Morningglory, red	Spanish needles	Wormseed
Creeping Beggarweed	Morningglory, tall	Speedwell, bi-lobed	Yarrow, common
Creeping Jenny (creeping charlie)	Morningglory, wolly	Speedwell, common	Yarrow, western
	Mugwort		Yarrow, yellow
	Mustard, ball		
	Mustard, black		

21/34

[Back Panel] [Restatement of information in sections 1 and 2.]

Getting Started

Kills:

Crabgrass, foxtail and most troublesome broadleaf weeds including dandelion, clover, thistle and over XXX other weeds listed.

Where To Use:

On established lawns without harming lawn grasses when used as directed (Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.) [See Inside Label for Complete Directions]

Amount To Use:

Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:

Entire Lawn Treatments: This container covers XXX sq.ft.

Coverage:

This [size] bottle treats XXXX sq.ft.

Telephone Icon

Computer Icon

For questions or comments call toll-free 1-800-821-7925 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST or visit [our website at] www.pbigordon.com [GordonsUSA.com]

SUBLABEL 4
Alternate to Traditional Layout

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USE.

SHAKE WELL BEFORE USING

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- [Reentry Icon] Do not allow people or pets to enter the treated area until sprays have dried.
- Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not spray exposed roots of ornamentals and trees.

USE PRECAUTIONS:

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- Certain hard-to-kill weeds may need to be spot treated in 14 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Avoid applications to bermudagrass when daytime temperatures exceed 85°F.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected.
- Use on established lawns.

HOW IT WORKS/PRODUCT INFORMATION:

EH-1437 Herbicide/Product Name contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE:

EH-1437 Herbicide/Product Name/This product can be applied to residential lawns with the following turfgrass types:

Lawns To Be Treated:	
Cool Season Turfgrass (Northern Lawns)	Warm Season Turfgrass (Southern Lawns)

Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Fine fescues	Bermudagrass ¹ , common and hybrid Zoysiagrass Buffalograss
¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.	

DO NOT apply EH-1437 Herbicide to the following:
Bahiagrass Bentgrass Carpetgrass Centipedegrass Dichondra Gardens including vegetables, fruit trees, vines and berries Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs) Seashore paspalum St. Augustinegrass Turfgrass species that are not listed on this label.

HOW MUCH TO USE:

Use 2.5 fl.oz. per gallon of water for every 400 sq.ft.

Limitations: The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.0 lb 2,4-D acid equivalent per acre per application]. Do not apply more than 2 applications per year. The second application may be made no sooner than 21 days after the first application. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.0 lb 2,4-D acid equivalent per acre].

WHEN TO USE:

Apply this product when weeds are small and actively growing in the spring or fall.

Crabgrass (large & smooth) – Apply in the spring before crabgrass is 4 inches tall or when crabgrass is first noticeable in the lawn. Make a second application after 3 weeks for dense weed populations.

GRASS AND BROADLEAF WEEDS CONTROLLED [KILLS WEEDS INCLUDING:]

[Some or all of the following weeds will appear on the product label]

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping Charlie)	Nettle	Spotted catsear
Bedstraw	Cudweed	Orange hawkweed	Spotted spurge
Beggarticks	Curly dock	Oriental cocklebur	Spurweed
Betony, Florida	Daisy, English	Oxalis	Stinging nettle
Bindweed, field	Daisy fleabane	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy, oxeye	Parsnip	Tall nettle
Bitter wintercress	Dandelion	Pearlwort	Tall vervain
Bittercress, hairy	Dichondra	Pennycress	Tansy ragwort
Bitterweed	Dogbane	Pennywort	Tansy mustard
Black-eyed Susan	Dogfennel	Peppergrass	Tanweed
Black medic	Dollarweed	Pepperweed	Thistle
Black mustard	False dandelion	Pigweed	Trailing crownvetch
Blackseed plantain	Falseflax	Pineywoods bedstraw	Tumble mustard
Blessed thistle	False sunflower	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed			Velvetleaf

BROADLEAF WEEDS			
Blue lettuce	Fiddleneck	Plantain	Venice mallow
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Wavyleaf bullthistle
Broadleaf dock	Goldenrod	Prairie sunflower	Western clematis
Broadleaf plantain	Ground ivy	Prickly lettuce	Western salsify
Broomweed	Gumweed	Prickly sida	White mustard
Buckhorn	Hairy fleabane	Prostrate knotweed	Wild aster
Buckhorn plantain	Hawkweed	Prostrate pigweed	Wild buckwheat
Bulbous buttercup	Healall	Prostrate spurge	Wild carrot
Bull thistle	Heartleaf drymary	Prostrate vervain	Wild four-o'clock
Bullnettle	Heathaster	Puncturevine	Wild garlic
Burclover	Hedge bindweed	Purslane, common	Wild geranium
Burdock	Hedge mustard	Ragweed	Wild lettuce
Burning nettle	Hemp	Red sorrel	Wild marigold
Burweed	Henbit	Redroot pigweed	Wild mustard
Buttercup	Hoary cress	Redstem filaree	Wild onion
Buttonweed	Hoary plantain	Rough cinquefoil	Wild parsnip
Canada thistle	Hoary vervain	Rough fleabane	Wild radish
Carolina geranium	Horsenettle	Russian pigweed	Wild rape
Carpetweed	Jimsonweed	Russian thistle	Wild strawberry
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild sweet potato
Catnip	Knotweed	Scotch thistle	Wild vetch
Catsear	Kochia	Sheep sorrel	Wild violet
Chickweed, common	Lambsquarters	Shepherd's purse	Woodsorrel
Chickweed, mouseear	Lespedeza	Slender plantain	Woolly croton
Chicory	Mallow	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Matchweed	Smooth dock	Woolly plantain
Clover, crimson	Mexicanweed	Smooth pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Morningglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed and other broadleaf weeds
Clover, sweet	Mugwort	Speedwell	
Clover, white	Musk thistle	Spiny amaranth	
Cockle	Mustard		

[Directions For Use Options Based On Applicator]

[Option 1: Dial 'N Spray/Dial 'N Spray with Direct Connect]

[Applicator] [Device] [Applicator Name] Directions/Instructions

HOW TO USE:

Set:

- Turn dial to 2.5 oz. [For Northern/Southern Lawns]

Fill:

- Pour desired amount of concentrate into sprayer jar.
- Turn sprayer nozzle to preferred spray.
- Do NOT add water.

Spray:

- Connect [garden] hose to sprayer.
- Turn on water.
- Press and hold trigger for continuous spray. Apply evenly.

Finish:

- Pour unused concentrate back into its original container.
- To clean: refer to the Ortho® Dial N Spray® cleaning instructions.

[Optional illustrations may be used in conjunction with the above steps]

AND/OR

[Applicator] [Device] [Applicator Name] Directions/Instructions

HOW TO USE:

Set: [Applicator Name]

- 1. Turn dial to 2.5 oz.

Connect:

- 1. Remove jar from [Ortho® Dial 'N Spray®] [applicator name] [applicator/sprayer].
- 2. Remove cap from the concentrate bottle. [Keep cap for storage.]
- 3. Twist [screw] the [Ortho® Dial 'N Spray®] [applicator name] [applicator/sprayer] onto concentrate bottle to attach.
- 4. Connect [garden] hose to the sprayer and turn on water.

Spray:

- 1. Press and hold trigger for continuous spray. Apply evenly.
- 2. Spray only until the surface [of the lawn] is wet.

Finish:

- 1. Untwist [unscrew] the [product name] concentrate bottle from the [Ortho® Dial 'N Spray®] [applicator name] [applicator/sprayer].
- 2. Replace cap on the [product name] concentrate bottle.
- 3. To clean: Refer to the Ortho® Dial 'N Spray® cleaning instructions.

[Optional illustrations may be used in conjunction with the above steps]

Refer to the Ortho® Dial 'N Spray label and booklet for additional information.

Important: Over-application or rates above those recommended on this label can cause grass [turf/lawn] injury. To prevent sprayer from clogging, always clean after use, as directed.

[Option 2: Tank Sprayer]

[Applicator] [Device] [Applicator Name] Directions/Instructions

HOW TO USE:

Add 2.5 fl.oz. per gallon of water for every 400 sq.ft.

Important:

Over-application or rates above those recommended on this label can cause grass [turf/lawn] injury.

[Option 3: Ortho® Ready-Spray® II System]

[Applicator] [Device] [Applicator Name] Directions/Instructions

HOW TO USE:

Connect:

- 1. Connect sprayer to [garden] hose.
 - 2. Turn on water.
- [User Tip: Start at the farthest point in your yard and work back.]

Spray:

- 1. To BEGIN spraying, slide/rotate [color] switch to "ON".
- 2. Spray only until the surface of the lawn is wet. Spray evenly.

Finish:

1. To STOP spraying, slide/rotate [color] switch to "OFF".
2. Turn off water.
3. Relieve water pressure by sliding/rotating [color] switch to "WATER/ON".
4. After water slows to a drip, slide/rotate [color] switch back to "OFF".
5. Disconnect sprayer from [garden] hose.

[Optional illustrations may be used in conjunction with the above steps]

USE TIPS:		
When to Apply:	For established lawns	Apply when daytime temperatures are below 90°F.
	For newly seeded areas	Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
	For newly sodded, sprigged, or plugged areas	Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
Seeding:		The treated area can be seeded at 4 weeks after the application of this product.
Irrigation:	Non-stressed turfgrass	Adequate soil moisture and favorable growing conditions enhance the performance of this product.
	Stress from high temperatures, drought, insects or diseases.	Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing:		Delay mowing 2 days before and until 2 days after the application of this product.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on the retail containers of supplemental registrants.

Efficacy Claims

- Controls crabgrass, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills Crabgrass and over XXX Listed Weeds in Country & Suburban Lawns* *See inside booklet for a complete list of weeds controlled
- Kills [controls] [Crabgrass,] [Dandelion,] [Clover,] and other common [lawn weeds] [weeds found in your lawn] (See inside booklet for a complete list of weeds controlled)
- Controls crabgrass and broadleaf weeds in lawns (See inside booklet for a complete list of weeds controlled)
- Kill crabgrass & [any listed weed(s)] too!
- Kills Crabgrass [plus [over XXX] Broadleaf Weeds] (See inside booklet for a complete list of weeds controlled)
- Kills both Crabgrass & Broadleaf weeds (See inside booklet for a complete list of weeds controlled)
- Kills lawn weeds plus crabgrass. (See inside booklet for a complete list of weeds controlled)
- Kills [Controls] 200+ broadleaf weeds. (See inside booklet for a complete list of weeds controlled)
- Controls (Number) weeds! (See inside booklet for a complete list of weeds controlled)
- **CONTROLS**
 - **SENSITIVE WEEDS:** Crabgrass, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.
 - **HARD-TO-KILL WEEDS:** Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and other broadleaf weeds. (See inside booklet for a complete list of weeds controlled)
- Kills [Lawn] Weeds [plus] [including] Crabgrass in One [Easy] Step (See inside booklet for a complete list of weeds controlled)
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step (See inside booklet for a complete list of weeds controlled)
- Kills [Controls] Listed Weeds (See inside booklet for a complete list of weeds controlled)
- Kills Hard-to-Control Weeds (See inside booklet for a complete list of weeds controlled)
- Kills [Controls] Both Broadleaf and Grass Weeds [in lawns] (See inside booklet for a complete list of weeds controlled)
- [Also] Kills [Dandelion], [Crabgrass], [Clover], plus XXX Other [Listed] Weeds [As Listed] (See inside booklet for a complete list of weeds controlled)
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds (See inside booklet for a complete list of weeds controlled)
- Effectively Kills Both Broadleaf and Certain Weed Grasses (See inside booklet for a complete list of weeds controlled)
- This product kills Crabgrass PLUS broadleaf weeds. The special Trimec® dual-action formula goes beyond crabgrass and foxtail to kill over XXX other broadleaf weeds listed in country and suburban lawns. (See inside booklet for a complete list of weeds controlled)
- Lawn Weed & Crabgrass Killer (See inside booklet for a complete list of weeds controlled)
- Kills Dandelions, Clover, and Other Broadleaf Weeds! Plus Kills Crabgrass and Other grass Weeds! (See inside booklet for a complete list of weeds controlled)
- Kills the Toughest Weeds in these Lawns (any lawn listed on label) (See inside booklet for a complete list of weeds controlled)
- Kills Over 200 Lawn Weeds plus [including] Crabgrass [in one easy step]. (See inside booklet for a complete list of weeds controlled)
- Kills Major [Lawn] Broadleaf Weeds (See inside booklet for a complete list of weeds controlled)
- Kills the weeds not the lawn when used as directed
- Kill [control] the most common weeds in your lawn [grass] (See inside booklet for a complete list of weeds controlled)

- Kills [controls] [hard-to-control] [tough] weeds (See inside booklet for a complete list of weeds controlled)
- Gentle on lawns [grass], tough on weeds!
- Tough weed killer
- Kills [controls] all types of [lawn] weeds [in your grass] (See inside booklet for a complete list of weeds controlled)
- Controls [Kills] even the tough weeds...roots and all!
- Kills the root.
- Kills Even Tough Weeds Down to the root
- Kills Crabgrass and Broadleaf weeds...-Down to the root
- Gets to the root [of your weed problem]
- Kills [lawn] weeds [down] to the root[s]
- Kills to the root
- Kills [controls] XX types of weeds [- broadleaf or grass weeds] (See inside booklet for a complete list of weeds controlled)
- Kill[s] [controls] [over] [up to] [XXX+] [lawn] [types of] weeds (See inside booklet for a complete list of weeds controlled)
- Won't harm lawn grass when used according to label directions.
- Won't harm [lawn] grass when used as directed.
- Won't harm lawns* *When used as directed
- Kills Weeds, Not Lawns(TM)
- [Even] kills [controls] poison ivy and poison oak
- [Only] targets the weeds, leaving your lawn [grass] unharmed when used as directed
- Targets weeds invading your lawn [grass]
- [Visible] [See] results in X hours
- Kills by contact
- Starts working on contact
- Results [works] with 1 [one] application
- Simple [and easy] steps to a weed free [lawn] [yard]
- One [1] [easy] solution for crabgrass, dandelions and clover

Formula Claims

- Keep from freezing.
- Water based.
- Mixes with Water - Concentrate
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Contains Trimec Herbicide
- 2-in-1
- All-in-One [complete] [total] [2-in-1] weed control [for lawns]
- All in One Weed Solution
- [With] Lawn Guard™ [Lawn Protect™]
- Colorless formula
- Non-staining
- Stain-free [formula]

Characteristics

- XX covers up to XXXX square feet.
- Makes more than XX Gallons!
- One quart covers up to 5,000 square feet.
- One [1] [bottle size] covers up to XXXX square feet
- 1 quart treats [up to] 5,000 sq.ft.
- Covers [treats up to] [XXXX] sq.ft.
- [XX] [Pint, Quart, etc] Treats Up To [YYYY] Sq.Ft.

- Bonus Size
- 25% More Free!
- Easily covers large areas
- Economical – use to cover large areas
- [Great] [Ideal] for broadcast treatment [application]
- Great value [- use to cover small or large areas]
- Designed for use on Southern lawns [warm season grasses/list approved grass types here]
- Treats many Northern & Southern grass types
- For use on [many Northern & Southern] grasses [lawns] including XXX
- XX% more than X gallon size
- The value of XX [size RTU] in one concentrate

Weather & Speed

- Starts working [overnight] [immediately]
- Rainfast in 1 Hour
- [Rainproof] [rainfast] in [1] [X] Hour[s]
- [See] results in hours
- Overnight results
- Fast-acting [formula]

Applicator Claims

- [Easy] [Ergonomic] Comfort Grip [Design] [Handle]
- Ergonomic [design]
- Easy grip [control]
- Easy [to use]
- Simple to use
- Convenient [Simple] [Design]
- Simple [and easy] [to use]
- [Simply] attach to [garden] hose and spray [go]
- Gentle shower [applicator] [spray pattern]
- Hand doesn't get wet
- Hand stays dry
- Hands stay dry
- No messy clean-up
- No concentrate contact with Ortho® Dial 'N Spray®
- Includes water [only] setting [option]
- Instant/Easy [One-handed] on/off applicator/switch/action
- One-handed operation [use] [spraying]
- Ready-Spray®/Ready-to-Spray/Applicator Name
- Ready-Spray
- Direct connect to the Ortho® Dial 'N Spray®
- Direct connect to Ortho® Dial 'N Spray® means no more measuring
- No more measuring concentrates with Ortho® Dial 'N Spray®
- [Requires] no measuring, [no mixing]
- No measuring! Attaches directly to [the] Ortho® Dial 'N Spray®
- With a quick twist, attach directly to the Ortho® Dial 'N Spray®
- Simple, direct connect capability with Ortho® Dial 'N Spray®
- Exact measurement and improved efficacy with Ortho® Dial 'N Spray®
- No measure, direct connect capability with Ortho® Dial 'N Spray®
- No over/under application with Ortho® Dial 'N Spray®
- No over/under application, direct connect capability with Ortho® Dial 'N Spray®
- Quick[ly] [and easily] to attach to Ortho® Dial 'N Spray®
- Direct[ly] [Easily] connects to [the] Ortho® Dial 'N Spray®

- No mixing required
- Just connect to garden hose & spray
- [Quick and] Easy [and simple] to use
- [The] Easy way to spray
- Easy [Simple] to attach [to [garden] hose]
- One step prep
- [Applicator name]
- XX [oz] Dial N Spray setting
- Save[s] time [and energy]
- Mess free cap [top] [lid] [cup]
- Measure with ease
- Easy-Measure [Cap] [Top] [Lid] [Cup]
- Drain-Back [Cap] [Top] [Lid] [Cup]
- Quick, convenient, and mess free
- [reusable] [drain-back/measuring] cap
- Quickly [and easily] attaches [connects] to Ortho® Dial N Spray®

Other

- * when used as directed
- Local Solutions for Local Problems
- For more Lawn and Garden Products Visit [www.website.com]
- Open Resealable Label for Directions & Precautions (Press to Reseal)
- Ortho Weed-B-Gon MAX Concentrate Plus Crabgrass Control [Killer]
- For [Use on] Lawns
- For home lawn care
- For Home Use [Only]
- Weed control for your lawn
- Weed control made easy
- For outdoor use
- Easy to mix
- Proven performance
- Systemic [Herbicide] [Activity]
- Before [illustration]/After [illustration]
- Be in control of weeds in your yard
- For a beautiful [weed-free] lawn
- [For] [Get] a weed free lawn
- Helps keep lawns weed free
- Weed free [worry free]
- From the lawn care experts at Scotts
- From the pest experts at Ortho
- Help[s] [keep[s]] the lawn look[ing] beautiful
- Lawn friendly [weed control] when used as directed
- Guaranteed [Results] *
- Guaranteed not to harm the lawn when used as directed
- Ortho guarantee
- NOTICE: To the extent consistent with applicable law, Buyer assumes all risks of use, storage, or handling of this product not in accordance with [label] directions
- ORTHO & Your Environment - Your yard is a place for your family and pets to enjoy. That's why ORTHO products are designed with care to provide effective solutions to weed problems. [For best results] please follow instructions for appropriate usage, storage and disposal.
- Won't harm [the] lawn, guaranteed*

- How it works: The [insert brand name] formula kills BOTH broadleaf weeds and Crabgrass in one easy step [without harming your lawn - guaranteed]. (See inside booklet for a complete list of weeds controlled)
- Ortho's #1 Weed Killer for Lawns
- Ortho's [Our] most powerful weed killer for lawns
- Ortho's simple weed killer
- Read Entire label before use
- Recyclable bottle
- [Recycle symbol]
- Takes the guesswork out of weed control [in the lawn]
- XX% less plastic . . . As compared to [alternate product]
- **Optional Instructions for Side Clip Tape**
 PULL / PEEL HERE [TO REMOVE TAPE]
 Open Resealable Label For Directions and/ & Precautions
 Press to Reseal
- **Questions & Comments:** [icon phone] [icon computer]
- Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com
- [Superscript is first letter of lot number]

For ORTHO Weed-B-Gon MAX Plus Crabgrass Control [Killer] Singles:

- NET 5 FL.OZ.
- Pre-Measured Concentrate Bottles
- Treats 5,000 sq.ft.
- No Measuring, No Mess
- Weed-B-Gon® Max(TM) Singles makes using concentrates easier than ever. No more measuring and no more mess. Mix (contents) (each pre-measured bottle) with two gallons of water to deliver consistent results. And its' revolutionary formula won't harm your lawn - Guaranteed.
- Designed for tank sprayers
- 1 bottle makes 2 gallons

[The following Product Facts format is optional and the information may appear inside or outside of a box.]

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)

When to Use: Apply when daytime temperatures are below 90°F. (do not apply above 85°F for bermudagrass).

Questions & Comments: [icon phone] [icon computer]
Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com

3. Trademark acknowledgements

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

32/32

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00896.20131004.amend-proposed-clean.doc
2. **Reason for Issue:** EPA Comments #1