

2217-903

4/22/2010

10/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

James Kunstman
PBI/Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101

APR 22 2010

Subject: Dicamba and MCPP-p RED Label Amendments
EH-1446 Herbicide
EPA Reg. No. 2217-903

The labeling referred to above is in compliance with the Dicamba and MCPP-p REDs and is amended under the REDs provided that you:

1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following revisions to the product label:
 - . Make the following revisions to the final printed labeling:
 - a. The "IF ON SKIN" first aid heading must be revised to read: "IF ON SKIN OR CLOTHING"
 - b. The acid equivalent amount of Dicamba and 2,4-D should be added to the text under the heading, "Limitations on spot treatments to residential turfgrass."
 - c. Remove the word "General" from the subheading, "General Precautions and Restrictions" as this word is an implied safety claim and makes all associated text unenforceable.
 - d. Per the product chemistry review, the following statement in the "Storage and Disposal" section should be removed from the label:
"Do not contaminate water, food, or feed by storage and disposal."
In addition, the Storage and Disposal statements should be placed in a text box for increased prominence.

Please Note: Final product reregistration cannot be considered until after all active ingredients in this product are eligible for reregistration.


Continued on Page 2

20/12

Page 2
EPA Reg. No. 2217-903
RED Amendments

A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Sincerely,



Kathryn V. Montague
Project Manager 23
Herbicide Branch
Registration Division (7505P)

4012



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Hygiene Statements:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to algae, fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

PRODUCT FACTS:	
<ul style="list-style-type: none"> Pint Containers: Treats up to 8,000 square feet of southern turfgrass and/or up to 4,000 square feet of northern turfgrass. Quart Containers: Treats up to 16, 000 square feet and/or up to 8,000 square feet of northern turfgrass. Gallon Containers: Treats up to 64,000 square feet and/or up to 32,000 square feet of northern turfgrass. 	
KILLS WEEDS:	Dandelion, clover, chickweed plus over xxx additional broadleaf weeds
WHERE TO USE:	Intended for use only on home lawns.
AMOUNT TO USE:	<p>For Southern turf: [bentgrass, hybrid bermudagrass, buffalograss, St. Augustinegrass (except 'Floratam') and centipedegrass] Use: 2 fluid ounces (4 Tablespoons) / 1,000 square feet Or Use: 0.5 fl. oz. per gallon of water per 250 sq.ft.</p> <hr/> <p>For Northern turf: [Kentucky bluegrass, fescues, perennial ryegrass, bahiagrass, common bermudagrass and zoysiagrass] Use: [3 to] 4 fluid ounces ([6 to] 8 Tablespoons) / 1,000 square feet [use higher rate for dense weed populations] Or Use: [0.75 to] 1.0 fl. oz. per gallon of water per 250 sq.ft. [use higher rate for dense weed populations]</p>
	Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx • http://www.company_name.com

WHERE TO USE:

EH-1446 Herbicide is intended for use only on residential (home) lawns including Kentucky bluegrass, perennial ryegrass, fescues, common bermudagrass, hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass (except 'Floratam').

A. For [hand-operated] tank sprayers

[SPOT TREATMENTS WITH HAND-OPERATED TANK SPRAYERS, PUMP-UP SPRAYERS OR COMPRESSED AIR SPRAYERS:]

Spot treat broadleaf weeds with this product as they appear. Repeat treatment for hard-to-control weeds in 2 to 3 weeks 30 days. Do not exceed specified dosages for any area.

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (do not use on 'Floratam' St Augustinegrass):

- Mix 1/2 fluid ounce (1 Tablespoon) of product per gallon of water to cover 250 square feet of lawn.

60912

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

- Mix [0.75 to] 1 fluid ounce ([1.5 to] 2 Tablespoons) of product per gallon of water to cover 250 square feet of lawn. [Use the higher rate for dense weed populations.]

[Equal measures: 1 Tablespoon (Tbs.) = 3 Teaspoons (tsp.) = ½ fluid ounce (fl. oz.)]

Limitations on spot treatments to residential turfgrass:

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl. oz. per 1,000 sq.ft. per application (0.08 lb. MCP-P acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

B. For Ortho® Dial-N-Spray® Applicators:

[optional - THREE EASY STEPS TO MIX AND APPLY

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying.
3. See below for FILLING the Ortho® Dial-N-Spray® hose-end sprayer. Spray evenly over the measured area.]

[Narrative format:]

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)

- ◆ Determine the *total* lawn area to be sprayed.
- ◆ RATE PER GALLON: Set the dial to 1 oz.
- ◆ FILLING: Add 2.0 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar.
- ◆ Spray evenly over the measured area.

Or

- ◆ RATE PER GALLON: Set the dial to 1 oz.
- ◆ FILLING: Add 10 fl.oz of product into sprayer jar. Do not add water to jar.
- ◆ Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

- ◆ Determine the *total* lawn area to be sprayed.
- ◆ RATE PER GALLON: Set the dial to 1½ oz.
- ◆ FILLING: Add [3.0 to] 4.0 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar. [Use the higher rate for dense weed populations]
- ◆ Spray evenly over the measured area.

Or

- ◆ RATE PER GALLON: Set the dial to 1½ oz.
- ◆ FILLING: Add [15 to] 20 fl.oz of product into sprayer jar. Do not add water to jar. [Use the higher rate for dense weed populations]
- ◆ Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.

Optional Tabular Format

Lawn Grass	Sprayer Type	Dial Setting	Product To Measure	Area to Treat
Bentgrass, hybrid bermudagrass, buffalograss centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)	Tank Sprayer	N/A	½ fl.oz. (1 Tbs.) into 1 gallon of water	250 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	1 oz.	Add 2.0 fl.oz. (4Tbs.) into sprayer jar for each 1,000 square feet. Do not add water to jar.	1,000 sq.ft.
Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass	Tank Sprayer	N/A	[0.75 to] 1 fl.oz. ([1.5 to] 2 Tbs.) into 1 gallon of water. [Use the higher rate for dense weed populations]	250 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	1½ oz.	Add [3.0 to] 4.0 fl.oz. ([6 to] 8Tbs.) into sprayer jar for each 1,000 square feet. Do not add water to the jar. [Use the higher rate for dense weed populations]	1,000 sq.ft.

[Equal Measures: 1 Tablespoons = 3 Teaspoons (tsp.) = ½ fl. oz.]

C. Ready-to-use hose-end sprayers

[optional: THREE EASY STEPS TO MIX AND APPLY

1. Measure the *total* (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. See below for directions for the ready-to-use hose-end sprayer
3. Spray evenly over the lawn. }

AMOUNT TO SPRAY

Use the amount indicated for your type of lawn:

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floritam'):

(Quart size): This container covers approximately 16,000 square feet.

(Pint Size): This container covers approximately 8,000 square feet

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

(Quart size): This container covers approximately 8,000 square feet.

(Pint size): This container covers approximately 4,000 square feet.

BROADLEAF WEEDS CONTROLLED:

EH-1446 Herbicide will control or suppress the following broadleaf weeds. A partial list is presented below.

Amaranth, purple	Daisy, English	Mustard, Indian	Speedwell, slender
Amaranth, slender	Daisy, oxeye	Mustard, tansy	Speedwell, snow
Aster, heath	Dandelion, common	Mustard, tumble	Speedwell, thyme-leaf
Aster, white heath	Dandelion, false	Mustard, white	Spiny amaranth
Aster, purple	Day-flower	Mustard, wormseed	Spurge, spotted
Aster, white prairie	Deadnettle	Nettle, stinging	Spurge, prostrate
Artichoke	Dichondra	Nettle, tall	Star of Bethlehem
Austrian fieldcress	Dock, broadleaf	Nettle, wood	Strawberry, Indian mock
Bedstraw, catch-weed	Dock, curly	Parsley-piert	Tansy ragwort
Bedstraw, smooth	Dogbane	Parsnip	Tanweed
Bedstraw, northern	Dog-fennel	Pearlwort	Thistle, blessed
Bedstraw, pineywoods	Dollarweed (Pennywort)	Pennycress	Thistle, bull
Beggar-ticks, hairy	Dove-weed	Pepperweed, perennial	Thistle, Canada
Betony, Florida	Elderberry	Pepperweed, Virginia	Thistle, Flodman's
Bindweed, field	Falseflax	(Peppergrass)	Thistle, musk
Bindweed, hedge	False sunflower	Pigweed, prostrate	Thistle, plumeless
Bitter Wintercress	Fiddleneck	Pigweed, redroot	Thistle, prairie
Bitter-cress, hairy	Filaree, redstem	Pigweed, Russian	Thistle, Russian
Bitterweed	Filaree, whitestem	Pigweed, smooth	Thistle, Scotch
Black-eyed Susan	Fleabane, daisy (annual)	Pigweed, tumble	Thistle, wavy-leaf
Black medic	Fleabane, rough	Pineapple weed	Thistle, yellow
Blood-flower milkweed	Galinsoga, hairy	Plains coreopsis (tickseed)	Trailing crown-vetch
Blue lettuce	Galinsoga, smallflower	Plantain, black-seed	Velvetleaf
Brassbuttons	Goldenrod	Plantain, bracted	Vervain, blue
Bristly oxtongue	Ground ivy	Plantain, broadleaf	Vervain, hoary
Broomweed	Groundsel	Plantain, buckhorn	Vervain, prostrate
Buckhorn	Gumweed	Plantain, hoary	Vervain, tall
Bull-nettle	Hairy fleabane	Plantain, narrow-leaf	Vetch, bird
Bur-clover	Hawkweed, meadow	Plantain, slender	Vetch, hairy
Burdock	Hawkweed, mouse-ear	Plantain, woolly	Vetch, milk
Burweed, lawn	Hawkweed, orange	Poison ivy	Vetch, narrow-leaf
Buttercup, bulbous	Hawkweed, yellow	Poison oak	Vetch, wild
Buttercup, creeping	Healall	Pokeweed	Virginia creeper
Buttonweed, common (poorjoe)	Heart-leaf drymary	Prairie sunflower	Western clematis
Buttonweed, Virginia	Hemp	Prickly lettuce (compass plant)	Western salsify
Carolina geranium	Henbit	Prickly sida	Wild aster
Carpetweed	Hoary-cress	Prostrate knotweed	Wild buckweat
Catnip	Horsenettle	Puncturevine	Wild carrot
Catsear, spotted (false dandelion)	Horseweed	Purslane, common	Wild four-o'clock
Chamber bitter	Jimsonweed	Pusley, Brazilian	Wild garlic
Chickweed, common	Johnny-jumpup violet	Pusley, Florida	Wild lettuce
Chickweed, mouse-ear	Knawel	Pusley, large flower	Wild marigold
Chickweed, sticky	Knotweed, prostrate	Ragweed, bur	Wild mustard
Chicory	Kochia	Ragweed, common	Wild onion
Cinquefoil	Lambsquarters	Ragweed, lance-leaf	Wild parsnip
Clover, crimson	Lespedeza	Ragweed, western	Wild radish
Clover, Alyce	Mallow, alkali	Redstem filaree	Wild rape
Clover, hop	Mallow, bristly	Rough cinquefoil	Wild strawberry
Clover, rabbit-foot	Mallow, common	Scarlet pimpernel	Wild sweet potato
Clover, red	Mallow, dwarf	Shepherd's purse	Wild violet
Clover, strawberry	Mallow, Venice	Smartweed, ladythumb	Woodsorrel, creeping
Clover, yellow sweet	Marcela	Smartweed, pale	Woodsorrel, violet
Clover, white sweet	Matchweed	Smartweed, Pennsylvania	Woodsorrel, yellow (<i>Oxalis</i>)
Clover, white	Mexican-weed	Smooth chaff-flower	Woolly croton
Cockle	Moneywort	Smooth dock	Wormseed
Cocklebur	Morningglory, bigroot	Sorrel, red (sheep)	Yarrow, common
Cocklebur, spiny	Morningglory, ivyleaf	Sowthistle, annual	Yarrow, western
Common mullein	Morningglory, red	Sowthistle, spiny	Yarrow, yellow
Corn spurry	Morningglory, tall	Spanish needles	Yellow Rocket
Creeping beggar-weed	Morningglory, wolly	Speedwell, bi-lobed	
Creeping Jenny (creeping Charlie)	Mugwort	Speedwell, common	
Cupid's shaving brush	Mustard, ball	Speedwell, corn	
Cudweed	Mustard, black	Speedwell, ivyleaf	
	Mustard, blue	Speedwell, Persian	
	Mustard, hedge	Speedwell, purslane	

90A12

[Optional text for further use clarification]

Spray when broadleaf weeds are young and actively growing for best results.

EH-1446 Herbicide may be applied in the spring, summer or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at ~~2-6 week~~ 30 day intervals are recommended suitable for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.

- **For Established Lawns:** Apply when daytime temperatures are below 90°F.
- **For Newly Seeded Areas:** Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- **For Newly Sodded, Sprigged or Plugged areas:** Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
- **Reseeding:** Treated areas can be reseeded at 2 weeks after the application of this product.
- ~~Limitations: The maximum per application rate is 4.0 fl. oz. of product per 1,000 sq. ft., the maximum number of broadcast applications is two (2) per year, the maximum seasonal rate is 8.0 fl. oz. of product per 1,000 sq. ft. excluding spot treatments.~~
- **Limitations on broadcast applications to residential turfgrass:** The maximum application rate is 4.0 fl. oz. of product per 1,000 sq. ft. per application (0.45 lb 2,4-D ae, 0.08 lb MCP-P ae and 0.05 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl. oz. of product per 1,000 sq. ft. (0.90 lb 2,4-D ae, 0.16 lb MCP-P ae and 0.10 lb dicamba ae per acre per year).
- **Irrigation:** Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- **Mowing:** Avoid mowing within 1-2 days before and after the application of this product.
- **Non-target plants:** Do not apply this product to carpetgrass, dichondra, food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. ~~Avoid spraying exposed roots of ornamentals and trees.~~ [end optional text]

STORAGE & DISPOSAL

~~Do not contaminate water, food, or feed by storage and disposal.~~

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

~~CONTAINER DISPOSAL:~~

- ~~If empty: Do not reuse this container. Place in trash or offer for recycling if available.~~
- ~~If partially filled: Call your local solid waste agency for toll free number for disposal instructions. Never place unused product down any indoor or outdoor drain.~~

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Advertising claims that may be used by supplemental registrants: These claims may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds.
- ◆ Kills weeds - not lawn grasses.
- ◆ Kills weeds - not lawns.
- ◆ For lawns containing Kentucky bluegrass, fescues, bentgrass, perennial ryegrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam'), and buffalograss.
- ◆ Can be applied with hose-end or pump-up (tank) sprayers.
- ◆ Kills (broadleaf) weeds.
- ◆ For northern and southern grasses.
- ◆ Won't harm lawn grasses when used as directed.
- ◆ Kills weeds in your lawn.
- ◆ Formulated for use on northern and sensitive southern lawns.
- ◆ Kills major broadleaf lawn weeds: List of the weed species.
- ◆ Helps provide a beautiful lawn.
- ◆ For cool and warm season lawn grasses.
- ◆ Kills major broadleaf lawn weeds including dandelion, chickweed, and knotweed in lawns.
- ◆ For use on grass lawns especially Kentucky bluegrass, fescues, perennial ryegrass, bentgrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam') and buffalograss.
- ◆ Can be used on St. Augustinegrass (except 'Floritam'), bentgrass, and buffalograss.
- ◆ A Trimec® Turf Herbicide Formulation.
- ◆ Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- ◆ "One quart covers up to xxxx square feet (of southern turfgrass)" or "One pint covers up to xxxx square feet (of southern turfgrass)".
- ◆ Treats up to (Number) square feet.
- ◆ Kills over xxx broadleaf lawn weeds including dandelion and others.
- ◆ Makes up to (Number) gallons of spray.
- ◆ Now for northern and southern lawns.
- ◆ ~~New~~ Formula for northern and southern grasses/lawns.
- ◆ Kills major broadleaf weeds including dandelion, clover, oxalis, chickweed, knotweed plantain, henbit, spurge and others.
- ◆ Concentrate
- ◆ Specially formulated for residential use (home lawns)
 - ◇ Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge and many others.
 - ◇ One easy step to a weed free lawn - Helps promote a beautiful lawn.
 - ◇ Simple easy to use/one step application
 - ◇ Simple and easy.
 - ◇ Use (Brand Name) to kill broadleaf weeds in lawns including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge, and many others. Won't harm bahiagrass, bentgrass, bermudagrass, bluegrass, buffalograss, centipedegrass, fescues, ryegrass or zoysiagrass when used as directed. Can also be used on St. Augustinegrass except 'Floritam'.
 - ◇ 1 pint covers 4,000 – 8, 000 sq.ft. (1 quart covers 8,000 – 16,000 sq.ft.) depending on lawn. See inside booklet for complete instructions. Specially formulated for residential use.

- ◇ CONVENIENT.
- ◇ ECONOMICAL - Use to cover small or large areas.
- ◇ Rapid and effective weed control for common and troublesome weed species in turfgrass.
- ◇ Superior cool weather performance.
- ◇ Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed in hours and plant death can occur within 7-14 days.
- ◇ Kills the weeds, not the lawn.
- ◇ Kills broadleaf weeds
- ◇ Won't harm your lawn when used as directed.
- ◇ Kills hard to control weeds.
- ◇ Kills over xxxx tough lawn weeds.
- ◇ Maximum weed control
- ◇ Kills weeds, not lawns
- ◇ The performance leader
- ◇ Treats up to xxxx sq.ft.
- ◇ Tough on weeds, easy on lawns
- ◇ Results in 24 hours
- ◇ Kills over 250 weeds
- ◇ Kills the toughest weeds

~~◇ Rain-Proof™ Rain or watering more than 1 hour after application will not wash away effectiveness~~

~~◇ Rain-Proof™ in 1 hour~~

~~◇ Rain-Fast in 1 hour~~

◆ **Notice:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

Guarantee: If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group