


2217-904

05/04/2007

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number
2217-904

Date of Issuance:
MAY 4 2007

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance: conditional

Name of Pesticide Product:
EH-1447 Herbicide

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation
P.O. Box 014090
Kansas City, Missouri 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label change:

a. Revise the EPA Registration Number to read: "EPA Reg. No. 2217-904."

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

MAY 4 2007

2/11

Page Number 2
EPA Reg. No. 2217-904

c. Under the "Environmental Hazard" section of the labeling, delete the "Consumer Label Initiative Option" statement.

3. Submit storage stability and corrosion characteristics studies (830.6317 and 830.6320), respectively within one year of the date of this registration notice.

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(c). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

3/11

EH-1447 Herbicide

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	4.82%
Mecoprop-p acid	0.59%
Dicamba acid	0.32%
Carfentrazone-ethyl	0.19%
INERT INGREDIENTS:	94.08%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

0.27 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 3.20%.
Isomer Specific by AOAC Methods.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

MAY 4 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

2217-904

First Aid

If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed: <i>(optional for Toxicity Category IV)</i>	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

See side panels for additional Precautionary Statements

**NET CONTENTS: 1.0 pint, 1.0 quart and
1.0 U.S. Gallon**

AP-----

EPA File Symbol No. 2217-

EPA Est. No. 2217-KS-1

MANUFACTURED BY:



Telephone: 1-800-821-7925

4/11



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves.

Personal Hygiene Statements:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS:

[Traditional Statements Option]

This pesticide is toxic to algae, fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Consumer Label Initiative Option]


This pesticide is toxic to fish and aquatic invertebrates. Do not apply near water, storm drains or drainage ditches. Do not apply when windy or prior to heavy rainfall. Rinse applicator over lawn away from storm drains.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

PRODUCT FACTS:	
<ul style="list-style-type: none"> • <i>Pint Containers:</i> Treats up to 6,400 square feet of southern turfgrass and/or up to 3,200 square feet of northern turfgrass. • <i>Quart Containers:</i> Treats up to 12,800 square feet and/or up to 6,400 square feet of northern turfgrass. • <i>Gallon Containers:</i> Treats up to 51,200 square feet and/or up to 25,600 square feet of northern turfgrass. 	
KILLS WEEDS:	Dandelion, clover, chickweed plus over xxx additional broadleaf weeds
WHERE TO USE:	Intended for use only on home lawns.
AMOUNT TO USE:	<p>For Southern turf: [bentgrass, hybrid bermudagrass, buffalograss St. Augustinegrass (except 'Floritam') and centipedegrass] Use: 2.5 fluid ounces (5 Tablespoons) / 1,000 square feet Or Use: 1.0 fl. oz. per gallon of water per 400 sq.ft.</p> <hr/> <p>For Northern turf: [Kentucky bluegrass, fescues, perennial ryegrass, bahiagrass, common bermudagrass and zoysiagrass] Use: [4 to] 5 fluid ounces ([8 to] 10 Tablespoons) / 1,000 square feet [use higher rate for dense weed populations] Or Use: [2.0 to] 2.5 fl. oz. per gallon of water per 500 sq.ft. [use higher rate for dense weed populations]</p>
	Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx • http://www.company_name.com

WHERE TO USE:

EH-1447 Herbicide is intended for use only on residential (home) lawns including Kentucky bluegrass, perennial ryegrass, fescues, common bermudagrass, hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass (except 'Floritam').

A. For [hand-operated] tank sprayers

[SPOT TREATMENTS WITH HAND-OPERATED TANK SPRAYERS, PUMP-UP SPRAYERS OR COMPRESSED AIR SPRAYERS:]

Spot treat broadleaf weeds with this product as they appear. Repeat treatment for hard-to-control weeds in 2 to 3 weeks. Do not exceed specified dosages for any area.

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floritam'):

- Mix 1 fluid ounce (2 Tablespoon) of product per gallon of water to cover 400 square feet of lawn.

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

- Mix [2.0 to] 2.5 fluid ounce ([4 to] 5 Tablespoons) of product per gallon of water to cover 500 square feet of lawn. [Use the higher rate for dense weed populations.]

[Equal measures: 1 Tablespoon (Tbs.) = 3 Teaspoons (tsp.) = 1/2 fluid ounce (fl. oz.)]

6/11

B. For Ortho® Dial-N-Spray® hose-end sprayers

[optional - THREE EASY STEPS TO MIX AND APPLY

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying.
3. See below for FILLING the Ortho® Dial-N-Spray® hose-end sprayer. Spray evenly over the measured area.]

[Narrative format:]

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)

- ◆ Determine the *total* lawn area to be sprayed.
- ◆ RATE PER GALLON: Set the dial to 1 oz.
- ◆ FILLING: Add 2.5 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar.
- ◆ Spray evenly over the measured area.

Or

- ◆ RATE PER GALLON: Set the dial to 1 oz.
- ◆ FILLING: Add 12.5 fl.oz of product into sprayer jar. Do not add water to jar.
- ◆ Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

- ◆ Determine the *total* lawn area to be sprayed.
- ◆ RATE PER GALLON: Set the dial to 2 oz.
- ◆ FILLING: Add [4.0 to] 5.0 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar. [Use the higher rate for dense weed populations]
- ◆ Spray evenly over the measured area.

Or

- ◆ RATE PER GALLON: Set the dial to 2 oz.
- ◆ FILLING: Add [20 to] 25 fl.oz of product into sprayer jar. Do not add water to jar.
- ◆ Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.

Tabular Format

7/11

Lawn Grass	Sprayer Type	Dial Setting	Product To Measure	Area to Treat
Bentgrass, hybrid bermudagrass, buffalograss centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)	Tank Sprayer	NA	1 fl.oz. (2 Tbs.) into 1 gallon of water	400 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	1 oz.	Add 2.5 fl.oz. (5Tbs.) into sprayer jar for each 1,000 square feet. Do not add water to jar.	1,000 sq.ft.
Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass	Tank Sprayer	NA	2 fl.oz. (4 Tbs.) into 1 gallon of water.	400 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	2 oz.	Add [4.0 to] 5.0 fl.oz. ([8 to] 10Tbs.) into sprayer jar for each 1,000 square feet. Do not add water to the jar. [Use the higher rate for dense weed populations]	1,000 sq.ft.

[Equal Measures: 1 Tablespoons = 3 Teaspoons (tsp.) = ½ fl. oz.]

C. Ready-to-use hose-end sprayers

[optional: THREE EASY STEPS TO MIX AND APPLY

1. Measure the *total* (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the lawn.]

AMOUNT TO SPRAY

Use the amount indicated for your type of lawn:

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floritam'):

(Quart size): This container covers approximately 12,800 square feet.

(Pint size): This container covers approximately 6,400 square feet

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

(Quart size): This container covers approximately 6,400 square feet

(Pint size): This container covers approximately 3,200 square feet

BROADLEAF WEEDS CONTROLLED:

EH-1447 Herbicide will control or suppress the following broadleaf weeds. A partial list is presented below.

Amaranth, purple	Daisy, English	Mustard, Indian	Speedwell, slender
Amaranth, slender	Daisy, oxeye	Mustard, tansy	Speedwell, snow
Aster, heath	Dandelion, common	Mustard, tumble	Speedwell, thyme-leaf
Aster, white heath	Dandelion, false	Mustard, white	Spiny amaranth
Aster, purple	Day-flower	Mustard, wormseed	Spurge, spotted
Aster, white prairie	Deadnettle	Nettle, stinging	Spurge, prostrate
Artichoke	Dichondra	Nettle, tall	Star of Bethlehem
Austrian fieldcress	Dock, broadleaf	Nettle, wood	Strawberry, Indian mock
Bedstraw, catch-weed	Dock, curly	Parsley-piert	Tansy ragwort
Bedstraw, smooth	Dogbane	Parsnip	Tanweed
Bedstraw, northern	Dog-fennel	Pearlwort	Thistle, blessed
Bedstraw, pineywoods	Dollarweed (Pennywort)	Pennycress	Thistle, bull
Beggar-ticks, hairy	Dove-weed	Pepperweed, perennial	Thistle, Canada
Betony, Florida	Elderberry	Pepperweed, Virginia	Thistle, Flodman's
Bindweed, field	Falseflax	(Peppergrass)	Thistle, musk
Bindweed, hedge	False sunflower	Pigweed, prostrate	Thistle, plumeless
Bitter Wintercress	Fiddleneck	Pigweed, redroot	Thistle, prairie
Bitter-cress, hairy	Filaree, redstem	Pigweed, Russian	Thistle, Russian
Bitterweed	Filaree, whitestem	Pigweed, smooth	Thistle, Scotch
Black-eyed Susan	Fleabane, daisy (annual)	Pigweed, tumble	Thistle, wavy-leaf
Black medic	Fleabane, rough	Pineapple weed	Thistle, yellow
Blood-flower milkweed	Galinsoga, hairy	Plains coreopsis (tickseed)	Trailing crown-vetch
Blue lettuce	Galinsoga, smallflower	Plantain, black-seed	Velvetleaf
Brassbuttons	Goldenrod	Plantain, bracted	Vervain, blue
Bristly oxtongue	Ground ivy	Plantain, broadleaf	Vervain, hoary
Broomweed	Groundsel	Plantain, buckhorn	Vervain, prostrate
Buckhorn	Gumweed	Plantain, hoary	Vervain, tall
Bull-nettle	Hairy fleabane	Plantain, narrow-leaf	Vetch, bird
Bur-clover	Hawkweed, meadow	Plantain, slender	Vetch, hairy
Burdock	Hawkweed, mouse-ear	Plantain, woolly	Vetch, milk
Burweed, lawn	Hawkweed, orange	Poison ivy	Vetch, narrow-leaf
Buttercup, bulbous	Hawkweed, yellow	Poison oak	Vetch, wild
Buttercup, creeping	Healall	Pokeweed	Virginia creeper
Buttonweed, common (poorjoe)	Heart-leaf drymary	Prairie sunflower	Western clematis
Buttonweed, Virginia	Hemp	Prickly lettuce (compass plant)	Western salsify
Carolina geranium	Henbit	Prickly sida	Wild aster
Carpetweed	Hoary-cress	Prostrate knotweed	Wild buckweat
Catnip	Horsenettle	Puncturevine	Wild carrot
Catsear, spotted (false dandelion)	Horseweed	Purslane, common	Wild four-o'clock
Chamber bitter	Jimsonweed	Pusley, Brazilian	Wild garlic
Chickweed, common	Johnny-jumpup violet	Pusley, Florida	Wild lettuce
Chickweed, mouse-ear	Knawel	Pusley, large flower	Wild marigold
Chickweed, sticky	Knotweed, prostrate	Ragweed, bur	Wild mustard
Chicory	Kochia	Ragweed, common	Wild onion
Cinquefoil	Lambsquarters	Ragweed, lance-leaf	Wild parsnip
Clover, crimson	Lespedeza	Ragweed, western	Wild radish
Clover, Alyce	Mallow, alkali	Redstem filaree	Wild rape
Clover, hop	Mallow, bristly	Rough cinquefoil	Wild strawberry
Clover, rabbit-foot	Mallow, common	Scarlet pimpernel	Wild sweet potato
Clover, red	Mallow, dwarf	Shepherd's purse	Wild violet
Clover, strawberry	Mallow, Venice	Smartweed, ladysthumb	Woodsorrel, creeping
Clover, yellow sweet	Marcela	Smartweed, pale	Woodsorrel, violet
Clover, white sweet	Matchweed	Smartweed, Pennsylvania	Woodsorrel, yellow (<i>Oxalis</i>)
Clover, white	Mexican-weed	Smooth chaff-flower	Woolly croton
Cockle	Moneywort	Smooth dock	Wormseed
Cocklebur	Morningglory, bigroot	Sorrel, red (sheep)	Yarrow, common
Cocklebur, spiny	Morningglory, ivyleaf	Sowthistle, annual	Yarrow, western
Common mullein	Morningglory, red	Sowthistle, spiny	Yarrow, yellow
Corn spurry	Morningglory, tall	Spanish needles	Yellow Rocket
Creeping beggar-weed	Morningglory, wolly	Speedwell, bi-lobed	
Creeping Jenny (creeping Charlie)	Mugwort	Speedwell, common	
Cupid's shaving brush	Mustard, ball	Speedwell, corn	
Cudweed	Mustard, black	Speedwell, ivyleaf	
	Mustard, blue	Speedwell, Persian	
	Mustard, hedge	Speedwell, purslane	

9/11

[Optional text for further use clarification]

Spray when broadleaf weeds are young and actively growing for best results.

EH-1447 Herbicide may be applied in the spring, summer or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2 - 6 week intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.

- **For Established Lawns:** Apply when daytime temperatures are below 90°F.
- **For Newly Seeded Areas:** Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- **For Newly Sodded, Sprigged or Plugged areas:** Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
- **Reseeding:** Treated areas can be reseeded at 2 weeks after the application of this product.
- **Limitations:** The maximum per application rate is 5.0 fl. oz. of product per 1,000 sq.ft., the maximum number of broadcast applications is two (2) per year; the maximum seasonal rate is 10.0 fl. oz. of product per 1,000 sq.ft. excluding spot treatments.
- **Irrigation:** Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- **Mowing:** Avoid mowing within 1-2 days before and after the application of this product.
- **Non-target plants:** Do not apply this product to carpetgrass, dichondra, food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. Avoid spraying exposed roots of ornamentals and trees. [end optional text]

STORAGE & DISPOSAL

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency [or toll free number (1-800-CLEANUP)] for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

i. Advertising claims that may be used by supplemental registrants: These claims may be presented on the retail container label or on the labeling accompanying the product.

- ◆ Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds.
- ◆ Kills weeds - not lawn grasses.
- ◆ Kills weeds - not lawns.
- ◆ For lawns containing Kentucky bluegrass, fescues, bentgrass, perennial ryegrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam'), and buffalograss.
- ◆ Can be applied with hose-end or pump-up (tank) sprayers.
- ◆ Kills (broadleaf) weeds.
- ◆ For northern and southern grasses.
- ◆ Won't harm lawn grasses when used as directed.
- ◆ Kills weeds in your lawn.
- ◆ Formulated for use on northern and sensitive southern lawns.
- ◆ Kills major broadleaf lawn weeds: List of the weed species.
- ◆ Helps provide a beautiful lawn.
- ◆ For cool and warm season lawn grasses.
- ◆ Kills major broadleaf lawn weeds including dandelion, chickweed, and knotweed in lawns.
- ◆ For use on grass lawns especially Kentucky bluegrass, fescues, perennial ryegrass, bentgrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam') and buffalograss.
- ◆ Can be used on St. Augustinegrass (except 'Floritam'), bentgrass, and buffalograss.
- ◆ A Trimec® Turf Herbicide Formulation.
- ◆ Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- ◆ "One quart covers up to xxxx square feet (of southern turfgrass)" or "One pint covers up to xxxx square feet (of southern turfgrass)".
- ◆ Treats up to (Number) square feet.
- ◆ Kills over xxx broadleaf lawn weeds including dandelion and others.
- ◆ Makes up to (Number) gallons of spray.
- ◆ Now for northern and southern lawns.
- ◆ New formula for northern and southern grasses/lawns.
- ◆ Kills major broadleaf weeds including dandelion, clover, oxalis, chickweed, knotweed plantain, henbit, spurge and others.
- ◆ Concentrate

- ◆ Specially formulated for residential use (home lawns)
 - ◇ Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweec, knotweed, plantain, henbit, spurge and many others.
 - ◇ One easy step to a weed free lawn - Helps promote a beautiful lawn.
 - ◇ Simple easy to use/one step application
 - ◇ Simple and easy.
 - ◇ Use (Brand Name) to kill broadleaf weeds in lawns including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge, and many others. Won't harm bahiagrass, bentgrass, bermudagrass, bluegrass, buffalograss, centipedegrass, fescues, ryegrass or zoysiagrass when used as directed. Can also be used on St. Augustinegrass except 'Floritam'.
 - ◇ 1 pint covers 4,000 – 8, 000 sq.ft. (1 quart covers 8,000 – 16,000 sq.ft.) depending on lawn. See inside booklet for complete instructions. Specially formulated for residential use.
 - ◇ *CONVENIENT.*
 - ◇ *ECONOMICAL* - Use to cover small or large areas.
 - ◇ Rapid and effective weed control for common and troublesome weed species in turfgrass.
 - ◇ Superior cool weather performance.
 - ◇ Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed in hours and plant death can occur within 7-14 days.
 - ◇ Kills the weeds, not the lawn.
 - ◇ Kills broadleaf weeds
 - ◇ Won't harm your lawn when used as directed.
 - ◇ Kills hard to control weeds.
 - ◇ Kills over xxxx tough lawn weeds.
 - ◇ Maximum weed control
 - ◇ Kills weeds, not lawns
 - ◇ The performance leader
 - ◇ Treats up to xxxx sq.ft.
 - ◇ Tough on weeds, easy on lawns
 - ◇ Results in 24 hours
 - ◇ Kills over 250 weeds
 - ◇ Kills the toughest weeds
 - ◇ Rain-Proof™ - Rain or watering more than 1 hour after application will not wash away effectiveness
 - ◇ Rain-Proof™ in 1 hour
 - ◇ Rain Fast in 1 hour
- ◆ **Notice:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

Guarantee: If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group