



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

March 25, 2025

Eric D. Smith
Director, Regulatory Affairs
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, KS 66286

Subject: Label Amendment - Registration Review Mitigation for Carfentrazone-ethyl & Mecoprop-p
Product Name: EH-1447 Herbicide
EPA Registration Number: 2217-904
Application Dates: May 1, 2018; July 5, 2022; & February 2, 2024
Decision Numbers: 590534 & 596738
Case Numbers: 474330

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Carfentrazone-ethyl & Mecoprop-p Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at (202) 566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read 'Linda Arrington', with a stylized flourish at the end.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

EH-1447 HERBICIDE

EPA Reg. No. 2217-904

ACCEPTED

Mar 25, 2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 2217-904

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	4.82%
Mecoprop-p acid	0.59%
Dicamba acid	0.32%
Carfentrazone-ethyl	0.19%
INERT INGREDIENTS:	94.08%
TOTAL.....	100.00%

THIS PRODUCT CONTAINS:

0.27 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 3.20%.
Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

First Aid	
If on skin:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 – 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Fish Advisory Statement: This product may be hazardous to aquatic organisms, particularly in clear, shallow water bodies that are adjacent to treated areas. Transport to water by runoff or spray drift of this product in areas where surface water is present, or intertidal areas below the mean high water mark, should be avoided. Do not contaminate water when disposing of equipment wash water or rinsate.

Non-target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Only use for sites, pests, and application methods specified on this labeling.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- The maximum broadcast application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application [(0.46 lb 2,4-D ae, 0.09 lb MCPP-p ae, 0.05 lb dicamba ae, and 0.03 carfentrazone ai per acre per application)].
- The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications.
- The maximum annual broadcast application rate is 10.0 fl. oz. of product per 1,000 sq.ft. [(0.92 lb 2,4-D ae, 0.18 lb MCPP-p ae, 0.10 lb dicamba ae, and 0.06 carfentrazone ai per acre per year)].
- Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 5.0 fl. oz. per 1,000 sq.ft. per application [(0.46 lb 2,4-D ae, 0.09 lb MCPP-p ae, 0.05 lb dicamba ae, and 0.03 carfentrazone ai per acre per application)]. *{Note to Reviewer: This bullet may be omitted if Option 1 directions are not presented on the market label.}*

- The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications. *{Note to Reviewer: This bullet may be omitted if Option 1 directions are not presented on the market label.}*

PRODUCT FACTS	Pint Containers: Treats up to 6,400 square feet of southern turfgrass and/or up to 3,200 square feet of northern turfgrass. Quart Containers: Treats up to 12,800 square feet and/or up to 6,400 square feet of northern turfgrass. Gallon Containers: Treats up to 51,200 square feet and/or up to 25,600 square feet of northern turfgrass.
KILLS WEEDS:	Dandelion, clover, chickweed plus over XXX additional broadleaf weeds
WHERE TO USE:	Intended for use only on home lawns
AMOUNT TO USE:	For southern turf: [bentgrass, hybrid bermudagrass, buffalograss, St. Augustinegrass (except 'Floratam') and centipedegrass] Use: 2.5 fl.oz. (5 Tbsp)/1,000 sq.ft. OR Use: 1.0 fl.oz. per gallon of water per 400 sq.ft. For Northern turf: [Kentucky bluegrass, fescues, perennial ryegrass, bahiagrass, common bermudagrass and zoysiagrass] Use: [4 to] 5 fl.oz. ([8 to] 10 Tbsp)/1,000 sq.ft. [use higher rate for dense weed populations] OR Use: [2.0 to] 2.5 fl.oz. per gallon of water per 500 sq.ft. [use higher rate for dense weed populations]
	Questions, Comments or Medical Information? Call: 1-800-XXX-XXXX http://www.companyname.com

WHERE TO USE:

EH-1447 Herbicide is intended for use only on residential (home) lawns including Kentucky bluegrass, perennial ryegrass, fescues, common bermudagrass, hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass (except 'Floratam').

OPTION 1. For [hand-operated] tank sprayers:

[SPOT TREATMENTS WITH HAND-OPERATED TANK SPRAYERS, PUMP-UP SPRAYERS OR COMPRESSED AIR SPRAYERS:]

Spot treat broadleaf weeds with this product as they appear. Repeat treatment for hard-to-control weeds in 30 days. Do not exceed specified dosages for any area.

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floratam'): Mix 1 fluid ounce (2 Tablespoon) of product per gallon of water to cover 400 square feet of lawn.

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass: Mix [2.0 to] 2.5 fluid ounce ([4 to] 5 Tablespoons) of product per gallon of water to cover 500 square feet of lawn. [Use the higher rate for dense weed populations.]

[Equal measures: 1 Tablespoon (Tbs.) = 3 Teaspoons (tsp.) = ½ fluid ounce (fl. oz.)]

OPTION 2. For Ortho® Dial-N-Spray® hose-end sprayers:

[Note to Supplemental Distributor: Choose an option below, various formats presented]

[THREE EASY STEPS TO MIX AND APPLY]

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying.
3. See below for FILLING the Ortho® Dial-N-Spray® hose-end sprayer. Spray evenly over the measured area.]

[For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)]

- Determine the *total* lawn area to be sprayed.
- RATE PER GALLON: Set the dial to 1 oz.
- FILLING: Add 2.5 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar.
- Spray evenly over the measured area.

OR

- RATE PER GALLON: Set the dial to 1 oz.
- FILLING: Add 12.5 fl.oz of product into sprayer jar. Do not add water to jar.
- Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.]

[For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

- Determine the *total* lawn area to be sprayed.
- RATE PER GALLON: Set the dial to 2 oz.
- FILLING: Add [4.0 to] 5.0 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar. [Use the higher rate for dense weed populations]
- Spray evenly over the measured area.

OR

- RATE PER GALLON: Set the dial to 2 oz.
- FILLING: Add [20 to] 25 fl.oz of product into sprayer jar. Do not add water to jar.
- Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.]

[Optional Tabular Format]

Lawn Grass	Sprayer Type	Dial Setting	Product To Measure	Area to Treat
Bentgrass, hybrid bermudagrass, buffalograss centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)	Tank Sprayer	NA	1 fl.oz. (2 Tbs.) into 1 gallon of water	400 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	1 oz.	Add 2.5 fl.oz. (5Tbs.) into sprayer jar for each 1,000 square feet. Do not add water to jar.	1,000 sq.ft.
Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and	Tank Sprayer	NA	2 fl.oz. (4 Tbs.) into 1 gallon of water.	400 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed	2 oz.	Add [4.0 to] 5.0 fl.oz. ([8 to] 10Tbs.) into sprayer jar for each	1,000 sq.ft.

zoysiagrass	Sprayette®		1,000 square feet. Do not add water to the jar. [Use the higher rate for dense weed populations]	
-------------	------------	--	--	--

[Equal Measures: 1 Tablespoons = 3 Teaspoons (tsp.) = ½ fl. oz.]

OPTION 3. Ready-to-use hose-end sprayers

[THREE EASY STEPS TO MIX AND APPLY

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the lawn.

AMOUNT TO SPRAY

Use the amount indicated for your type of lawn:

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floritam'): [Quart Size] This container covers approximately 12,800 square feet. [Pint Size] This container covers approximately 6,400 square feet

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass: [Quart Size] This container covers approximately 6,400 square feet. [Pint Size] This container covers approximately 3,200 square feet

BROADLEAF WEEDS CONTROLLED:

EH-1447 Herbicide will control or suppress the following broadleaf weeds.

Amaranth, purple	Cudweed	Mustard, blue	Speedwell, ivyleaf
Amaranth, slender	Daisy, English	Mustard, hedge	Speedwell, Persian
Aster, heath	Daisy, oxeye	Mustard, Indian	Speedwell, purslane
Aster, white heath	Dandelion, common	Mustard, tansy	Speedwell, slender
Aster, purple	Dandelion, false	Mustard, tumble	Speedwell, snow
Aster, white prairie	Day-flower	Mustard, white	Speedwell, thyme-leaf
Artichoke	Deadnettle	Mustard, wormseed	Spiny amaranth
Austrian fieldcress	Dichondra	Nettle, stinging	Spurge, spotted
Bedstraw, catch-weed	Dock, broadleaf	Nettle, tall	Spurge, prostrate
Bedstraw, smooth	Dock, curly	Nettle, wood	Star of Bethlehem
Bedstraw, northern	Dogbane	Parsley-piert	Strawberry, Indian mock
Bedstraw, pineywoods	Dog-fennel	Parsnip	Tansy ragwort
Beggar-ticks, hairy	Dollarweed (Pennywort)	Pearlwort	Tanweed
Betony, Florida	Dove-weed	Pennycress	Thistle, blessed
Bindweed, field	Elderberry	Pepperweed, perennial	Thistle, bull
Bindweed, hedge	Falseflax	Pepperweed, Virginia	Thistle, Canada
Bitter Wintercress	False sunflower	(Peppergrass)	Thistle, Flodman's
Bitter-cress, hairy	Fiddleneck	Pigweed, prostrate	Thistle, musk
Bitterweed	Filaree, redstem	Pigweed, redroot	Thistle, plumeless
Black-eyed Susan	Filaree, whitestem	Pigweed, Russian	Thistle, prairie
Black medic	Fleabane, daisy (annual)	Pigweed, smooth	Thistle, Russian
Blood-flower milkweed	Fleabane, rough	Pigweed, tumble	Thistle, Scotch
Blue lettuce	Galinsoga, hairy	Pineapple weed	Thistle, wavy-leaf
Brassbuttons	Galinsoga, smallflower	Plains coreopsis (tickseed)	Thistle, yellow
Bristly oxtongue	Goldenrod	Plantain, black-seed	Trailing crown-vetch
Broomweed	Ground ivy	Plantain, bracted	Velvetleaf
Buckhorn	Groundsel	Plantain, broadleaf	Vervain, blue
Bull-nettle	Gumweed	Plantain, buckhorn	Vervain, hoary
Bur-clover	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Burdock	Hawkweed, meadow	Plantain, narrow-leaf	Vervain, tall
Burweed, lawn	Hawkweed, mouse-ear	Plantain, slender	Vetch, bird
Buttercup, bulbous	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Buttercup, creeping	Hawkweed, yellow	Poison ivy	Vetch, milk

Buttonweed, common (poorjoe)	Healall	Poison oak	Vetch, narrow-leaf
Buttonweed, Virginia	Heart-leaf drymary	Pokeweed	Vetch, wild
Carolina geranium	Hemp	Prairie sunflower	Virginia creeper
Carpetweed	Henbit	Prickly lettuce (compass plant)	Western clematis
Catnip	Hoary-cress	Prickly sida	Western salsify
Catsear, spotted (false dandelion)	Horsenettle	Prostrate knotweed	Wild aster
Chamber bitter	Horseweed	Puncturevine	Wild buckweat
Chickweed, common	Jimsonweed	Purslane, common	Wild carrot
Chickweed, mouse-ear	Johnny-jumpup violet	Pusley, Brazilian	Wild four-o'clock
Chickweed, sticky	Knawel	Pusley, Florida	Wild garlic
Chicory	Knotweed, prostrate	Pusley, large flower	Wild lettuce
Cinquefoil	Kochia	Ragweed, bur	Wild marigold
Clover, crimson	Lambsquarters	Ragweed, common	Wild mustard
Clover, Alyce	Lespedeza	Ragweed, lance-leaf	Wild onion
Clover, hop	Mallow, alkali	Ragweed, western	Wild parsnip
Clover, rabbit-foot	Mallow, bristly	Redstem filaree	Wild radish
Clover, red	Mallow, common	Rough cinquefoil	Wild rape
Clover, strawberry	Mallow, dwarf	Scarlet pimpernel	Wild strawberry
Clover, yellow sweet	Mallow, Venice	Shepherd's purse	Wild sweet potato
Clover, white sweet	Marcela	Smartweed, ladysthumb	Wild violet
Clover, white	Matchweed	Smartweed, pale	Woodsorrel, creeping
Cockle	Mexican-weed	Smartweed, Pennsylvania	Woodsorrel, violet
Cocklebur	Moneywort	Smooth chaff-flower	Woodsorrel, yellow (<i>Oxalis</i>)
Cocklebur, spiny	Morningglory, bigroot	Smooth dock	Woolly croton
Common mullein	Morningglory, ivyleaf	Sorrel, red (sheep)	Wormseed
Corn spurry	Morningglory, red	Sowthistle, annual	Yarrow, common
Creeping beggar-weed	Morningglory, tall	Sowthistle, spiny	Yarrow, western
Creeping Jenny (creeping Charlie)	Morningglory, wolly	Spanish needles	Yarrow, yellow
Cupid's shaving brush	Mugwort	Speedwell, bi-lobed	Yellow Rocket
	Mustard, ball	Speedwell, common	
	Mustard, black	Speedwell, corn	

[Optional text for further use clarification]

Spray when broadleaf weeds are young and actively growing for best results.

EH-1447 Herbicide may be applied in the spring, summer or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 30 day intervals are suitable for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.

- **For Established Lawns:** Apply when daytime temperatures are below 90°F.
- **For Newly Seeded Areas:** Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- **For Newly Sodded, Sprigged or Plugged areas:** Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
- **Reseeding:** Treated areas can be reseeded at 2 weeks after the application of this product.
- **Irrigation:** Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- **Mowing:** Avoid mowing within 1-2 days before and after the application of this product.
- **Non-target plants:** Do not apply this product to carpetgrass, dichondra, food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.

[end optional text]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

Herbicide Benefit Claims:

- Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds.
- Kills weeds - not lawn grasses.
- Kills weeds - not lawns.
- Kills (broadleaf) weeds.
- Kills weeds in your lawn.
- Kills major broadleaf lawn weeds: List of the weed species.
- Kills major broadleaf lawn weeds including dandelion, chickweed, and knotweed in lawns.
- A Trimec® Turf Herbicide Formulation.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills over XXX broadleaf lawn weeds including dandelion and others.
- Kills major broadleaf weeds including dandelion, clover, oxalis, chickweed, knotweed plantain, henbit, spurge and others.
- Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge and many others.
- One easy step to a weed free lawn - Helps promote a beautiful lawn.
- Use (Brand Name) to kill broadleaf weeds in lawns including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge, and many others. Won't harm bahiagrass, bentgrass, bermudagrass, bluegrass, buffalograss, centipedegrass, fescues, ryegrass or zoysiagrass when used as directed. Can also be used on St. Augustinegrass except 'Floritam'.
- Rapid and effective weed control for common and troublesome weed species in turfgrass.
- Kills the weeds, not the lawn.
- Kills broadleaf weeds
- Kills hard to control weeds.
- Kills over XXX tough lawn weeds.
- Maximum weed control
- Kills weeds, not lawns
- Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed in hours and plant death can occur within 7-14 days.
- Kills over 250 weeds
- Kills the toughest weeds
- Tough on weeds, easy on lawns

General Marketing Claims:

- For lawns containing Kentucky bluegrass, fescues, bentgrass, perennial ryegrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam'), and buffalograss.
- Can be applied with hose-end or pump-up (tank) sprayers.
- For northern and southern grasses.
- Won't harm lawn grasses when used as directed.
- Formulated for use on northern and sensitive southern lawns.
- Helps provide a beautiful lawn.
- For cool and warm season lawn grasses.

- For use on grass lawns especially Kentucky bluegrass, fescues, perennial ryegrass, bentgrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam') and buffalograss.
- Can be used on St. Augustinegrass (except 'Floritam'), bentgrass, and buffalograss.
- "One quart covers up to XXX square feet (of southern turfgrass)" or "One pint covers up to xxxx square feet (of southern turfgrass)".
- Treats up to XXX square feet.
- Makes up to XXX gallons of spray.
- Now for northern and southern lawns.
- Formula for northern and southern grasses/lawns.
- Concentrate
- Specially formulated for residential use (home lawns)
- Simple easy to use/one step application
- Simple and easy.
- 1 pint covers 4,000 – 8, 000 sq.ft. (1 quart covers 8,000 – 16,000 sq.ft.) depending on lawn. See inside booklet for complete instructions. Specially formulated for residential use.
- CONVENIENT.
- ECONOMICAL - Use to cover small or large areas.
- Superior cool weather performance.
- Won't harm your lawn when used as directed.
- The performance leader
- Treats up to XXX sq.ft.
- Results in 24 hours
- **Notice:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.
- **Guarantee:** If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.
- TRIMEC® is a registered trademark of PBI/Gordon Corporation.
- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00904.20240202.amend-proposed-clean.doc

2. Reason for Issue: Comments on MCPPE and Carfentrazone-ethyl Interim Reg Review Decision

Internal Usage Notes
Associated Container Label: