1/14/2014

2217-906



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

JAN 14 2014

James L. Kunstman, Ph.D. PBI/ Gordon Corporation P.O. Box 014090 Kansas City, Missouri 64101

Subject:

Label Notification; Per PR-Notice 98-10

Product Name: EH-1449 Herbicide

EPA Reg. No. 2217-906

Date Submitted: June 27, 2013

Dear Mr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated June 27, 2013 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact Jasmine Branch at (703) 347-0351 or branch.jasmine@epa.gov.

Sincerely,

Kathryn Montague Product Manager 23

Herbicide Branch

Registration Division (7505P)

2)17

Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95 OPP Identifier Number Registration **United States** SEPA **Environmental Protection Agency Amendment** Washington, DC 20460 Other Application for Pesticide - Section I 2. EPA Product Manager 1. Company/Product Number 3. Proposed Classification 2217-906 Kathryn V. Montague PM# 4. Company/Product (Name) X None Restricted EH-1449 Herbicide Product Manager—Team 23 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) PBI/Gordon Corporation (b)(i), my product is similar or identical in composition and labeling to: Post Office Box 014090 EPA Reg. No. Kansas City, Missouri 64101 Product Name Check if this is a new address Section - II Final printed labels in response to Amendment - Explain below. Agency letter dated Resubmission in response to Agency letter dated _ "Me Too" Application. Other - Explain below. Notification - Explain below. Explanation: Use additional page(s) if necessary. (For section I and Section II.) e-mail to jkunstman@pbigordon.com FAX: 816-421-2731 Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments. Section II(A),(N) and IV(E). Please refer to cover letter for details. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA. Section - III 1. Material This Product Will Be Packaged In: 2. Type of Container Child-Resistant Packaging Unit Packaging Water Soluble Packaging Metal Yes* Yes Yes Plastic No Nο Glass * Certification must If "Yes" No. per If "Yes" No. per Paper container Unit Packaging wgt. Package wgt. Container be submitted Other (Specify) 3. Location of Net Contents Information 4. Size(s) Retail Container 5. Location of Label Directions 1 pint, 24 fl. Oz., 1 quart, 2 quarts On Label Label Container and 1 gallon On Labeling accompanying product Other 6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Telephone No. ('ncludo Area Code) Name James L. Kunstman, Ph.D. Director of Regulatory Services 816-460-6292 Certification 6. Date Application I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. Received I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or (Stamped) both under applicable law. 2. Signature Director of Regulatory Services 5. Date 4. Typed Name James L. Kunstman, Ph.D. June 27, 2013



An Employee-Owned Company

1217 W. 12th STREET · P.O. BOX 014090 KANSAS CITY, MISSOURI 64101-0090 816-421-4070 · 1-800-821-7925 FAX: 816-474-0462

June 27, 2013

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Kathryn V. Montague (PM-23)

NOTIFICATION: Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments. Section II(A),(N) and IV(E): for EH-1449 Herbicide (EPA Reg. No. 2217-906).

I. Labeling notification:

Section II (A): Brand Names

We ask to add a new alternate brand name.
 Trimec Crabgrass Plus Lawn Weed Killer Ready-To-Use

Section II (N): Other Revisions

• We ask to add advertising claims that are consistent with the previously approved advertising claims. Please refer to the appendix section of the enclosed draft label on pages 12 and 13.

Section IV (E): Redesign of Label Format

 Update the labeling format to make it consistent with currently approved labels.

II. Certification Statement

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR_Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

EH-1449 HERBICIDE

NOTIFICATION

JAN 1 4 2014

EPA Reg. No. 2217-906

Two sublabels represent the entire master label of EH-1449 Herbicide. Please refer to the appropriate section of the labeling as shown as follows:

Section	
Sublabel 1: Ready-to-use intended for use by homeowners	
Sublabel 2: Ready-to-use intended for use by homeowners and marketed by supplemental registrants.	
Appendix	
Document Control Information	

SUBLABEL 1 Ready-to-use intended for use by homeowners

EH-1449 HERBICIDE

EPA Reg. No. 2217-906

Alternate Brand Names: Gordon's Trimec Crabgrass Plus Lawn Weed Killer Ready-To-Use, Trimec Crabgrass Plus Lawn Weed Killer Ready-To-Use

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	0.313%
Quinclorac	0.104%
Dicamba, dimethylamine salt	0.029%
OTHER INGREDIENTS:	99.554%
TOTAL	

THIS PRODUCT CONTAINS:

- 2,4-dichlorophenoxyacetic acid equivalent, 0.260%
- 3,7-dichloro-8-quinolinecarboxylic acid, 0.104%
- 3,6-dichloro-o-anisic acid equivalent, 0.024%

Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chevring gum, or using tobacco. Remove and wash clothing before reuse.

Applicators and other handlers of this product must wear:

- Long sleeved shirt and long pants
- Shoes
- Socks

First Aid	
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
	ct container or label with you when calling a poison control center or doctor or going for may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
 Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Apply when daytime temperatures are between 50° and 90°F (do not apply above 85°F for bermudagrass). Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed, clover as well as crabgrass and foxtail. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed. [icon weeds]

When to Use: Apply when daytime temperatures are between 50° and 90°F, (do not apply above 85°F for bermudagrass).

Questions & Comments: [icon phone] [icon computer]

Call 1-800-xxx-xxxx or visit our website at www.companyname.com

SHAKE WELL BEFORE USING.

HOW IT WORKS:

EH-1449 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE:

EH-1449 Herbicide is a ready-to-use (RTU) product intended for the treatment of individual weeds, patches of weeds or sparse weeds scattered throughout home lawns. EH-1449 Herbicide can be applied to the following established turfgrass species:

Cool season turfgrass	Warm season turfgrass
Kentucky bluegrass	Bermudagrass, common and hybrid
Ryegrass, perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues	
(red, hard and chewings) and bentgrass	

This product is for spot treatment only.

DO NOT apply EH-1449 Herbicide to the following:

Warm season turfgrass	Other areas
St. Augustinegrass,	Lawns with desirable clovers or legumes,
Bahiagrass,	Dichondra,
Centipedegrass,	Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs),
Carpetgrass,	Turfgrass species that are not listed on this label,
Seashore paspalum	Gardens including vegetables, fruit trees, vines and berries.

WEEDS CONTROLLED

GRASS WEEDS:

crabgrass (large and smooth), foxtail (green, yellow and giant), kikuyugrass, and broadleaf signalorass

BROADLEAF WEEDS: A partial list is presented below:			
Annual yellow sweetclover	Cocklebur	Mustard	Spiny cocklebur
Aster	Common mullein	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Creeping Jenny (Creeping	Narrowleaf vetch	Spotted catsear
Bedstraw	Charlie)	Nettle	Spotted spurge
Beggarticks	Cudweed	Orange hawkweed	Spurweed
Betony, Florida	Curly dock	Oriental cocklebur	Stinging nettle



BROADLEAF WEEDS:	laves.		
A partial list is presented be		10 -1	[Ot-10]
Bindweed, field	Daisy, English	Oxalis	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsley-piert	Tall nettle
Bitter wintercress	Daisy, oxeye	Parsnip	Tall vervain
Bittercress, hairy	Dandelion	Pearlwort	Tansy ragwort
Bitterweed	Dichondra	Pennycress	Tansy mustard
Black-eyed Susan	Dogbane	Pennywort	Tanweed
Black medic	Dogfennel	Peppergrass	Thistle
Black mustard	Dollarweed	Pepperweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pigweed	Tumble mustard
Blessed thistle	False dandelion	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	Falseflax	Plains coreopsis	Velvetleaf
Blue lettuce	False sunflower	(tickseed)	Venice mallow
Blue vervain	Fiddleneck	Plantain	Virginia buttonweed
Bracted plantain	Florida pusley	Poison ivy	Virginia creeper
Brassbuttons	Frenchweed	Poison oak	Virginia pepperweed
Bristly oxtongue	Galinsoga	Pokeweed	Water pennywort
Broadleaf dock	Goathead	Poorjoe	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prairie sunflower	Western clematis
Broomweed	Ground ivy	Prickly lettuce	Western salsify
Buckhorn	Gumweed	Prickly sida	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild buckwheat
Bull thistle	Healall	Prostrate spurge	Wild carrot
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild four-o'clock
Burclover	Heathaster	Puncturevine	Wild garlic
Burdock	Hedge bindweed	Purslane, common	Wild geranium
Burning nettle	Hedge mustard	Ragweed	Wild lettuce
Burweed	Hemp	Red sorrel	Wild marigold
Buttercup	Henbit	Redroot pigweed	Wild mustard
Buttonweed	Hoary cress	Redstem filaree	Wild onion
Canada thistle	Hoary plantain	Rough cinquefoil	Wild parsnip
Carolina geranium	Hoary vervain	Rough fleabane	Wild radish
Carpetweed	Horsenettle	Russian pigweed	Wild rape
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild strawberry
Catnip	Knawel	Scarlet pimpernel	Wild sweet potato
Catsear	Knotweed	Scotch thistle	Wild vetch
Chickweed, common	Kochia	Sheep sorrel	Wild violet
Chickweed, mouseear	Lambsquarters	Shepherdspurse	Woodsorrel
Chicory	Lespedeza	Slender plantain	Woolly croton
Cinquefoil	Mallow	Smallflower galinsoga	Woolly morningglory
Clover, crimson	Matchweed	Smooth dock	Woolly plantain
Clover, hop	Mexicanweed	Smooth pigweed	Wormseed
Clover, red	Milk vetch	Sorrel	Yarrow
Clover, strawberry	Morningglory	Sowthistle	Yellow rocket
Clover, sweet	Mouseear hawkweed	Spanishneedles	Yellowflower peppervieed
Clover, white	Mugwort	Speedwell	and other broadleaf
Cockle	Musk thistle	Spiny amaranth	weeds

WHEN TO USE:

Apply this product when weeds are small and actively growing.

CRABGRASS (large and smooth):

• **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at 14 days later for dense populations.

FOXTAIL (GREEN, GIANT AND YELLOW):

• Spring and early summer: Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at 14 days later for dense populations.

BROADLEAF WEEDS:

- Spring: Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- Fall: Apply when henbit, chickweed, white clover, ground ivy, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

[HOW TO USE]

TRIGGER SPRAYERS:

- 1. Shake well. [if bottle already has sprayer attached only include steps 1 & 2 (not 1a & 1b)]
- 1a. Connect trigger sprayer. Unscrew and discard the white cap on the bottle. Insert the hose into the bottle and tighten the cap securely.
- 1b. Prime Sprayer: Turn the sprayer nozzle one-half turn counter-clockwise. Hold the sprayer lower than the bottle and squeeze the trigger several times until it sprays.
- 2. [Adjust the nozzle to coarse spray pattern to minimize wind drift. (graphic)]. Spray the weeds only. Spray [Apply] to lightly cover broadleaf and grass weeds. Excessive spraying can injure some lawns. Do not apply when daytime temperatures are above 90°F as lawn injury can occur (do not apply above 85°F for bermudagrass).

CULTURAL TIPS:

For newly seeded areas:

• Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas:

• The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the
application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

Delay moving 2 days before and until 2 days after the application of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

SUBLABEL 2

Ready-to-use intended for use by homeowners and marketed by supplemental registrants.

EH-1449 HERBICIDE

NOTIFICATION
JAN 1 4 2014

EPA Reg. No. 2217-906

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	0.313%
Quinclorac	0.104%
Dicamba, dimethylamine salt	0.029%
OTHER INGREDIENTS:	99.554%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 2,4-dichlorophenoxyacetic acid equivalent, 0.260%
- 3,7-dichloro-8-quinolinecarboxylic acid, 0.104%
- 3,6-dichloro-o-anisic acid equivalent, 0.024%

Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

Editorial Note-[Bracketed Text] is optional language

KEEP OUT OF REACH OF CHILDREN

CAUTION

OPEN Re-sealable Label for Directions & Precautions **PRESS TO RE-SEAL**

[EH-1449 Herbicide's exclusive and unique-formula kills both broadleaf plus grassy weeds in one easy step without harming your lawn-Guaranteed!*. Ready-to-use—no mixing, no measuring no messy cleanup. Rainproof in one hour. Controlling weeds including crabgrass in your lawn has never been easier!].



QUICK FACTS	
[Editorial Note – M	larketing Claims Positioned Here]
KILLS	Broadleaf Weeds: Dandelion, Chickweed, and Clover [Over 200+ weeds, including dandelions, chickweeds, clovers.] Grass Weeds: Crabgrass, and over 200 other weeds. [Crabgrass, foxtail and many others.]
WHERE TO USE	ON LAWNS without harming lawn grasses when used as directed (Do not use on bahiagrass, carpetgrass, centipedegrass, St. Augustinegrass [including 'Floratam'], or dichondra.)
COVERAGE	This container treats an average size lawn (X,000 to Y,000 sq.ft.).
☆ □	For questions or comments, call toll-free 1-800-XXX-YYYY or visit our website at www.CompanyName.com

[This product contains: 0.260% 2,4-D acid equivalent, 0.104% quinclorac, and 0.024% dicamba acid equivalent. Isomer specific by AOAC Methods.]

DIRECTIONS	S EOD LISE
	deral law to use this product in a manner inconsistent with its labeling.
HOW TO USE	Turn spray nozzle to coarse spray. Hold sprayer about 12 inches from weed, aim at center of weed, and spray to lightly cover.
WHEN TO APPLY	 Apply when weeds are [young] [small] and actively growing. Apply when the daily high temperature doesn't exceed 90°F (85°F for Bermudagrass). Rain or watering one hour after application will not wash away effectiveness.
WHERE TO APPLY	 To lawns - including: bermudagrass, buffalograss, Kentucky bluegrass, ryegrass, fescue, zoysiagrass and bentgrass. (Application to bermudagrass may cause temporary yellowing or discoloration, but full recovery can be expected) (Do not use on bahiagrass, carpetgrass, centipedegrass, St. Augustinegrass [including 'Floratam'], or dichondra.)
FOR BEST RESULTS	 [Apply when wind is calm to avoid spray drift onto desirable plants (fruit, vegetables, flowers, ground covers, shrubs and trees)]. [Repeat application in 2 weeks if needed for hard-to-control weeds.] [Do not apply while grass is emerging from winter dormancy.] [Newly seeded lawns may be seeded 4 weeks after using this product.] [Bare spots may be seeded 4 weeks after using this product.] [Do not mow for 1 to 2 days after spraying [to allow weed killer to move throughout the weed]. Do not use as compost or mulch]. [If lawn is dry, water 1 to 2 days before using this product. (Weeds take up herbicide best when not under stress.]
REFILL	Add 1.6 fl.oz. (9½ teaspoons) of EH-1432 Herbicide to this empty [24 fl.oz.] container. Top off with water, screw the sprayer back on and shake well. (note: refill does not contain SmartTrack(TM) spray marker.
HOW IT WORKS	EH-1449 Herbicide contains three proven weed killers to control lawn weeds & crabgrass, roots and all. And it won't harm your lawn when used as directed. Weeds begin to wilt and die within [24] hours and are dead within 14 to 28 days.

WEEDS [Kills over 200 weeds including]:			
Black Medic	English Daisy	Pennywort	Wild garlic
Carpetweed	False dandelion	Plantains	Wild geranium
Chickweeds	Florida pusley	Purslane	Wild onion
Clovers	Foxtails	Red sorrel	(& many other broadleaf
Crabgrass	Ground ivy	Sheep sorrel	weeds)
Creeping Charlie	Heartleaf drymary	Spurges	(see appendix for a more
Cudweed	Henbit	Spurweed	complete list of weeds
Curly dock	Kikuyugrass	Thistles	controlled)
Dandelion	Oxalis	Wild carrot	




[alternative people/pet warning statements]

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not allow people (other than applicator) or pets on treatment area during application and until spray has dried.

[optional instructions for use with battery-operated sprayer]

• [Battery-operated trigger sprayer: This product is applied directly from the container using the battery-operated trigger sprayer. The sprayer is packaged with the battery installed. To use, remove the cap from the container, insert the pickup tube and thread cap attached to pickup tube to the container. Move the safety lock located below the trigger to the "ON" position. Pull trigger, waiting several seconds for fluid to prime. Adjust the sprayer tip to deliver coarse spray droplets. To store, move the safety lock located below the trigger to the "LOCK" position.]

IMPORTANT! - READ BEFORE USE




How to Attach Sprayer to Bottle mernove Power Sprayer and hose from side carrier. melace bottle cap with cap attached to Power Sprayer hose.



Prepare to Spray Point Sprayer away from body.

Adjust nozzle to desired setting. Do not turn past 'X" position. Slide safety switch to ON position.



Spray ■ Squeeze trigger to dispense product.
Sprayer may require several seconds to prime before spraying.

After Use

■ Slide safety switch to LOCK position. ■ Return Power Sprayer to side carrier for storage.



4 AA Batteries Included For important battery information – see booklet.

PROPER USE OF BATTERIES

<Drawing of Battery>

- · Never mix old batteries with new.
- Do Not mix Alkaline with Standard (carbon zinc) or Rechargeable (nicker cadmium) batteries.
- When replacing batteries, follow diagram inside battery compartment that is located at the bottom of the sprayer.
- When storing sprayer at end of season, remove batteries to prevent leakage.
- Follow manufacturer's recommendations for disposal of batteries.

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash clothing before reuse.

Applicators and other handlers of this product must wear:

- Long sleeved shirt and long pants
- Shoes
- Socks

First Aid	
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
	alling the poison control center, have this product label accessible. If seeking treatment y room or doctor's office, bring this product label to show medical personnel. You may

call toll-free 1-877-229-3763 for medical emergency information.

Active ingredients: 0.313% 2,4-D, dimethylamine salt (CAS# 2008-39-1), 0.104% quinclorac (CAS# 84087-1-4), and 0.029% dicamba, dimethylamine salt (CAS# 2300-66-5).]

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

Important – Directions for Storage and Disposal Do not contaminate water, food, or feed by storage and disposal.	
Storage Conditions	[Return spray nozzle to closed position.] Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
PESTICIDE DISPOSAL AND CONTAINER HANDLING:	Nonrefillable container. Do not reuse or refill this container If empty: Do not reuse this container [except to refill with EH-1432 Herbicide.] Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

APPENDIX

- 1. Statements which may appear on different label components depending on packaging configuration.
 - · See next panel for additional Precautionary Statements and First Aid
 - Net Contents: ____EPA Est. No. ____
- 2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:
 - Kills [Lawn] Weeds plus Crabgrass in One [Easy] Step
 - All-in-One Weed Solution
 - Kills Crabgrass plus Broadleaf Weeds
 - Two Products in One!
 - Effective [Broadleaf] Weed Protection
 - · Effectively Protects Lawns from Broadleaf Weeds
 - Weed Kill System
 - Kills Listed Weeds
 - Kills [Hard-to-Control] [Tough] Weeds
 - Will not Harm [Your] Lawn[s]
 - · Kills the Weeds, Not the Lawn
 - · All-in-One Lawn Weed Killer
 - Kills to the Root
 - Systemic [Herbicide] [Activity]
 - Kills [Both] Broadleaf and [AND] Grassy Weeds [in One [Easy] Step]
 - · All-in-One Weed Killer for Lawns
 - Gets the Root
 - Dual Action Weed Control Kills the Plant and the Root
 - Kills the Weed, Kills the Root
 - · Kills Weeds not Lawns
 - Kills All Major Broadleaf Weeds
 - Won't Harm Lawns* *When used as directed

- Kills Weeds Won't Harm Lawns
- Two-in-One Product that Kills Crabgrass & Broadleaf Weeds
- · Kills both Crabgrass & Broadleaf weeds
- Double [Dual] [Two-Way] Action Kills Broadleaf and Crabgrass Weeds
- · Kills Both Broadleaf and Crabgrass Weeds
- When You Are Going to Kill Weeds, Why Not Get Them All [Broadleaf & Grass]
- Kills Lawn Weeds Plus Crabgrass
- Two Types of Weeds; One Solution
- Kills Listed Weeds
- · All-In-One Weed Killer
- Kills Even Tough Weeds roots & all
- [Also] Kills [Dandelion], [Crabgrass], and [Clover], Plus [200] Other [Listed] Weeds [As Listed]
- · Controls over 200 Broadleaf Weeds
- · Controls crabgrass and broadleaf weeds in lawns
- · Controls dandelions, crabgrass and other listed hard to kill grassy or broadleaf weeds
- Kills 200+ Weeds
- Kills over 200 Weeds [Plus Crabgrass in One Easy Step]
- Starts Working on Contact
- Unique Formula
- Fast Results
- Kills Crabgrass & Other Weeds in Your Lawn
- · Ideal for Spot Treatment
- Visible Results
- Easy to Use Power Sprayer
- Easy to Use
- No Pumping
- · Battery Powered Sprayer
- Targets Weeds
- Fast Coverage
- Easy & Fast Coverage
- Faster [Coverage]Than Pull & Spray
- Just Point & Shoot
- Kills Weeds Fast
- Save Time & Energy
- Always Ready-To-Spray
- No More Squeeze, Squeeze, Squeeze
- The Easiest Way to Spray
- No Pulling Just Point & Shoot
- Revolutionary [Applicator] [Sprayer]
- Ideal for small areas.
- No messy cleanup.
- Control lawn weeds plus crabgrass in your lawn with ease!
- Rainproof in [1] hour[s]
- SmartTrackTM Marker Shows Treated Weeds
- Contains SmartTrackTM, a Non Staining Marker [that Indicates Where You've Sprayed]
- Guaranteed* Not to Harm Your Lawn
- Guaranteed* Not to Harm Lawns
- Guaranteed* Results
- Kills Weeds the First Time, Every Time, Guaranteed*
- 100% Satisfaction Guaranteed* (or your money back)
- Also Kills Dandilion and Clover Plus 200 Other Listed Weeds
- Kills crabgrass and over XXX listed weeds in country and suburban lawns* *See inside booklet for a complete list of weeds controlled

Broadleaf Weeds: (a partial list is presented below)

Annual Yellow Sweetclover, Aster, Austrian Fieldcress, Barnyardgrass, Bedstraw, Beggarticks, Florida Betony, Field Bindweed, Bird Vetch, Bitter Wintercress, Hairy Bittercress, Bitterweed, Black-eyed Susan, Black Medic, Black Mustard, Blackseed Plantain, Blessed Thistle, Bloodflower Milkweed, Blue Lettuce, Blue Vervain, Bracted Plantain, Brassbuttons, Bristly Oxtongue, Broadleaf Dock, Broadleaf Plantain, Broadleaf Signalgrass, Broomweed, Buckhorn Plantain, Bulbous Buttercup, Bull Thistle, Bullnettle, Burclover, Burdock, Burning Nettle, Burweed, Buttercup, Buttonweed, Canada Thistle, Carolina Geranium, Carpetweed, Catchweed Bedstraw, Catnip, Catsear, Common Chickweed, Mouseear Chickweed, Chicory, Cinquefoil, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White Clover, Cockle, Cocklebur, Common Mullein, Creeping Charlie, Creeping Jenny, Cudweed, Curly Dock, English Daisy, Fleabane Daisy, Oxeye Daisy, Dandelion, Dichondra, Dogbane, Dogfennel, Dollarweed, Elderberry, False Dandelion, False Flax, False Sunflower, Fiddleneck, Florida Pusley, Frenchweed, Galinsoga, Giant Foxtail, Goathead, Goldenrod, Green Foxtail, Ground Ivy, Gumweed, Hairy Fleabane, Hawkweed, Healall, Drymary Heartleaf, Heath Aster, Hedge Bindweed, Hedge Mustard, Hemp, Henbit, Hoary Cress, Hoary Plantain, Hoary Veryain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Large Crabgrass, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk Vetch. Morningglory, Mouseear Hawkweed, Mugwort, Musk Thistle, Mustard, Narrowleaf Plantain, Narrowleaf Vetch, Nettle, Orange Hawkweed, Oriental Cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods Bedstraw, Plains Coreopsis (Tickseed), Plantain, Poison Ivy, Poison Oak, Pokeweed, Poorjoe, Prairie Sunflower, Prickly Lettuce, Prickly Sida, Prostrate Knotweed, Prostrate Pigweed, Prostrate Spurge, Prostrate Vervain, Puncturevine, Common Purslane, Ragweed, Red Sorrel, Redroot Pigweed, Redstem Filaree, Rough Cinquefoil, Rough Fleabane, Russian Pigweed, Russian Thistle, Scarlet Pimpernel, Scotch Thistle, Sheep Sorrel, Shepherdspurse, Slender Plantain, Smallflower Galinsoga, Smooth Crabgrass, Smooth Dock, Smooth Pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny Amaranth, Spiny Cocklebur, Spiny Sowthistle, Spotted Catsear, Spotted Spurge, Spurged, Stinging Nettle, India Mock Strawberry, Tall Nettle, Tall Vervain, Tansy Mustard, Tansy Ragwort, Tanweed, Thistle, Trailing Crownvetch, Tumble Mustard, Tumble Pigweed, Velvetleaf, Venice Mallow, Virginia Buttonweed, Virginia Creeper, Virginia Pepperweed, Wavyleaf Bullthistle, Western Clematis, Western Salsify, White Mustard, Wild Aster, Wild Buckwheat, Wild Carrot, Wild Four-o'clock, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Marigold, Wild Mustard, Wild Onion, Wild Parsnip, Wild Radish, Wild Rape, Wild Strawberry, Wild Sweet Potato, Wild Vetch, Wild Violet, Woodsorrel, Woolly Croton, Woolly Morningglory, Woolly Plantain, Wormseed, Yarrow, Yellow Foxtail, Yellowflower Pepperweed, Yellow Rocket, and other broadleaf weeds

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00906.20130627.notif-proposed-highlighted.doc

2. Reason for Issue: Re-format label, ad claims, ABN